



76. 34 80

CLINICAL STUDIES
OF
DISEASE IN CHILDREN

BY
EUSTACE SMITH, M.D.

FELLOW OF THE ROYAL COLLEGE OF PHYSICIANS
PHYSICIAN TO HIS MAJESTY THE KING OF THE BELGIANS
PHYSICIAN TO THE EAST LONDON CHILDREN'S HOSPITAL
AND TO THE VICTORIA PARK HOSPITAL FOR DISEASES OF THE CHEST

SECOND EDITION



LONDON
J. & A. CHURCHILL
11, NEW BURLINGTON STREET
1887

[*All Rights reserved*]

PREFACE TO THE SECOND EDITION.

THIS little work has been out of print now for nearly five years, but the pressure of other duties has hitherto interfered with its revision.

In the present edition many alterations have been introduced. In parts these have been so extensive as to amount to a re-writing of portions of the book. This may be said especially of Chapters III., IV., and X., which will be found to differ greatly from the first edition.

It was originally the Author's intention to increase the size of the work in successive issues, so that it might come in time to include most of the diseases to which children are liable. On reflection, however, it seemed more convenient to deal with the whole subject of Disease in Children in an independent work.

Accordingly, the scope of the present volume has not been altered, and the only changes made have been such as an enlarged clinical experience has shown to be required.

GEORGE STREET, HANOVER SQUARE,
February, 1887.

CONTENTS.

CHAPTER I.

INTRODUCTION.

	PAGE
Peeuliarities of eonstitution in early life	1
Children not exempt from degeneration of tissue	2
Exeitability of the nervous system in the child	2
Wide diffusion of symptoms in aeute disease	3
Intensity of symptoms no guide to their eause	3
General and loeal symptoms	3
Loeal symptoms sometimes deecitful	3
Frequeney of funetional derangements	4
Effect of ehronic disease on nervous exeitability	5
Clinical examination of young ehildren	6
Value of the mother's aeount of the illness	6
Method of taking the temperature	7
Respiration in infants	8
Examination of ehest	9
Examination of mouth	11
General remarks on the diagnosis of disease	11
General remarks on prognosis	14
Treatment of disease in ehildren	14
Objects of treatment	15
Indications for treatment	15
Evils of too early stimulation	16
Value of alkalis	17
Tolerance of ehildren for eertain drugs	18
Doses	18

CHAPTER II.

COLLAPSE OF THE LUNG—(POST-NATAL ATELECTASIS).

Collapse of lung, a eommon lesion in the child	21
Method of production	21

	PAGE
<i>Causes</i>	22
Inability to cough and expectorate	22
Weakness of the inspiratory act	22
Mechanical impediment to expansion of the chest	23
Influence of rickets	23
Diseases which are apt to be complicated by atelectasis	24
<i>Morbid anatomy</i>	25
<i>Symptoms</i>	26
If collapse extensive may cause sudden death	26
Convulsions	26
Insufficient aëration of blood	26
Difficulty of swallowing	27
<i>Physical signs</i>	27
Sometimes absent	28
<i>Case I.</i> —Sudden death from pulmonary collapse in a weakly, wasted infant	28
<i>Case II.</i> —Collapse of the lung occurring in a case of pertussis	29
Comment	31
Mechanism of collapse in pertussis	31
<i>Case III.</i> —Collapse of the lung with laryngismus stridulus	32
<i>Case IV.</i> —Bronchitis and pulmonary collapse in a rickety child	33
Comment	36
<i>Diagnosis</i> of pulmonary collapse	37
of diffused collapse	38
of lobular collapse	39
<i>Prognosis</i>	40
<i>Treatment</i>	41
Mechanical treatment	41
Emetics	41
Stimulation	42

CHAPTER III.

CROUPOUS PNEUMONIA.

Not uncommon under two years	43
Often accompanied by signs of nervous disturbance	44
<i>Symptoms</i>	44
Temperature	44
Cough	44
Pulse-respiration ratio	45
Breathing rapid but not laborious	45
Prostration	45

CONTENTS.

vii

	PAGE
<i>Physical signs</i>	46
Seldom complete before the third day	46
Importance of using two fingers in percussion	46
Sequence of physical signs	47
Situation of physical signs	47
Resolution of the inflammation	48
Sudden fall in the temperature	49
Date of the crisis	49
Mode of beginning with convulsions	49
<i>Case V.</i> —Pneumonia introduced by a convulsive attack	50
Other nervous symptoms	51
<i>Case VI.</i> —Pneumonia of apex with severe cerebral symptoms	52
Comment	55
Value of irregularity of pulse as a symptom	55
<i>Case VII.</i> —Rapid course of pneumonia of apex	56
<i>Case VIII.</i> —Pneumonia complicated by gastro-intestinal catarrh	57
Comment	59
Diagnosis from typhoid fever	59
Danger of this complication in babies	60
<i>Case IX.</i> —Pneumonia of left lung in an infant : gastro-intestinal catarrh	60
Comment	61
Postponement of resolution	62
Relapses	62
<i>Prognosis</i>	63
Slight gravity of nervous symptoms in the primary disease	63
<i>Diagnosis</i>	64
<i>Treatment</i>	67
Poultices	67
Apcrients	68
Diet	68
Reduction of heat	69
Quinine	70
Treatment of secondary pneumonia	71

CHAPTER IV.

PLEURISY.

Physical signs very similar to those of pneumonia	72
Usually limited to one side of the chest	73
<i>Symptoms</i>	73
Pain	74
Respiration	74

	PAGE
Pulse	74
Pulse-respiration ratio	74
Digestive derangement	74
Cough	75
Temperature	75
Complexion	76
Degree of prostration	76
<i>Physical signs</i>	76
Displacement of organs	77
Dilatation of the side	77
Alteration of shape of chest	77
Vocal fremitus	78
Dulness, its quality and distribution	78
Movability of dulness	78
Auscultation	79
Friction	79
Breath-sounds	79
<i>Case X.—Ordinary attack of pleurisy</i>	81
Comment	84
Localised pleurisy	85
<i>Case XI.—Localised pleurisy</i>	85
Comment	87
Empyema	89
<i>Case XII.—Empyema—absorption of fluid</i>	89
Comment	92
Empyema—symptoms	92
Its consequences	93
<i>Case XIII.—Empyema—spontaneous evacuation of the fluid</i>	94
Source of danger in spontaneous evacuation	96
Retraction of chest	97
<i>Case XIV.—Great retraction of the chest from empyema</i>	97
Comment	98
<i>Case XV.—Empyema—Secondary acute tuberculosis</i>	98
Comment	102
Fibrinous pleurisy	104
Extension of inflammation from the pleura to the peritoneum	104
<i>Case XVI.—Empyema—Extension of inflammation to the peri-</i> <i>toneum</i>	105
Comment	107
<i>Diagnosis</i>	108
From pneumonia	108
From fibroid induration with secondary pneumonia	109
Of localised pleurisy	112
Of empyema	113

CONTENTS.

x

	PAGE
From phthisis	113
From enlarged liver	114
From hydrothorax	115
<i>Prognosis</i>	115
<i>Treatment</i>	116
Hypodermic injection of morphia	117
Counter-irritants	118
Dry diet	118
Paracentesis	119
Use of aspirator	120
<i>Case XVII.</i> —Traumatic pneumo-thorax from puncture of the lung during paracentesis	121
Use of exploring needle	122
Causes of failure to find fluid	122
Use of drainage-tube	123
Correction of fœtor of pus	124
Prevention of deformity of chest	124
Diet in empyema	125
Secondary pleurisy	125

CHAPTER V.

ACUTE CATARRHAL PNEUMONIA.

Especially the pneumonia of infancy	126
Distinction between croupous and catarrhal pneumonia	127
Pathology	127
Modes of origin	128
Extension of catarrhal process	128
From collapse of lung	128
Always a secondary lesion	129
Symptoms and course	130
Temperature	130
Pulse-respiration ratio	130
Cough	130
<i>Physical signs</i>	131
<i>Case XVIII.</i> —Ordinary case of acute catarrhal pneumonia in an infant	131
<i>Case XIX.</i> —Acute catarrhal pneumonia secondary to collapse of the lung	132
Comment	134
Apathy of the patient	135
Subacute course of the disease	135

	PAGE
<i>Case XX.—Subacute type of catarrhal pneumonia</i>	135
Comment	137
Favourable termination	139
<i>Case XXI.—Favourable ending of acute catarrhal pneumonia</i> .	140
Comment	142
Dilatation of bronchi	143
<i>Case XXII.—Acute catarrhal pneumonia with dilatation of</i> bronchus	143
Comment	144
Catarrhal pneumonia complicating tuberculous bronchitis . .	145
<i>Diagnosis</i>	146
From lobular collapse of lung	147
From croupous pneumonia	147
<i>Prognosis</i>	148
Unfavourable symptoms	149
Favourable symptoms	149
<i>Treatment</i>	149
Of pulmonary catarrh	149
Counter-irritants	150
Expectorants	150
Danger of too early stimulation	150
Of catarrhal pneumonia	152
Diet	152
Danger of over-feeding	153

CHAPTER VI.

CHRONIC CATARRHAL PNEUMONIA AND UNABSORBED PNEUMONIC DEPOSITS—PNEUMONIC PHTHISIS.

The principal form of chronic lung disease	154
<i>Pathology</i>	155
Origin of caseous masses	156
Process of chronic catarrhal pneumonia	156
Caseation of pulmonary deposits	156
“Tubercular infiltration” of Laennec	156
Softening	157
Excavation	158
Calcification	158
Fibroid induration	159
Secondary changes	159
Ulceration of bowels	159
Secondary Tuberculosis	159

	PAGE
<i>Symptoms</i>	160
Conformation of body	160
Peculiarities of chronic lung disease in children	160
Sources of error in diagnosis	161
General symptoms mild	162
Physical signs	163
Temperature	163
<i>Case XXIII.—Chronic catarrhal pneumonia. Rapid absorption</i>	
of the deposit	164
Comment	165
Seat of consolidation	165
Symptoms of softening	165
Signs of excavation	167
Diagnosis of cavity from a dilated bronchus	167
Imperfect evidence of cavity	167
Cavernous breathing without a cavity	168
General symptoms of softening	168
Rarity of hæmoptysis	168
<i>Case XXIV.—Chronic catarrhal pneumonia.—Softening and</i>	
excavation	169
Comment	170
<i>Case XXV.—Caseous consolidation with dilatation of bronchus</i>	173
Comment	176
Absorption of pneumonic deposits	178
Secondary tuberculosis	178
<i>Case XXVI.—Chronic catarrhal pneumonia, with plastic</i>	
pleurisy and secondary tuberculosis	178
Comment	181
<i>Diagnosis</i>	183
In the early stage	184
From acute tuberculous	185
In the advanced stage	185
From empyema	185
<i>Prognosis</i>	186
<i>Treatment</i>	187
Dry air and soil	188
Diet	188
Medicines	189

CHAPTER VII.

FIBROID INDURATION OF THE LUNG (CIRRHOSIS).

<i>Pathology</i>	192
----------------------------	-----

	PAGE
A result of catarrhal pneumonia	193
of pleurisy	194
Fibroid phthisis	194
<i>Case XXVII.</i> —Fibroid thickening of lung following an attack of catarrhal pneumonia	194
Comment	202
Physical signs	205
Symptoms	205
Fetid expectoration	205
State of nutrition	206
Absence of pyrexia	206
Interference with pulmonary circulation	206
Edema	207
<i>Case XXVIII.</i> —Fibroid induration of apex of lung with con- traction of chest	207
<i>Case XXIX.</i> —Fibroid induration of apex of lung with contrac- tion	209
Comment	210
<i>Case XXX.</i> —Extreme fibroid induration and contraction	212
Comment	214
Diagnosis from pleurisy with retraction	215
<i>Case XXXI.</i> —Extreme fibroid induration without contraction of side	216
Comment	224
Excavation of lung	225
<i>Case XXXII.</i> —Fibroid induration with extensive excavation of lung (Fibroid Phthisis)	227
Comment	231
<i>Diagnosis</i>	232
<i>Prognosis</i>	233
<i>Treatment</i>	235
Avoidance of chills	235
Inhalations	235
Tonics and astringents	235
Emetics	236

CHAPTER VIII.

ACUTE GENERAL TUBERCULOSIS.

Not to be confounded with pulmonary phthisis	237
Resemblance to the acute specific diseases	238
Often occurs in apparently healthy children	238

CONTENTS.

xi

	PAGE
Pathology	238
Often a result of secondary infection	239
Sometimes a primary disease	239
Causation	240
Seat and distribution	241
<i>Symptoms</i>	242
General	242
Wasting	243
Temperature	243
Fever often remittent	243
Loss of elasticity of skin	244
Albuminuria	244
Edema	244
Mode of onset	245
Special	245
Pulmonary tuberculosis	245
<i>Symptoms</i>	245
Cough	245
Breathing	246
Secondary diseases	246
Bronchitis	246
Catarrhal pneumonia	247
Cerebral complications	247
<i>Case XXXIII.—Acute pulmonary tuberculosis with tubercular peritonitis.</i>	247
Comment	250
<i>Case XXXIV.—Acute general tuberculosis following old-standing disease of bone.</i>	251
Comment	253
Diagnosis of acute general tuberculosis	254
In infants	254
In older children	256
From typhoid fever	256
of acute pulmonary tuberculosis	257
of tuberculous bronchitis	257
of tuberculous broncho-pneumonia	258

CHAPTER IX.

ACUTE TUBERCULOSIS OF THE PIA MATER.—TUBERCULAR MENINGITIS.—ACUTE HYDROCEPHALUS.

Two forms of the disease	259
Secondary form	259
Primary form	259

	PAGE
Symptoms of the primary form	260
Premonitory symptoms	260
Stages of the disease	261
Special symptoms	261
Symptoms of first stage	261
Symptoms of second stage	262
Pulse	262
Headache	262
Respiration	263
Pupils	263
Cerebral flush	263
Symptoms of third stage	263
Stupor	263
Paralytic lesions	264
Convulsions	265
State of the fontanelle	265
Variations in the symptoms	266
In state of bowels	266
Pulse	266
Temperature	266
State of circulation	267
Duration	267
Case XXXV.—Ordinary attack of primary acute hydrocephalus	267
Comment	271
The secondary form	271
Case XXXVI.—Secondary tubercular meningitis	271
Comment	273
Case XXXVII.—Secondary tubercular meningitis	274
Comment	275
Seat of the meningitis	276
Usually at the base	276
Sometimes at the convexity	276
Case XXXVIII.—Tubercular meningitis occupying the convexity	276
Comment	278
Diagnosis	279
Value of special symptoms	280
Relative value of nervous symptoms	281
Diagnosis from typhoid fever	282
from exhaustion	283
from wasting with digestive derangement	283
from simple meningitis	283
cerebral tumour	284
cerebral abscess	284
Prognosis	285

CHAPTER X.

TUBERCULOSIS OF THE PERITONEUM.—TUBERCULAR PERITONITIS.

	PAGE
Course less acute than in the other forms of tuberculous disease	286
Early symptoms very obscure	287
<i>Symptoms</i>	288
Pain and tenderness	288
Temperature	288
Abdominal swelling	288
Fluctuation	289
Increased sense of resistance	289
Shape of abdomen	290
Diarrhœa	291
<i>Case XXXIX.—Tubercular peritonitis</i>	292
Comment	295
<i>Case XL.—Tubercular peritonitis</i>	296
Comment	298
<i>Diagnosis</i>	299
From abdominal derangement	299
Case of common functional derangement	300
Value of tenderness as a symptom	301
From ulceration of bowels	302
Treatment of acute general tuberculosis	302
Preventive treatment	303
When disease established	304
Diarrhœa	304
Inflammatory chest affections	305
Tubercular meningitis	305
Tubercular peritonitis	306

CLINICAL STUDIES

OF

DISEASE IN CHILDREN.

CHAPTER I.

INTRODUCTION.

DISEASE in early life often rendered obscure by the peculiarities of the childish constitution—Intense vital activity of organs and excitability of the nervous system—Wide diffusion of symptoms in cases of acute disease—Intensity of symptoms no guide to their cause—Importance of local symptoms—But these sometimes misleading.

Frequency of functional derangements—Loss of normal nervous excitability under the influence of chronic disease.

Clinical examination of young children—Method of proceeding—Order of examination—Pulse—Temperature—Examination of chest.

General remarks on diagnosis, prognosis and treatment—Indications for treatment—Evils of too early stimulation—Drugs—Tolerance of children for certain drugs and susceptibility to others—Doses.

THE diseases of infancy and childhood are often regarded as presenting peculiar difficulties ; and certainly, to one accustomed only to disease as it presents itself in the adult subject, the symptoms will frequently appear obscure and sometimes contradictory. Children differ from adults in many ways. The intense nervous excitability in a robust child will often communicate a false appearance of gravity to a trivial ailment, while the lowered nervous sensibility which is found in a child much depressed by chronic disease will so obscure the symptoms that a serious intercurrent

malady may excite little notice or even be completely overlooked.

The diseases of children are in some respects simpler, in others more obscure, than those of older persons. In the first place, children are less subject than adults to degenerations of tissue. They are not, however, exempt from such degenerations, for all the consequences of impaired nutrition may be found in the child, as in the adult ; and not only may morbid products undergo a degenerative process, but also the normal tissues of the body may become the seat of structural changes such as in older persons are ascribed to the influence of advancing years. Thus, commencing atheroma of the aorta may be found in children of two or three years old ; amyloid degenerations are far from uncommon in early life ; and the degenerative processes occurring in pulmonary morbid deposits may be present in the youngest infants.

Still, as compared with the adult, degenerations are rare in the child ; growth and development are in active progress, and the organs are gathering strength and maturing themselves for the efficient performance of their duties. The intense vital activity of organs, to which the child owes this comparative immunity, increases his susceptibility to disease by rendering him more liable to be affected by slight causes ; it also renders the course of disease more rapid, the attack more sudden, and the symptoms more severe. Moreover, the rapid development through which the whole nervous system is passing seems to link together the various organs into a closer sympathy, so that disease of one of these will often cause a general commotion throughout the body, and the most prominent symptoms may emanate from some distant organ, and not from the part which is the actual seat of trouble.

The access of acute disease in the young child is always accompanied by signs of general disturbance ; and it is very important in the examination to distinguish between

the general and the local symptoms—for the latter, although sometimes obscure, are yet much more exact than are those which arise from the general perturbation of the system. Thus, in the beginning of an attack of pneumonia, the cough, the rapid breathing, and the indications of pain in the side point pretty clearly to the lungs as the part attacked ; while the high temperature, the vomiting, the headache, the rapid pulse, and in many cases the convulsions, are not special symptoms of chest disease, but are common to the beginning of many other febrile affections.

In entering upon the study of children's diseases it is very important to bear in mind this wide diffusion of symptoms, and also to remember that their severity bears no relation at all to the severity of the local disorder of which they are the manifestation. A mere functional disturbance will often, for the time, excite as wild a tumult in the system as the severest organic lesion ; and we often find it necessary in diagnosis to wait for the general symptoms to diminish before we are able to gather from the local symptoms any sufficient tokens of the cause and nature of the ailment. Local symptoms are not, however, always present. In infants, the weakness of the organism is so great, and the hold upon life so feeble, that death may take place before the more special symptoms have had time to reveal themselves ; and thus a child thrown into violent convulsions by some trifling disorder may actually die from the shock to the system, and no post-mortem appearances may be afterwards found to account for the illness.

Even local symptoms themselves are not always to be trusted. The organ from which the more decided symptoms appear to arise may not be, as has been already stated, the organ in which the morbid process is actually seated, and these deceptive manifestations may throw the real lesion completely into the shade. Thus, in some cases of pneumonia the more prominent symptoms may be those of cerebral disturbance ; violent convulsions, delirium and

stupor may so disguise the real nature of the complaint that the slight cough may be either unnoticed or unregarded, and active measures may be taken for the suppression of a supposed meningitis, to the neglect of the organ which is the real seat of disease.

Again, the local symptoms, although arising from the actual seat of trouble, may convey by themselves no trustworthy indication of the extent or even the nature of the mischief. A child who went to bed healthy, except for a slight cold, wakes up suddenly after a few hours' sleep with fever, stridulous breathing, and a hoarse, barking cough. His breathing is very difficult, and sometimes becomes completely arrested, and we see the child with head thrown back, open mouth, and nostrils widely expanded, making ineffectual efforts to draw a breath, while his staring eyes, his frightened look, and the lividity of his face prove the reality of the impediment to the entrance of air. The spasm after a time relaxes, the symptoms subside, and the child falls asleep. In the morning he may appear as well as on the eve of the attack, or may have a slight hoarseness and some cough, which subside after a few days. In this example the real morbid condition is merely catarrh of the laryngeal mucous membrane, while all that is alarming in the symptoms is produced by functional spasm from excentric irritation: it is an instance of the important part played by mere disorder of function in the diseases of childhood. This frequency of mere functional derangements is another peculiarity of early life which it is important to be aware of. On account of the extreme impressionability of the nervous system in young children, and the delicacy of their organs—especially their singular sensitiveness to changes of temperature—such disturbances form a large proportion of their ailments, and it is of the utmost importance for both prognosis and treatment that we should be able to form an accurate estimate of the real nature of such complaints. In this way gastric, intestinal and pulmonary

catarrhs arise, and a violent attack of vomiting or diarrhœa will soon reduce an infant to a very dangerous state of weakness. Even in cases where the derangement is less acute and severe it may still be attended by actual danger to life. An infant is so dependent, not only for growth and development, but for actual existence, upon a frequent supply of nourishment, that any derangement which temporarily checks the assimilation of nutritive material is an event always to be regarded with anxiety. Such a condition is not an uncommon sequence to severe operations, and often causes death, even after the more immediate effects of the operation have been recovered from. In the case of pulmonary catarrh there is a special danger which is independent of the severity of the catarrh. Collapse of the lung is always threatened in every case of cold on the chest in a weakly infant; and if the collapse be extensive and quickly produced, the child may die suddenly. It is in consequence of this dangerous character of mere functional disorders that post-mortem examinations in young children are so often unsatisfactory. Nothing at all may be found to indicate the cause of death, or the appearances observed may be quite insufficient in themselves to account for the fatal result.

The intense excitability of the nervous system, which under ordinary circumstances impresses their peculiar character upon the disorders of early life, gradually diminishes and disappears under the influence of chronic enfeebling disease, and in its stead we find a condition exactly the opposite to that which has been described. Here, the nervous system appears nearly insensible to all sources of irritation; the general symptoms are almost completely absent, and the local symptoms are so slight and obscure that they may easily escape notice. Thus, inflammation of the lung occurring secondarily in a cachectic child may present hardly any symptoms at all. The temperature may be only slightly elevated, and quickly fall: the cough is

rare or absent ; and an increase in the general weakness is the principal sign by which this serious complication is manifested.

In the child, then, we must be prepared for these two very opposite conditions : in the one case, a violent reaction of the whole system against the most trifling disturbance ; in the other, a profound insensibility to nervous impressions, the system assuming for the time many of the physiological characters of old age, and responding faintly, if at all, to the stimulus of local irritations.

The clinical examination of infants is seldom difficult, for very young children as a rule can be easily managed with a little patience. Older children, however, are often very suspicious of "the doctor," and sometimes, especially if badly brought up and petted and spoiled, may resist all our efforts to examine them ; but even in these cases much may be done by gentleness and deliberation. It is important not to be in a hurry ; and, besides, it is well to accustom the patient to our presence in the room before proceeding to a physical examination. This interval may be conveniently occupied by inquiring into the history of the attack ; and, if there be other children of the same family, it is advisable to see them, for information may in this way be gained as to constitutional and hereditary family peculiarities. The information to be derived from the mother is often of the greatest value, and her account of the symptoms should be always received with attention and carefully weighed. If she be a person of intelligence, many minute points in the history of the beginning of the attack can thus be collected ; and these, in an obscure case, may be of the utmost importance. If the patient be an infant, it is advisable to see him first during sleep, for those parts of the examination during which perfect quiet is indispensable can seldom be satisfactorily conducted unless the child is unconscious of what is passing around him. At this time,

then, we can note the expression of the face, can observe the attitude and any involuntary movements of the body ; we can count the number of respirations, feel the pulse, and ascertain the temperature of the body by gently introducing the ball of the thermometer, wetted or oiled, into the rectum. It is better to take the temperature in the rectum rather than in the axilla ; for, besides that the result thus obtained is more to be relied upon, as the skin of infants is quickly cooled by slight exposure, and does not therefore always furnish trustworthy information as to the heat of the body generally, the method is also more convenient. Infants usually resent very strongly the amount of constraint necessary to keep the arm firmly in contact with the side, and are often so noisy and unmanageable in consequence that little afterwards can be done with them in the way of examination. With regard to the temperature of children, it may be noted that we must not allow ourselves to be deceived by sudden and rapid rises of temperature into the belief that the patient must necessarily be suffering from serious disease. Very slight causes will in infants produce a remarkable increase of heat ; and during natural dentition, just before the passage of the tooth through the gum, a temperature of 104° or 105° Fahr., even in the morning, is not at all an uncommon circumstance. Besides, the normal temperature of young children is rather higher than that of the adult. In a perfectly healthy child of three or four years old the thermometer will often register a morning temperature of $99\frac{1}{2}^{\circ}$.

The pulse of infants can seldom be counted except during sleep, and even then involuntary movements of the arm and twitching of the tendons of the wrist render the operation not always an easy one. The finger should be laid very lightly upon the artery, and any temporary restlessness which may be excited in the limb by the touch should be allowed to subside before we begin to take the number of pulsations. The pulse is rather slower during sleep, and is

then often irregular in force ; but the slightest movement of the body causes a sudden quickening in the number of beats, and at the same time the force of the beats becomes equalised, and remains so for a few seconds, or as long as the increase in rapidity is maintained.

Whatever may be the disease for which we are called to see an infant, we should always make a thorough examination of the whole body ; and therefore, when we have satisfied ourselves as to the points which can only be successfully noted during sleep, it is well to have the clothes entirely removed, so that we may have the whole body exposed to view. If the child cry during this process, we should pay attention to the quality of the voice, for hoarseness is one of the early signs of inherited syphilis.

Placing the infant naked on his nurse's lap, the body must be examined for signs of eruption or any appearance of swelling or local inflammation ; the condition of the skin must also be noted—as to softness and suppleness, dryness, or moisture—for a harsh, dry skin is a frequent accompaniment of chronic abdominal disease. At the same time the condition of the belly can be ascertained, paying attention to the hardness or softness of the parietes, and to the size, whether normal or otherwise, of the liver and spleen. The existence of abdominal pain or tenderness can also be detected without much difficulty.

Children breathe more quickly than adults. Under the age of two years the number of respirations in the minute may be taken at about thirty. After that age the rapidity gradually diminishes ; but at any moment in young subjects the respiratory movement is apt, like the pulse, to be temporarily quickened by trifling causes.

The respiration of infants is chiefly abdominal, on account of the shape of the thorax. In a new-born child the antero-posterior diameter of the chest is scarcely less than the lateral, and this nearly circular shape allows of little further lateral expansion. The circular shape, however,

soon changes to the elliptical, and the chest becomes broad out of proportion to its depth. In a perfectly healthy child there is at each inspiration a certain recession of the lower part of the chest-walls, for the yielding ribs are forced inwards by the pressure of the air before they can be sufficiently supported by the expanding lungs. Disease, however, may greatly increase this recession, and in certain cases we notice a deep furrow appearing across the chest at each inspiration, and marking the upper borders of the abdominal viscera. The extent of the recession is a sign of the degree of difficulty with which the air enters the lungs, and should always be noted during inspection of the chest. The impediment may be created by softening of the ribs, as in rickets, which renders difficult the proper expansion of the thoracic cavity, or by disease of the lungs and air-passages, which presents a direct obstacle to the entrance of air.

The examination of the chest in an infant is not attended with any great difficulty. In severe chest-complaints the child is almost always quiet, as crying interferes with the respiratory act, upon which his whole attention is concentrated. Much noise and resistance, then, may be taken as a sign that the lungs are not seriously affected. If a child with any abnormal dulness at the base of the chest cry loudly, the disease is almost certain to be pleurisy.

The infant should be first placed in the nurse's arms, with his chin resting upon her left shoulder, and his left arm round her neck. If necessary, the chin can be kept down by the nurse's hand placed upon the child's head. In this position the muscles of both shoulders are relaxed, and percussion of the supra-spinous fossæ can be relied upon. As the resonance differs considerably, according as to whether percussion be made during inspiration or expiration, it is important that this should be made on each side at the same period of the respiratory movement. Infants are fond, when under constraint of any kind, of holding

their breath in the intervals of crying ; and as they always do so at the end of an inspiration, this is a favourable opportunity for employing percussio of the two sides. Percussio should always be *mediate*, and it is by far the most convenient to use the first finger of the left hand as a pleximeter. It is best to use two fingers, as well as one, in striking : for by this means, without using any greater force, a larger volume of sound is elicited, and an amount of dulness which might escape notice were only one finger employed, can by this means be readily detected.

It must be remembered that during expiration the liver rises higher in children than it does in adults, and that therefore the right base is naturally less resonant than the left immediately before a fresh breath is taken.

In auscultation the stethoscope should always be made use of. It has the great advantage of limiting the area within which the different respiratory sounds are perceived ; and this, when, as in a child, the area is small, and sounds from the nose and throat are so rapidly conducted, is a matter of considerable importance. If not seriously ill, the child often cries during the examination ; but this will not prevent our acquiring all necessary information, for during inspiration the cry usually ceases, and in this interval of silence we can ascertain the quality of the breathing, and the presence or absence of adventitious sounds. Afterwards we notice whether or not the cry is unnaturally resonant.

To examine the front and sides of the chest the infant should be laid upon a pillow ; and it is convenient to have this latter placed upon the table so as to raise up the patient to a convenient height. By this means the examination can be made thoroughly and without that inconvenience which the practitioner experiences when he stoops down to examine a child lying upon his mother's lap. After exploring the lungs, the heart-sounds can be listened to during inspiration in the intervals of the cry.

In older children an examination of the chest should never

be omitted, whatever may be the nature of the symptoms; for inflammation of the lung often begins very insidiously, and the symptoms of pleurisy are frequently obscure.

The respiratory sounds are of a more blowing quality in children than they are in adults, and this quality is more remarkable in inspiration than it is in expiration, which latter, indeed, is often very weak, and sometimes hardly perceptible. In cases of inflammatory affections of the lungs the rhythm of respiration may become altered, the pause taking place at the end of inspiration instead of at the end of expiration. This is not characteristic of any special disease, but is common to all cases where the function of the lungs is interfered with. During examination of the chest the patient's mouth should, if possible, be open, as by this means the transmission of sounds generated in the throat is to a considerable extent prevented.

The examination of the mouth should be reserved for the end of the investigation, as it often excites very great resistance on the part of the patient. By placing the finger upon the child's chin, and using a little gentle pressure, the mouth opens and the tongue is often protruded at the same time; if not, we can with a pencil-case press down the lower jaw and tongue, and thus obtain a very good view. Afterwards we should run the forefinger along the gums, and ascertain their condition as to heat and swelling. If there be fever, the cause of which our previous examination has left undetected, the throat should always be inspected, depressing the tongue with the handle of a spoon for the purpose. Lastly, if vomiting or diarrhoea be present, the discharges should be looked at: in diarrhoea especially this is of importance, as the method of treatment to be adopted will depend very much upon the exact character of the stools.

The *diagnosis* of disease is not more difficult in the child than it is in the adult; in some respects it is easier, for although the infant cannot himself describe his symptoms, we have in the mother a ready recorder of all changes visible

to the eye, while the facility with which the whole body of the patient can be examined leaves us little excuse if, from want of due diligence, any affection appreciable by the eye, the ear, or the finger be passed over. In all cases of acute disease, before the removal of the clothes, we should be careful to take the temperature with a thermometer, and to count the respirations, and, if possible, the pulse. The respirations especially should not be neglected, for, if the breathing be not laboured, mere rapidity of breathing may easily be overlooked unless special attention be directed to it.

It is at the beginning of the attack that diagnosis is especially difficult. In determining the nature of the complaint at this time we must take into account the peculiarities of children, and, remembering the nervous excitability of young subjects, must not allow our attention to be diverted by the signs of general excitement from noting the more special symptoms which point to local distress. In all cases we should endeavour to obtain from the mother a clear account of the history of the attack, so as to be able, if possible, to fix a definite time for its beginning. A careful examination should then be made of the whole body, and by this means a diagnosis can often be arrived at immediately.

One point of which it is always necessary to take note is the general expression of the child. In a young subject a haggard face is almost invariably an indication of physical distress ; seldom or never of mental worry. If then a child "looks ill," that is to say, has a worn, haggard expression, we may be sure that he is really ill, and however apparently trifling his symptoms may be, should carefully abstain from speaking lightly of his condition. In some serious diseases, such as tubercular meningitis, the early symptoms may be insignificant ; but when we look at the patient's face and see his features pinched and drawn, we may be sure that it will not be long before something more positive will be noted. It is then essential so to train our powers of observa-

tion as to catch readily these changes of expression, for by this means we may sometimes foretell the outbreak of grave illness.

In estimating the nature of the complaint we shall often derive assistance from our general knowledge of the course of disease; of the special complications by which certain disorders are apt to be accompanied; and of the pathological consequences which may follow the occurrence of acute illness. Thus, in rickets we remember the extreme susceptibility of the body to changes of temperature and anticipate catarrhal derangements. Being aware, also, of the singular convulsive tendency in this disease, we are not alarmed by the occurrence of convulsive attacks, but at once search for some external source of irritation by which such attacks may be explained. Again, knowing the frequency of rheumatic fever in early life, and the rapidity with which the heart becomes affected secondarily, although the articular symptoms may be insignificant, the slightest stiffness in the joints, accompanied by a rise in the temperature, should at once suggest a most careful examination of the heart and chest generally. After an attack of measles or of whooping-cough, we should be on the watch for catarrhal pneumonia and acute tuberculosis. After scarlatina we look for acute desquamative nephritis and dropsy. In all cases where the symptoms are obscure and indefinite, and the most thorough examination fails to supply any sufficient materials for a positive conclusion, we must note the presence or absence of diathetic conditions—whether or not there be in process of development any constitutional state which might be itself the cause of disease, or might predispose special organs to be more readily affected by injurious influences; secondly, we must take into account any prevailing epidemics to which the patient may have been exposed; and lastly, we must inquire into the hygienic and dietetic conditions under which the child has lately been living. Still, in many cases it will be necessary to wait

before we can decide upon the nature of the ailment. In case of doubt no positive opinion should be hazarded, for few mistakes are more fatal to the reputation of the practitioner than to class as trifling a complaint which is afterwards discovered to be of a serious character.

In the *prognosis* of disease in children it is necessary to speak with much caution, for changes in them take place with great rapidity, and the expectations of one hour may be completely falsified by the next. In the early stage of an acute disease we must not allow ourselves to be deceived by the seeming severity of the symptoms, nor, when the disease is fully established, must we too hastily assume, however grave the condition of the patient, that the case is necessarily a hopeless one. In early life the re-adjusting power of nature is very great, and instances of unlooked-for recovery must be familiar to everyone whose practice has lain much amongst young children. When physic has become useless, careful nursing still remains; and the aphorism, "as long as there is life, there is hope," is especially true in the case of a child. On the other hand, we must be careful not to err in the opposite respect, and to pronounce too favourable an opinion in a case of serious disease because the symptoms for the time are mild; for, if cases frequently change rapidly for the better, they also often change rapidly for the worse. Indeed, sudden death is not at all uncommon in very young children, especially in cases of acute disease where the strength has been very suddenly reduced; and chronic diseases sometimes end very abruptly.

Children are more amenable than adults to the action of drugs, and the *treatment* of their diseases is consequently more easy. But treatment does not consist entirely, or even principally, in the administration of medicines. By attention to diet, hygiene, and the general management of a child, we can exercise a very important influence upon the course

and termination of an ailment ; and the mere giving of physic is only a small part of the duties of a physician.

In treatment, our first object should be to remove, if possible, the cause of the disorder. The cause is, however, often obscure, or, if detected, may be no longer in existence, the disease being merely the effect of some transient impression which has ceased to operate. In most functional derangements, however, especially those affecting the alimentary canal, some error in feeding or in management may usually be discovered, the remedying of which will produce an immediate improvement in the symptoms ; and in all cases the cause or causes should be diligently searched for, and the removal of them, when found, should be looked upon as the first and most important step in the treatment. Without this preliminary step, indeed, treatment will often meet with little success. Thus, in the case of rickets, we shall find very little benefit from the administration of tonics and cod-liver oil so long as the child continues to be fed on indigestible food, and to live in a close, ill-ventilated residence. Here treatment consists essentially in removing the causes which are producing the disease ; for cleanliness, fresh air, and a nutritious, digestible diet will by themselves be sufficient in mild cases to cure the disease without any aid from drugs being required.

Indications for treatment are also furnished by the nature and seat of the disease ; but in following such indications we must not allow ourselves to be led away too much by narrow rules as to drug-giving. In all cases we should be careful to treat the patient rather than the disease. Our object is to hasten the restoration of the child to health ; and this end will be gained far more effectually by attending carefully to symptoms—general as well as local—rather than by pursuing any one rigid line of treatment to which the malady is expected to succumb. It is, indeed, especially the case with children that the consideration of symptoms is of the first importance, whatever be the nature

of the disorder ; and in many cases treatment must be directed exclusively to these, although they may arise from a distant organ, and be the result of the general disturbance produced in the system through sympathetic irritation excited by some general or local ailment. Thus, the vomiting which frequently accompanies the onset of fevers and local inflammations may assume so violent a character as to place the life of the patient in peril. The beginning of variola is especially liable to be accompanied by excessive vomiting. In such a case treatment must be directed at once to control the dangerous symptom, and this irrespective of the seat or nature of the complaint in the course of which this complication has arisen. So also in cases of sympathetic convulsions occurring in the course of an acute disease, measures must at once be taken to quiet a perturbation of the nervous system which might exercise a very unfavourable influence upon the termination of the illness.

In acute cases the treatment will vary very much according to the age of the child, according to the presence or absence of diathetic or constitutional conditions, and especially according to whether the case be one of primary or secondary disease. If the child be very young, or be of feeble constitution, or be already weakened by previous disease, we should be careful not to depress ; but, on the other hand, we cannot always hope by mere stimulation to tide over the evil day, in the expectation that by such means the patient may be enabled to outlive the disease. Too early stimulation in certain diseases, such as bronchitis and croupous pneumonia, will be found merely to increase the severity of the inflammation, and may by itself be sufficient to determine the fatal issue. In bronchitis, especially, death has, I believe, been often caused in young children by a too early administration of carbonate of ammonia. Medicines—and stimulants do not belong to the least valuable class—are beneficial when selected with judgment,

and given at a proper time ; but they may be very hurtful when given imprudently and without discrimination. It is, then, of special importance in dealing with children, upon whom the action of medicines is so powerful, that we should have a distinct idea of the object sought to be obtained by the administration of a drug—that we should have a definite therapeutic purpose in all we do. Therefore, until we find some trustworthy indication for the use of a drug, we should be careful to give nothing of importance, but should content ourselves with ordering some harmless fluid, watching meanwhile for any symptoms which may arise to guide us to the employment of more active measures.

There is one class of remedies which is of singular value in the treatment of the diseases of young children—viz., the alkalies. In all children (in infants especially) there is a constant tendency to acid fermentation of their food. This arises partly from the nature of their diet, into which milk and farinaceous matters enter so largely ; partly from the peculiar activity of their mucous glands, which pour out an alkaline secretion in such large quantities. An excess of farinaceous food, therefore, soon begins to ferment, and an acid is generated, which stimulates the mucous membrane to further secretion. In all chronic diseases, and in many of the acute disorders, this sour condition of the stomach and bowels is present. Alkalies are therefore useful—firstly, in neutralising the acid products of this fermentation ; and secondly, in checking the too abundant secretion from the mucous glands. A few grains of soda or potash, given an hour or two after taking food, will quickly remedy this derangement and remove the distressing symptoms which arise from it. In the chronic diseases, indeed, attention to this point is of especial importance : for by placing the stomach and bowels in a healthy state, and insuring a proper digestion of food, we put the child in a fair way of recovery ; and prepare the way for the administration of

tonic and strengthening medicines, by which his restoration to health is to be brought about.

In prescribing for infants, an aromatic should always be included in the mixture. The aromatics are useful, not only for their flavouring properties, but also for their value in all those cases of abdominal derangement where flatulence, pain, and spasm, resulting from vitiated secretions and undigested food, are present to increase the discomfort of the patient. Such dyspeptic phenomena are usually relieved rapidly by the use of these agents ; and aniseed, cinnamon, caraway seed, or even tincture of capsicum in minute doses, will be found important additions to the prescription in all cases where alkalies are required.

With regard to quantity :—The proper dose of a medicine cannot always be calculated according to the age of the child, and does not in all cases bear the same proportion to the quantity suitable for an adult. For certain drugs children show a remarkable tolerance, while to the action of others they show as remarkable a susceptibility. Thus, opium, it is well known, acts upon a child more powerfully than would be expected, judging from the mere difference of age. It should therefore be given to infants with a certain caution, especially if the child be enfeebled by disease. It is, however, a medicine which is of especial value in the treatment of the diseases of infancy, and may be given without fear if care be taken not to repeat the dose too frequently. Belladonna, on the contrary, can be taken by children in large quantities. A child of two or three years old will bear without inconvenience a dose which in an adult might produce very uncomfortable symptoms.* Lobelia, again, is a remedy which is very well borne by children. Dr. Ringer has given it to “very young children” in doses of five minims every hour, and in

* It is important to remember this in giving belladonna for its sedative effects, as in pertussis.

no case has he noticed any ill effects to follow its administration. Arsenic should be given to children over five years of age in the same dose as that used to adults, and infants of a month or two old will take one drop of Fowler's solution three times a day with great benefit in cases of gastric catarrh. The influence of mercury upon young children deserves remark. It seldom in them produces stomatitis or salivation; but an excess of the drug is not therefore harmless: its influence is seen in the irritation of the alimentary canal which it so often excites, and in the profound anæmia which it induces. The anæmia which is so common a sequence of constitutional syphilis in infants is no doubt often a result of too long-continued mercurial treatment.

When the physician has prescribed the necessary medicines, and given directions as to the diet and general management of the child, his duties are not necessarily ended. It is his part not only to advise, but also, as far as possible, to see that his directions are carried out. The soundest advice is useless if it be not put into practice, and all the art of the physician may be neutralised by weakness or indifference on the part of the attendants. From a feeling of injudicious fondness a mother will sometimes neglect the duties intrusted to her, through a fear of inflicting pain or an unwillingness to subject the child to annoyance. In these cases the medical attendant should impress upon her the importance of the measures to be taken, and should remind her that he is there only as an adviser, while it is her task to put his advice into practice. During an illness a child is too often considered as one whose slightest desire is to be instantly gratified, on the ground that opposition might aggravate the severity of his complaint, even if it did not exercise an unfavourable influence upon the issue. A fretful, petted child well knows how to take advantage of this feeling, and the unwise indulgence of those around him is often the most serious

obstacle to his recovery. If, however, a child be well enough to form unreasonable wishes, he is not too ill to bear the disappointment of a refusal, and the mother should endeavour with gentle tact to divert her child's thoughts from any hurtful inclination of the moment, remembering that the truest kindness consists in abiding strictly by the directions of the physician, and in allowing nothing which would in any way interfere with recovery.

CHAPTER II.

COLLAPSE OF THE LUNG (POST-NATAL ATELECTASIS).

COLLAPSE of the lung a common lesion in the child—Method of production and causes which induce it—Anatomical characters—Symptoms and physical signs.

Case I.—Sudden death from pulmonary collapse in a weakly, wasted infant.

Case II.—Collapse of lung occurring in a case of pertussis—Comment.

Case III.—Collapse of lung with laryngismus stridulus—Comment.

Case IV.—Bronchitis and collapse of lung in a rickety child—Comment.

Diagnosis of pulmonary collapse—Treatment.

COLLAPSE of the lung is one of the most important diseases of early life. When occurring after the lungs have been fully expanded at birth, it is almost always the consequence of pulmonary catarrh, and is one of the causes which render bronchitis so fatal a disease in the child.

The immediate cause of collapse of the lung is the presence of mucus in the tubes, and to Dr. Gairdner, of Glasgow, we are indebted for a simple and conclusive explanation of the mechanism by which this exhaustion of the lobules is effected. A plug of mucus is carried inwards by the force of inspiration along an air-tube of constantly diminishing calibre, and becomes at last arrested, obstructing the whole channel of the tube. Although, however, the narrowness of the air-tube prevents its further advance, the plug of mucus can still be moved in a direction contrary to that in which it has lately passed. At each expiration, then, it is dislodged sufficiently to allow the passage outwards of some of the air still left in the air-cells ; but at each inspiration it is driven back again into the tube, so as completely

to close it against any air entering to replace that which has just escaped. By several repetitions of this to-and-fro movement the air in the lobules beyond the obstruction becomes completely exhausted, and collapse is the result.

This mechanism may take place in several spots in the lung at the same time, and the extent of the collapse in each instance depends upon the size of the plug and the point at which the obstruction of the bronchial tube takes place.

The above is the immediate cause of collapse of the lung, but the process is assisted by anything which renders difficult the entrance of air into the bronchi. There are, therefore, predisposing causes which promote the retention of mucus in the tubes, and assist its obstructive action. Of these the first is an inability to cough and expectorate. New-born infants do not know how to cough, for the act of coughing is only in part an involuntary act; it is in great part due to an effort of volition to remove an impediment to free respiration. The infant, then, is conscious of the impediment to the free passage of air, but has not yet acquired a knowledge of the means by which it may be remedied, and the mucus remains in the tubes because the child does not exert the force necessary to expel it.

Again, the child may be unable to cough. For the act of coughing, a quantity of air must be drawn in past the obstruction; the glottis is then closed, and pressure is brought to bear upon the lungs by the expiratory muscles. While the pressure is at its height, the glottis is relaxed, and a blast of expired air rushes out, carrying with it the mucus which caused the obstruction. If, however, a sufficient quantity of air cannot be drawn in past the collection of mucus, or if the general weakness be so great that adequate pressure cannot be brought to bear upon the lung, the effort is insufficient to cause expulsion of the phlegm in the tubes.

Another predisposing cause of collapse is weakness of the inspiratory act. This may be a result of general exhaus-

tion, the patient being too feeble to take a deep breath, and is often the cause of extensive collapse in dying persons. Air is drawn but imperfectly past the mucus at each inspiration ; while at each expiration the air in the vesicles is expelled by the natural elasticity of the lung. After a time the cells become completely exhausted of their air, and sink in.

The act of inspiration may, however, be weak from other causes than actual muscular debility. The patient may be prevented mechanically from expanding his chest-cavity, either from some impediment in the abdomen to the descent of the diaphragm, or from deficient rigidity of the thoracic walls, which cannot resist the pressure of the external air. Thus, ascites or tumours distending the abdomen may so interfere with the descent of the diaphragm that the inspiratory effort is too feeble to draw in air past the mucus ; and in such cases a comparatively trifling bronchitis may lead to very extensive collapse, for the residual air soon becomes exhausted, and the vesicles sink in. In infants, whose nearly circular chest allows of little lateral expansion the respiration is principally diaphragmatic ; any obstacle to the action of this important muscle is therefore in them attended by the most serious consequences.

Again, a want of rigidity of the chest-wall in the child greatly assists the production of collapse. Even in a healthy child the parietes of the chest are very flexible, as is seen in all cases where there is any difficulty of respiration. The ribs at the lower parts sink in under such circumstances, so that very imperfect dilatation takes place of the lower lobes of the lungs. If, however, the ribs are softened by rickets, the flexibility of the thoracic parietes is very much increased. No dilatation of the lower part of the chest takes place at all : instead, there is at each inspiration so much recession of the parietes of the thorax in the inferior regions that very little air indeed is able to penetrate into the lower lobes of the lungs. If bronchitis come on in

such a patient, the recession of the chest walls becomes greater than ever, for the presence of secretion in the tubes offers an additional obstacle to the entrance of air. Any air, therefore, which the lower lobes may contain soon becomes expelled by the elasticity of the lung, and collapse of that portion of the lung takes place.

Infants are especially liable to atelectasis, because their bronchi are small and easily obstructed; besides which, they are feeble and quickly depressed, and their chest-walls are naturally deficient in rigidity and yield readily to external atmospheric pressure.

Collapse of the lung is always a secondary lesion. The accident is especially liable to happen in cases of exhausting disease, and is prone to complicate those forms of illness in which catarrh of the lung is a prominent symptom. Thus, in whooping cough and measles, when these infectious complaints attack the infant, atelectasis is common. It is often also met with in the course of diseases which interfere with the passage of air through the windpipe, as diphtheria, spasmodic laryngitis, and retro-pharyngeal abscess. We may find it in every variety of morbid state which in any way lessens the force of the inspiratory act, either by opposing the descent of the diaphragm, as in ascites, flatulent distension, or abdominal growth; by reducing the rigidity of the chest-wall, as in advanced rickets; or by lowering the muscular strength of the patient, as in chronic exhausting disease. Often, many of these causes are in operation at once. Thus, a rickety child begins to suffer from catarrh of the bowel, which interferes with his nutrition and reduces his strength. In this case the force of the inspiratory act, already impaired by the softened state of the ribs, is still further weakened by loss of power in the inspiratory muscles. Suddenly, great accumulation of gas takes place in the belly from fermentation of food. Expansion of lung, carried on hitherto under great difficulties, ceases to be possible, and the child dies more or less rapidly from asphyxia.

In all these cases, no doubt, a certain amount of pulmonary catarrh must enter into the disturbance; but it is important to be aware that this catarrh may be of the most insignificant kind. A comparatively small plug of mucus may do all the mischief. Often the child has not been noticed to cough, or it is only when questioned specially upon the point that the nurse can remember that he has done so.

The size of the portion of the lung which has become collapsed varies, as has been said, according to the size of the plug of mucus, and the point at which the obstruction of the bronchial tube takes place. Collapse is therefore said to be either diffused or lobular, according as it affects numerous adjoining lobules over a considerable surface, or is limited to single ones or to single clusters. In the diffused form the greater part of a lobe is often affected, usually at the posterior region, and the circumference is less distinctly circumscribed than is the case with lobular collapse, the collapsed portion being seen to pass gradually at many points into the (usually) emphysematous tissue around. The lobular variety is often situated at the anterior edges of the lungs, although it may occupy any other part, and is very abruptly marked off by the interlobular septa. In either case the collapsed portion is solid, but still soft, to the feel; it is tough and not easily broken down, does not crepitate, and sinks instantly in water. It is depressed below the level of the surrounding lung, and is more or less purple in colour, the depth of tint depending upon the amount of blood contained in the tissue. When cut into, the section is smooth, and much blood or bloody serum exudes on pressure. Lastly, if the affection be recent, the collapsed portion of lung can be completely re-inflated through the bronchus. This is a very valuable test, and conclusively proves the nature of the lesion. If, however, the collapse be less recent, complete distension cannot always be effected, for where the lesion is of long standing

certain nutritive changes are apt to take place in the inactive vesicles, which to some extent alter the structure of the lung at the seat of disease; and in cases where pneumonic changes have begun in the tissue, perfect re-inflation becomes impossible.

The *symptoms* of pulmonary collapse vary to a certain extent, according to the area of the lung involved in the disease. If the collapse be very sudden and extensive, so as very rapidly to arrest the functions of a large part of both lungs, death may follow almost immediately. This is not at all an uncommon cause of sudden death in whooping-cough, or in cases of catarrh attacking young and weakly children whose strength has been rapidly reduced by some exhaustive disease, as a severe attack of diarrhoea. In cases where the portion of lung collapsed is less extensive, the occurrence of this complication is often indicated by a convulsive fit, and the fits may be repeated again and again until death ensues. Convulsions are indeed not an uncommon symptom, even in cases where the area of lung involved is small, although they are by no means present in every case.

A child in whom collapse of the lung occurs begins at once to exhibit evident signs that the pulmonary catarrh from which he has been previously suffering, is no longer uncomplicated. He becomes restless, and moves his arms uneasily about. His face shows signs of distress, and his features become pinched. The eyes are hollow, the lips bluish, and the eyelids and mouth livid. He whines plaintively, but seldom cries. The skin is cool, and often moist with a cold, clammy perspiration, and a thermometer placed in the rectum shows that the internal temperature of the body is little, if at all, elevated above the natural level. The pulse is very small and feeble, and rather hurried. The breathing is irregular, shallow, and exceedingly rapid, often reaching 60, 70, 80, or even more, in a minute; and this rapidity of breathing causes a perversion of the pulse-respiration ratio which is not exceeded even in cases of the

severest pneumonia. The nares dilate widely at each inspiration, but the inspiratory efforts are seldom violent, except in some cases of rickets when there is much softening of the ribs. The cough becomes very feeble, and sometimes seems almost suppressed. In some cases the attempt to cough is followed by gasping efforts, as if the child were being suffocated, while the lividity of his face is increased, and his limbs are agitated by spasmodic movements. When the collapse is extensive, the child will take little food, and, if an infant, generally refuses the breast. This is partly occasioned by his respiratory wants, but it often also proceeds from a difficulty in swallowing, which is a frequent symptom in the disease.

Physical signs, when they occur, are found at once, and the rapidity with which they arise is an important element in the diagnosis. On examination dulness is commonly found at one or the other base, with slightly increased resistance. This is in most cases limited to one region of the thorax, usually the posterior base. Sometimes the lesion is found in the two lungs at the same time; when thus symmetrical a favourite seat is a narrow vertical strip extending for two or three inches along each side of the spine. The dulness is best detected by gentle percussion with one finger. It is seldom complete, and is often difficult to estimate, on account of the presence of emphysema; for where the collapsed area is of small extent, local dilatation of the air-cells may render the sound elicited by the finger perfectly clear, or the percussion-note may have merely a slight wooden quality, which presents nothing characteristic. With the stethoscope we hear bronchial breathing, which may be loud and intense, or weak and faintly marked. There is usually a certain amount of coarse crepitation, which is produced, not in the collapsed portion itself, but in the tissue around. Vocal resonance may be increased over the seat of collapse, but more often it is annulled. The same may be said of vocal vibration

but in children vibration of the chest-wall is often absent, even in healthy subjects, and this sign is therefore seldom serviceable in diagnosis. It is only in cases where several contiguous lobules are affected—that is to say, where the collapse is diffused, and then only when the lesion is in contact with the chest-wall—that any physical signs are present. In cases where solitary lobules are exhausted of their air, or where the diseased condition is seated in the interior of the lung, no evidence at all is to be derived from examination of the chest.

For the production of this lesion it is not necessary that the pulmonary catarrh should be a severe one ; cases even of slight catarrh may be followed by rapid death, when the patient is very young and has been lately reduced by some exhausting disease.

CASE I.—SUDDEN DEATH FROM PULMONARY COLLAPSE IN A WEAKLY, WASTED INFANT.

Edward K., 10 months old, one tooth, a very small and miserable-looking child, was admitted into the East London Children's Hospital on October 15. He had been suffering for some months from chronic diarrhœa, and had been wasting more or less all his life. A rigid diet, from which milk was carefully excluded, combined with warmth and an astringent mixture, soon stopped the diarrhœa, and the child was just beginning to regain flesh, when on October 26 he was noticed to have a slight cough, with a little rattling of phlegm in the chest. He, however, took his food as usual, and nothing occurred to occasion the slightest anxiety, when early on the following morning he was found to be very greatly altered in appearance. His face was livid, and his breathing very rapid and shallow. On examination of the chest, the breath-sound was very weak and distant, but was unaccompanied by rhonchus. He had no convulsions, and died quietly very shortly afterwards. On

examination of the body, a pyramidal strip of collapse was found at the posterior surface of each lower lobe, reaching upwards to the upper border of the lobe. The breadth of each was about an inch and a half at the lower border, and the thickness about half an inch. They could not be entirely re-inflated. The large intestine throughout its whole extent was inflamed along the summits of the longitudinal folds of mucous membrane. In the small intestine Peyer's patches were injected, and there was some arborescent congestion of the mucous membrane. There were no ulcerations.

In the above case the physical signs were only imperfectly ascertained. When seen by the House-Surgeon, the child was evidently dying, and it was thought unadvisable to disturb him more than was absolutely necessary ; otherwise, no doubt, careful percussion would have revealed the existence of dulness over the collapsed portions of the lung.

CASE II.—COLLAPSE OF THE LUNG OCCURRING IN A CASE OF PERTUSSIS.

Albert S., age 1 year and 9 months, twelve teeth, was seized, on April 22, with a severe attack of diarrhœa, which considerably depressed his strength, for although he had for two months been able to walk, he was now scarcely able to stand. All his joints were nodular and his ribs beaded, but there was no actual deformity of the chest. The diarrhœa was soon arrested by treatment, but on April 29 he began to cough a good deal, and a week afterwards whooping-cough was completely developed.

May 10.—Whoops very much, and the fits of coughing are very frequent, and are invariably followed by vomiting, so that very little food is retained upon the stomach. Medium-sized bubbling is heard over both sides of back. Temperature in axilla, 100° Fahr.

May 14.—The fits of coughing are less frequent, but the

laryngeal spasm is very intense, and the child gets purple in the face after each attack. Even between the attacks the lips remain very dark in colour, and the eyelids purplish. Skin moist and not hot. Subcrepitant rhonchus is heard all over back.

May 17.—On the evening of May 15 the child had a fit of convulsions which lasted ten minutes. Since then he has seemed very weak, and has taken nothing but occasionally a little broth. Perspires a good deal. Temperature, 99.4° ; pulse, 180. Lies back with head low; pale and moist-looking in face; eyelids and lips dark red; lividity under eyes. Respiration very rapid, 120, and nares acting, but the breathing is very shallow, and there is no contraction of accessory muscles. The cough is evidently painful, for the child makes unusual efforts to prevent it. When the desire to cough comes on, the face becomes congested, the eyes get fixed, the brows contract, and sweat breaks out on forehead. Has not whooped since yesterday evening.

On examination of the chest there is a little dulness at the right base behind, and the breathing there is higher-pitched and drier in quality than on the opposite side. Subcrepitant rhonchus is heard at each base, back and front.

On the following day, May 18, the convulsive attacks returned with great violence, and the child died.

Autopsy.—The lower half of the inferior lobe of the right lung was found in a state of collapse, and there were smaller patches of collapse scattered about the upper part of the same lobe. There was also a little collapse of the lower lobe of the left lung, but this was much less extensive than on the right side. There was no trace of pneumonia, but the lining membrane of the larger tubes was a little reddened. At the anterior border of the inferior lobe of the right lung a string of quite recent lymph was found, and there was a little reddening of the opposed surfaces of the pleura in that situation.

Whooping-cough, when severe, has always a tendency to

cause collapse, for a violent fit of coughing with extreme spasm reduces the quantity of residual air to a very low point, while at the same time the exhaustion which follows a severe paroxysm weakens the force of the inspiratory act. The thick, ropy mucus contained in the air-tubes offers already an obstacle to the entrance of air, and the rapid, feeble breathing which is noticed during the periods of depression is little calculated to overcome the difficulty. In such cases the stages which mark the production of collapse are—

1. Partial exhaustion of the air-cells during the prolonged cough.

2. State of great depression, during which the inspiratory efforts are insufficient to draw in air past the mucus which loads the tubes.

3. Passage outwards at each expiration of some of the residual air left in the cells.

Unless, therefore, the strength of the inspirations can be increased, or the obstructing mucus can be dislodged, the occurrence of collapse may be looked upon as in the highest degree probable. If rickets be present in addition, the obstacle to full inspiration offered by the softened ribs increases still further the danger of the case.

In the above example the child was the subject of rickets, and had besides been already much depressed by the abdominal disorder before the whooping-cough began. The cough was very violent and the spasm severe, and the further reduction of strength which resulted from the excessive vomiting was so great that the occurrence of collapse could be looked upon as almost a certainty. The evident pain excited by the cough was an unusual feature in the case; for although children frequently strive in pertussis to repress the cough when they feel it to be coming on, yet they seldom manifest any sign of actual pain in respiration. The existence of pain was, however, quite accounted for by the local pleurisy discovered on examination of the body.

CASE III.—COLLAPSE OF THE LUNG WITH LARYNGISMUS STRIDULUS.

G. E., one month old, a fat, well-formed child, was first seen on July 28. He had been quite well until two days before, when he had begun to cough. The cough had since increased, and had sometimes been followed by what were described as "fits." He was said to take the breast well, but to vomit a part of the milk very shortly afterwards.

When first seen—Child rather red in face; seems rather depressed; pulse 116; respiration 82. Breathing shallow as a rule, but the respirations are not all of equal depth, and there is sometimes considerable recession of the lower part of the chest during inspiration. Skin does not feel at all hot to hand. Temperature in rectum 100.4° . The child gives a single or double hack at intervals, and sometimes seems as if he could not draw his breath immediately after it, so that his face gets congested and his eyelids and mouth become livid. On one occasion after the cough he had one of the so-called fits. His eyes became fixed, his limbs were moved uneasily about for a few seconds, and his face was livid; he afterwards lay motionless with closed eyes for some minutes, during which the pupils were contracted as in ordinary sleep; they acted readily with light. The fontanelle is slightly depressed. He perspires a little about the face. The bowels are rather confined, and he has not passed water all day.

On examination of the chest, a little incomplete dulness is found at the right base behind, with slightly increased resistance. The respiration is very weak, and when the child takes a deep breath, which occurs at rare intervals, a fine subcrepitating rhonchus is heard at the end of inspiration.

A mixture containing bromide of potassium and sal volatile was ordered; and the frequent application of weak mustard poultices was recommended to the chest and back.

July 30.—Since last report the child has had four “fits.” He seems very much in the same state as before. Still takes the breast well, but is said to pass very little water. The same difficulty of breathing and lividity are noticed after the cough. Pulse 144 : respiration very shallow and irregular, 76 ; temperature in rectum 98°. Lies with mouth closed, and nares not acting. No dulness is found now at the left posterior base, but all over both sides of back a fine sub-crepitant rhonchus is heard with inspiration, and, if the child takes a deeper breath, with expiration. It is not evolved in puffs, and is not at all like the crepitation of pneumonia.

A senega and ammonia mixture was ordered, and the sinapisms were repeated, but the child died the next day in convulsions.

Unfortunately, no post-mortem examination was obtained, but there could be little doubt as to the nature of the case. The incomplete dulness limited to the right base behind, the weak breathing, combined with the severe general symptoms, and the absence of pyrexia, all these occurring in the course of a pulmonary catarrh, could only be due to collapse of the lung. The dulness was not found after the first visit, probably owing to the occurrence of local emphysema. The so-called “fits” were no doubt slight attacks of laryngismus stridulus.

CASE IV.—BRONCHITIS AND COLLAPSE OF THE LUNG IN A RICKETY CHILD.

Charles W., aged 1 year and 8 months, twelve teeth, was admitted into the East London Children's Hospital on December 13. “The child has been brought up partially by hand, and has been fed from birth with farinaceous food. Never had pertussis or measles. Has always been very subject to cough. Three months ago had an illness which was

called choleraic fever. His breath was very bad during this time, and he was much pulled down. Since then his chest has been 'falling in.' For the last few days he has had a very bad cough."

State on Admission.—Child very rickety. Head long from before back; fontanelle size of a florin, depressed; ends of long bones enlarged, but limbs generally straight; muscles excessively flabby and wasted; spleen and liver can be felt, of natural size, below false ribs; antero-posterior diameter of chest very long, and lower part of sternum very prominent; chest deeply grooved laterally at both sides, and the enlarged ends of the ribs are seen lying on the inner face of the groove; below the sixth rib the chest is much expanded laterally; the back is very flat in the scapular regions, and the spine at the lower dorsal region is curved sharply backwards. At each inspiration there is great recession of the chest-walls at about the level of the sixth rib, so as to give the appearance of a deep furrow extending across the body in that situation. Heart's impulse can be felt from the left side of ensiform cartilage nearly to nipple. At the right base behind, the percussion-note is very high-pitched as far as the lower angle of the scapula. Respiration there bronchial, with subcrepitant rhonchus, and increased resonance of cry. In the right interscapular region the breathing is tubular, although there is no dulness. In left interscapular region respiration is bronchial; elsewhere on that side it is natural, and is accompanied over the lower two-thirds with subcrepitant rhonchus. There is no dulness on left side. In front the physical signs are perfectly natural on both sides of the chest. Temperature at 9 a.m., 99°; pulse 130; respiration 72. At 9 p.m., temperature 100°. The child was ordered a senega and ammonia mixture with five drops of vinum ipecac. every four hours, and twice a day the whole body was to be rubbed with the following liniment:—
℞ Olei morrhuae ʒj., olei terebinthinæ ʒij., liniment. saponis co. ʒj., m. ft. linimentum.

December 17.—Looks very much better ; lying on back with head low, and sucking all the fingers of his left hand ; less recession of chest-walls than was ; very little dulness at right base, and respiration there loud and slightly blowing, with coarse subcrepitant rhonchus accompanying the inspiration ; at lower angle of scapula, and from that point to supra-spinous fossa, diffused blowing respiration is heard, without rhonchus ; at the left back percussion is perfectly natural ; there is a little subcrepitant rhonchus at the left base, and at the angle of scapula the respiration is slightly blowing, although much less so than at right side ; respiration generally over both sides in front seems rather more bronchial than is natural, and vocal resonance rather increased. Temperature since last report has been almost uniformly 100° ; pulse 120 ; respiration 66 to 74. This morning, at 9 a.m.—temperature 98° ; pulse 114 ; respiration 66. Temperature at 9 p.m., 99° .

20th.—Percussion is now everywhere perfectly natural, but respiration is still rather more bronchial at right base than elsewhere, and in the right interscapular region diffused blowing ; no rhonchus heard about the chest except at right base, where there is an occasional clicking sound at the end of inspiration. Temperature at 9 p.m., 98° .

After this the child went on well till December 28, when he had an attack of diarrhoea which lasted two days. He also began again to cough, and seemed very much weakened by the complication.

31st.—Eyes rather sunken ; cheeks flushed ; nares acting ; cough loose ; percussion of back natural, but fine bubbling is heard all over both backs, most marked at bases. Respiration in right interscapular region very blowing, as before. 9 a.m. : Temperature 99.2° ; pulse 120 ; respiration 72. 9 p.m. : Temperature 101° .

On January 1 face became very livid ; he coughed a great deal, and was very restless. Temperature 101° ; pulse 150 ; respiration 50. At night he slept, but not very quietly,

and early on the following morning woke up with a scream, and died almost immediately.

Autopsy.—Right lung slightly emphysematous at apex of upper lobe. The greater part of the upper lobe, the whole of the middle and half the lower lobe deep purple in colour, firm, shrunken, and non-crepitant, evidently in a state of collapse. Redness of the mucous membrane, of larger bronchi, and many of the smaller bronchial tubes filled with plugs of thick mucus. In the left lung the posterior half of the upper lobe and part of the lower lobe also collapsed: the collapsed portions could be perfectly re-inflated. Right side of heart distended with black blood. Other viscera healthy.

In this case the child was the subject of profound rickets; the chest was much deformed, the ribs exceedingly soft, and the consequent difficulty of respiration extreme. In such a child on the occurrence of severe pulmonary catarrh collapse of the lung may, for reasons already explained, be confidently predicted. On his admission into the Hospital the child was already suffering from partial collapse of the lower lobe of the right lung, as was shown by the dulness on percussion limited to the posterior region, and the bronchial breathing, combined with the low temperature. The blowing breathing in each interscapular region was not due to disease of the lung, but was the consequence of the flattening of the back in that situation, which approximated the large divisions of the bronchi to the chest-wall. This flattening, as was made evident by a section of the chest drawn from the cyrtometer, was especially marked on the right side, and at that point the respiration was excessively blowing, almost tubular in quality. Under the influence of treatment, and especially of the frequent application of a highly stimulating liniment, the bronchitis was relieved, the lung resumed its functions, and the urgent symptoms disappeared. Unfortunately, however, the child caught a fresh cold. Diarrhoea came on, which greatly reduced his

strength ; the bronchitis returned ; and all the conditions tending to cause pulmonary collapse were again in full force. The inevitable result followed ; both lungs became extensively collapsed, and the child died. The temperature in this second attack was rather higher (101°) than it had been during the first. This was no doubt due to the early stage of the bronchitis, which has the effect of at first elevating the temperature. That it was not owing to any pneumonic complication was shown by the perfect re-inflation of the collapsed portions of lung in the post-mortem room.

From the cases of collapse of the lung above narrated, it is evident that in young subjects the physical signs differ considerably from the signs which would be present in similar cases in the adult. In pulmonary collapse occurring in the adult the percussion note is slightly dull, vocal vibration and vocal resonance are diminished or annulled, and the breathing is suppressed, or is very weak, distant, and bronchial. Occasionally, when a deep breath is taken, a dry, minute, crackling bronchus is caught faintly at the end of inspiration. In the child, on the contrary, and especially in the infant, sounds generated in the chest, owing to the small size of the thorax, are more readily diffused, and are conveyed with much greater distinctness to the ear. As a consequence, the intensity of the breathing, instead of being diminished, may even be increased ;* so, also, the resonance of the voice may be stronger than natural, and, unless the collapsed portion of lung be very

* A similar modification of the physical signs is seen in the case of pleurisy in the child, where, owing to the small area in which the phenomena occur, the physical signs may be very strikingly at variance with those to which we are accustomed in the adult. So great, indeed, is the difference, that any one unacquainted with the peculiarities of pleurisy in children would in all probability fail to estimate rightly the nature of the disease.

extensive, subcrepitant rhonchi generated in the immediate neighbourhood may be heard clearly over the diseased spot. While, therefore, the diagnosis of pulmonary collapse in the adult is an easy matter, in the child its detection is more difficult, owing to the resemblance borne by the physical signs to those furnished by other diseases.

Dulness, bronchial breathing, a fine crepitating rhonchus, and increased resonance of the cry might, on a hurried examination, suggest the presence of pneumonia or pleurisy. But the dulness is as a rule confined to the posterior base, and does not pass to the front of the chest as in pleurisy; the breathing never has the intense tubular quality so characteristic of pneumonia; and the crepitation, however fine it may be, is not evolved in puffs like the true pneumonic crepitus. The dulness is of a peculiar character, differing both from the incomplete dulness of pneumonia, and also from the dead, flat, toneless percussion-note of pleurisy. It is best detected by very light percussion with one finger, for, unless extensive, the collapse is limited to the surface, and does not penetrate deeply into the substance of the lung. When the collapse is symmetrical, and presents the pyramidal form which has been described, its detection is easy. When, however, it is confined to one lung the lesion is distinguished from pneumonia by the absence of tubular breathing and of true pneumonic crepitus. From pleurisy it differs by the very limited area occupied by the dulness, and by the fact that both the dulness and the sense of resistance are very much less than in cases where fluid is present.

Another and important distinguishing mark is furnished by the suddenness with which the signs occur. They appear and disappear with a rapidity which is not found in any other chest disease. But the diagnosis does not rest alone upon the characters of the physical signs. In all chest diseases, and especially in pulmonary collapse, it is important to take into account the general symptoms, and

to compare these carefully with the signs furnished by physical examination, before arriving at a positive conclusion. The general symptoms come on at once, and are very characteristic. The distress, the lividity, the rapid shallow breathing, the faint cough, and the feeble cry—these symptoms occurring suddenly in the course of a catarrh, and, combined with the absence of pyrexia, leave little room for doubt as to the nature of the disease. The use of the thermometer is in these cases of the highest importance. In pneumonia the temperature is very high. In pleurisy it may be natural, but in that disease, unless the accumulation of fluid be so great as to render its presence evident to the most inexperienced observer, there is no distress, no lividity, no great hurry of breathing,—no signs, in fact, to indicate any difficulty in the passage of the blood through the lungs.

Diffused collapse of the lung, then, differs from pneumonia by the absence of pyrexia, the absence of tubular breathing and true pneumonic crepitus, and by the suddenness with which the physical signs occur. From pleurisy it differs by the limited area and imperfect character of the dulness, the smaller sense of resistance, and by the much greater severity of the general symptoms; for these can only occur in pleurisy in cases where the accumulation of fluid is enormous, and where the physical signs are consequently so characteristic as to remove any difficulty from the diagnosis.

In cases where the pulmonary collapse is lobular, or where the lesion is not in contact with the chest-wall, the diagnosis must be made without any assistance from physical signs; but here, if many isolated lobules are involved, the very absence of physical signs supplies an important indication. The general symptoms which have been enumerated occurring suddenly in the course of a pulmonary catarrh, and without giving rise to any special physical signs by which they may be accounted for, suggest at

once the presence of lobular collapse of the lung. In catarrhal pneumonia, where special physical signs are also often absent, the temperature is elevated, and the breathing excessively laborious ; while in collapse the temperature is not elevated, and the breathing, although rapid, is shallow and quiet.

It is not always easy in cases of bronchitis to decide upon the presence or absence of accompanying lobular collapse : but in such cases collapse may be suspected if, with only moderate bronchitis and a normal temperature, the general symptoms are grave out of all proportion to the physical signs, or if, where the bronchitis is severe, the breathing changes from the laborious respiration of intense capillary bronchitis to the shallow, rapid breathing so characteristic of pulmonary collapse, while at the same time the lividity and general distress become still further aggravated, and the temperature remains low. If convulsions occur towards the end of an attack of bronchitis, it is probable that collapse of the lung has taken place.

In every case of pulmonary collapse occurring in the infant the prospects of the patient are very unfavourable. If the derangement, of which the atelectasis is a complication, amount to a severe bronchitis, the child's chances of escape are very small. So, also, if the resistance of the chest-wall be greatly reduced by rickety softening of the ribs, a comparatively mild catarrh may prove fatal through the inability of the child to inflate the lungs sufficiently to expel the obstructing mucus. In a young baby the danger in any particular case cannot be estimated merely by the amount of existing bronchitis ; indeed, it may happen that the apparent inoffensiveness of the primary derangement is itself an unfavourable element in the case. Thus, when atelectasis occurs in a wasted infant under the influence of a trifling pulmonary catarrh the prognosis is very serious, for the fact that under such circumstances atelectasis should have taken place indicates a very serious degree of weakness of the

inspiratory muscles. The occurrence of convulsions, of persistent increase in the lividity and torpor of the patient, and of inability to swallow, are all symptoms of the utmost gravity.

In children of three years old and upwards, collapse of the lung rarely occurs, unless the pulmonary catarrh is considerable. Therefore, in these cases the danger is in proportion to the amount of bronchitis. If this be not sufficiently extensive to cause death, the recovery of the patient may be safely calculated upon.

The *treatment* of pulmonary collapse consists in the adoption of means to increase the force of the inspiratory act, both by compelling the child to exert to its full extent the power he may possess, and also by adding to that power as much as possible, so as to effect a dilatation of the exhausted air-cells. In the first place, we must take care that there is no mechanical impediment to the expansion of the chest. The dress should be loose, and the child should be placed in such a position as to insure free movement of the thoracic walls. Thus, if the lesion be limited to one lung, the patient should be laid upon the back, inclining to the sound side. We should next adopt mechanical means for deepening the breathing. After certain acts, such as crying or vomiting, a deep breath is instinctively taken. It is therefore advisable to make the child cry by rough friction to the soles of the feet, or by the sudden application of a sponge dipped in cold water to the naked chest. The emetic should be one which acts quickly without producing much depression, such as oxymel of squill or one or two drachms of alum in syrup. Either of these may be given, and they will have the further advantage of dislodging and expelling some of the mucus which loads the air-tubes. If the feet are cold, they should be warmed by the application of a hot bottle; and the same may be applied, if necessary, under each arm as the child lies in his

cot. For increasing the child's inspiratory power, stimulation, both external and internal, is required. A highly stimulating liniment may be made by adding two drachms of oil of turpentine to four ounces of compound camphor liniment, or the linimentum ammoniæ of the Pharmacopœia may be used. Either of these should be well rubbed into the chest and back two or three times in the day. For internal administration the stimulating expectorants should be employed, especially carbonate of ammonia, senega, and squill. Four grains of quinine dissolved in half an ounce of sal volatile make a powerful diffusible stimulant, of which a few drops may be given frequently in a teaspoonful of milk or water. Five or ten drops of pale brandy may be given occasionally in addition. If the child be drowsy, he should not be allowed to sleep too long at one time, but he should be occasionally roused, so that some of the preceding measures may be adopted. The diet should consist of milk with (if the child be old enough) strong beef-tea. In bad cases, where there is difficulty of swallowing, the brandy-and-egg mixture of the Pharmacopœia, given at intervals with a teaspoon, is very useful. By the energetic use of the above measures very surprising results are sometimes obtained. But their employment must be energetic and immediate : no time must be lost, or the opportunity for successful interference will have passed away.

CHAPTER III.

CROUPOUS PNEUMONIA.

CROUPOUS pneumonia not common under two years of age—Symptoms and physical signs.

Case V.—Pneumonia introduced by a convulsive attack.

Case VI.—Pneumonia of apex with severe cerebral symptoms (cerebral pneumonia)—Comment.

Case VII.—Rapid course of pneumonia of apex.

Case VIII.—Pneumonia complicated by gastro-intestinal catarrh—Comment—Diagnosis from typhoid fever.

Case IX.—Pneumonia with diarrhoea in an infant—Comment.

Prognosis—Diagnosis—Treatment.

CROUPOUS pneumonia is a disease which is often met with in early life. It is less common in infancy than at a later period of childhood, but is seen with fair frequency even in children under two years of age. Croupous pneumonia is now regarded as a general disease of which the pulmonary inflammation is the local expression. Of the many arguments by which this opinion is supported one of the most convincing lies in the want of proportion which exists between the intensity of the general symptoms and the extent of the pulmonary mischief. The general symptoms, especially in young patients, may be serious and even alarming although the area of lung which becomes the seat of inflammatory consolidation is curiously limited. This inflamed area may occupy any part of the chest. It may be confined strictly to the apex or axilla of one side, and, indeed, is found in these situations little less often than at the posterior base. Therefore in any case we should never be satisfied until the whole thorax has been examined for signs of disease.

The onset of inflammation of the lung is especially apt to

be accompanied by signs of disturbance of the nervous system : indeed, in a young patient an illness beginning with high temperature, delirium, and rapid breathing, will in all probability prove to be a case of croupous pneumonia, and such a combination of symptoms should at once suggest this disease to our minds.

It is not always, however, that the beginning of the illness is accompanied by such violent phenomena. The onset is always abrupt ; but the early symptoms may be mild and increase only gradually in intensity. Thus, a child who is apparently in his usual health stops in his play, and, saying he is tired, asks to lie down. He then becomes thirsty and feverish, and usually vomits. In a few hours or the next day he begins to cough—a short, hacking cough ; his bowels are confined, his tongue is thickly furred, and he often complains of headache. The cough is generally accompanied by some pain, for, if old enough to speak, the child will often complain of pain in the side or belly, or will show by the distress in his face that the cough is painful to him. If the temperature be taken within a few hours of the attack the thermometer is found to mark 103° or 104° , or even higher, and it retains this level, with slight morning intermissions until resolution takes place, when the mercury falls as abruptly as it rose, and generally remains for some days afterwards at a level below the ordinary temperature of health.

At the same time that the cough is first heard the breathing begins to be hurried, and the nostrils dilate at each inspiration. The breathing is quickened out of proportion to the pulse ; and it is important to notice this point, because the altered relation between the pulse and the respiration is a valuable diagnostic sign, and when present with a high temperature would lead us at once to suspect the nature of the complaint. Instead of one breath to every four beats of the pulse, which is the ratio of health, there is one to three, or even sometimes one to only two.

This symptom is especially useful because it occurs early, and at a time when there may be no physical signs about the chest to lead us to a conclusion. About twenty-four hours after the first seizure the child is seen with flushed cheeks, burning skin, and red dry lips, which have often a patch of herpes. His breathing is quick, and its rhythm is often altered, the pause taking place at the end of inspiration instead at the end of expiration. With each inspiration his nares dilate widely. At frequent intervals he coughs a short hack, or has a succession of short hacks, which may last without intermission for five or ten minutes, leaving him pale and exhausted. The pulse is quick, varying from 130 to 150, and is often very unequal in the force of its beats, sometimes actually intermittent. The respiration is 50 or 60 in the minute, or even higher. But there is this peculiarity with regard to the breathing, as compared with the respiration of bronchitis, or of the other form of pneumonia called catarrhal pneumonia—viz., that however quick it may be, and however distinctly the lividity of the mouth and eyelids may prove the insufficient oxidation of the blood, the breathing is never laborious. None of the accessory muscles of respiration are brought into action; in fact, in the true sense of the word there is no dyspnoea. If there be much pain in the side the breathing is especially shallow, and Traube has attributed its rapidity partly to this cause and partly to the influence of the high temperature of the blood upon the nervous centres, for its quickness is found to be diminished by the application of cold. Still, although the child does not exhibit any profound consciousness of distress from the interference with the function of the lungs, yet he distinctly shows that his volition is now concerned in the act of respiration, and that the supply of air to the chest requires all his attention. If an infant, he sucks hurriedly, stopping every now and then to breathe with open mouth, as the nasal passages can no longer admit air in sufficient quantity to the lungs. Under examination,

too, he is quiet and docile, submitting himself without resistance to be percussed and ausculted, for he feels that crying and struggling would still further hurry his breathing, and this he cannot afford. This passiveness during examination of the chest is a valuable sign, and is characteristic of some interference with the function of respiration, whether that interference be due to disease of the lung, as in bronchitis and pneumonia, or of the chest-wall, as in rickets. In any case where an infant screams loudly during an examination of the chest, the probabilities are strong against the lungs being seriously diseased.

An older child lies quietly in bed, resting on his back or side. He takes little notice of what is going on around, and often refuses to answer the questions which may be put to him, for cases of pneumonia are distinguished by great muscular prostration, when all exertion becomes troublesome.

Examination of the chest does not at once reveal signs of lung consolidation. For the first day or two there may be no dulness, and by the stethoscope we may only hear weak or harsh breathing, if, indeed, the respiration be not perfectly natural. There is almost always sonoro-sibilant rhonchus diffused more or less widely over the lung. It is often not until the third day that positive signs of solidification can be detected, and even these may not all appear together. Thus, if the inflammation begins in the centre of a lobe, we may get bronchial breathing, with a puff of fine crepitation at the end of inspiration, but, should a layer of healthy lung-tissue intervene between the diseased spot and the chest-wall, there may at the same time be no dulness on percussion with one or even with two fingers. It is always important, as has been before remarked, to strike with two fingers as well as with one in examining the chests of infants and children. Percussion with one finger alone often yields negative results, when the use of two fingers will at once reveal the existence of dulness; and this practice is of especial importance in the disease.

under discussion, for in pneumonia the dulness is often far from being complete, and may easily escape notice if one finger only be made use of.

The sequence of the physical signs is much the same in infants and children as it is in the adult. At first the percussion is normal, and the respiration either natural or weak and harsh, accompanied by sonoro-sibilant rhonchus. In a day or two the percussion-note becomes more or less dull, with slightly increased resistance, the respiratory movements of the affected side are hampered, and the intercostal spaces sink in during inspiration. The respiration is weak or harsh or slightly bronchial, and fine crepitation is heard at the end of inspiration. The crepitation is perhaps coarser than the same sign as it occurs in the pneumonia of the adult, but its coarseness has been much exaggerated. At this period there is often a bronchitic subcrepitant rhonchus heard both with inspiration and expiration, and this, in ordinary gentle breathing is all that can be distinguished. It is only when the child takes a deep breath, as he occasionally does after a fretful whine or after a cough, that any true crepitation is heard at all, and then a puff of the true pneumonic crepitus can be caught at the end of the inspiration, and is found to differ little in its character from the same crepitation so familiar to us in the pneumonia of the adult.

The dulness then becomes more complete, the sense of resistance is greater, the crepitus ceases to be heard in the affected part, although it may still be detected at its confines, and the respiration assumes that intense sniffling, metallic character which is called tubular breathing; at the same time the voice of the child, or, in an infant, the cry, is conveyed with unnatural distinctness to the ear.

The situation of the physical signs is important. They may be found, as has been said, at any part of the lung—at the apex as often as at the base—or may be seated at some part of the middle lobe. As a rule, however, they are

limited—at any rate at first—to the posterior regions of the thorax, and unless the inflammation occupy the apex of the lung, are rarely to be detected in front. This is valuable, as serving to distinguish this disease from pleurisy, which it so much resembles, and with which it is so often confounded.

The physical signs, after persisting for a variable number of days, begin gradually to subside; and in a typical case of pneumonia the order of the changes is as follows:—Crepitation first returns, coarser and more distinctly bubbling than before. The respiration then becomes less sniffling, and gradually loses its bronchial quality. The dulness also diminishes, and finally disappears. The cry remains too resonant for a considerable time, persisting so long as there is left any dulness on percussion; and for some days a fine bubbling rhonchus, the remains of the redux crepitation, is heard at the affected part. The above phenomena are not, however, constant. Resolution is often marked by no returning crepitation, and the consolidation may be removed without other rhonchi than ordinary dry sonoro-sibilant râles being at any time detectable. This, indeed, is far from being an uncommon circumstance. The dulness is also sometimes very slow to disappear, and for many days, and perhaps weeks, after the return of normal resonance, the breath-sounds may be harsh and feeble at the affected part, and may only gradually re-assume the characters of health.

The symptoms during the course of the disease undergo little change. The child usually wanders a little at night, and starts and moans in his uneasy sleep. He seldom cries, having no breath to spare for noisy lamentations, but he sometimes emits a kind of grunt or moan at the end of each inspiration, which has been said by some writers to be characteristic of the disease. The flush on the face generally disappears after a few days and the cheeks remain pale, while the eyelids, lips, and sometimes the tips of the

fingers, become livid. The urine is scanty, and is thick from lithates ; it contains a large amount of urea, but little chloride of sodium, and there is often a trace of albumen. In many cases, however, the termination of the stage of exudation is marked by an evident improvement in the general condition of the child ; for although the temperature does not fall, yet certain changes, slight in themselves, are noticed at this time, showing that the destructive stage, if it may be so called, is at an end, and that recovery is shortly to begin. The expression of distress passes away, the child is less absorbed in his own sensations, loses his preoccupied look, and begins to take some interest in what is passing around him. The first symptom indicating that resolution has begun is a sudden fall of the temperature, which sinks to a level below that of health, and the thermometer often marks only 97.6° in the rectum. This fall generally precedes by about twenty-four hours, or even longer, any improvement in the physical signs. At about the same time the cough, which for some days had been less hacking, becomes quite loose, and there is often a profuse general perspiration. The pulse and respiration diminish in frequency, and return to their natural relation to one another. This change takes place in most cases on the fifth or seventh day, unless more than one lobe of the lung has been involved in the inflammation, in which case the crisis may be delayed until the eleventh, twelfth, or thirteenth day. The child is left very pale and prostrate, but the appetite quickly improves, and unless a relapse takes place the patient soon regains his flesh and strength.

The above may be called the normal course of simple primary croupous pneumonia, but we occasionally meet with variations in the symptoms which it is important to be acquainted with, as they may obscure the diagnosis.

The disease is sometimes ushered in with convulsions : and this is not a very uncommon mode of beginning. The convulsions may last for several hours, the child passing from

one fit into another. The cases where this symptom occurs need not be of unusual severity, nor is the evolution of the disease necessarily marked by any other sign of cerebral disturbance. The convulsions generally cease after a few hours, and the disease runs its usual course without any return of the fits, and without the occurrence of any circumstance calculated to excite unusual anxiety.

CASE V.—PNEUMONIA INTRODUCED BY A CONVULSIVE
ATTACK.

William A., aged 5 years, was seized with convulsions on November 28, which lasted for a quarter of an hour, and left him in a half-conscious state for several hours afterwards. He was seen on December 1, lying in bed with a flushed face, intensely hot skin, and loaded tongue. His bowels had been confined for two days, and he complained of frontal headache and of pains about the chest and belly. The pulse was 140, and the respiration 75. He had a frequent dry hard cough, and the nares acted with each inspiration. His nose had bled shortly before the visit. On examination of the chest, resonance was normal, and the respirations appeared quite natural both at the back and at the front.

On the next day, December 2, there was incomplete dulness at the right back, with bronchial respiration and some crepitation; the pulse was 138; the breathing was rapid, but could not be taken on account of irregularity from straining and sighing. He wandered at night, and lay dozing during the day. There were sordes on the lips, and his face was pale, with a tendency to lividity; but still he lay back in bed without any manifest oppression of his breathing. In this case the temperature fell on the seventh day, and at the same time redux crepitation was heard at the back. On the sixteenth day examination of the chest

showed no signs of disease, and the child was pronounced convalescent.

The epistaxis which occurred on the third day in this child is to be noted, because it is a symptom which is sometimes found early in the disease, and it is stated to be also occasionally seen at the period of resolution. The course of the disease was mild : the pneumonia was limited to one side, and the initial convulsion did not exercise any unfavourable after-influence upon the case.

The convulsions do not necessarily cease as the disease becomes developed, but may be repeated again and again. In the intervals of the fits the child may be drowsy and stupid, and pass his water and motions in his cot. I have even known a temporary squint to be left as a consequence of the convulsive attack. Such a symptom, combined with headache and drowsiness, might very possibly excite great alarm, especially in the absence of decided physical signs of pulmonary inflammation.

Convulsions do not always form part of the nervous symptoms met with at the beginning of the illness. In some cases drowsiness, approaching to stupor, may be noticed ; or there may be vomiting, with violent headache and delirium. The combination of headache and delirium is, indeed, far from uncommon at the beginning of a pneumonic attack in the child. If the cerebral symptoms are combined with a short hacking cough, active nares and a perverted pulse-respiration ratio, we should not overlook the probability that the patient is suffering from pneumonia even before the presence of consolidation can be ascertained. Special symptoms pointing to the lung are, however, occasionally absent. Instead of hacking cough and pain in the chest we find headache, vertigo and delirium or stupor,—symptoms all pointing to cerebral lesion and distracting our attention from the real seat of disease. On account of the predominance of brain symptoms, this form of inflammation of the lung has been called “cerebral pneumonia.” It is often a

source of considerable perplexity to the practitioner, for an examination of the chest may not at once discover the presence of pulmonary consolidation. The signs of local inflammation may be slow to appear, and, when at last present, may be curiously limited in extent. Therefore, unless the whole of the chest be subjected to repeated examination, the real nature of the disease may be misapprehended.

The following is a good instance of this form of pneumonia :—

CASE VI.—PNEUMONIA OF APEX WITH SEVERE CEREBRAL SYMPTOMS (CEREBRAL PNEUMONIA).

Ellen S., aged 2 years and 8 months, a fat, robust-looking child, after ailing for a week, began to be feverish, and had a slight cough. She was very restless at night, twitching, and sometimes starting up from her sleep with a scream. She refused food, but constantly required drink. From the beginning of her illness she had suffered much from vertigo. While lying or sitting on her mother's lap she would suddenly clutch at her mother's neck or dress, and exclaim she was falling. This, for two days before the visit, had been a very prominent symptom. The child was first seen on the morning of February 4, three days after the cough had been first noticed. "Looks ill; nares not acting; face not flushed, but is said to flush at times, and then to turn pale. Pressure on forehead produces no vivid redness, and when the finger-nail is drawn along the cheek a reddish line appears slowly, and is not brighter in colour than is compatible with health; muscles of face twitch. Tongue rather dry; and covered with a thick layer of yellowish-white fur. Skin feels hot to hand. Temperature in axilla before examination of chest, 101.8° . Pulse 156, irregular in force, but not in rhythm: it was

difficult to take, on account of subsultus tendinum. Respirations regular and not hurried. Belly not retracted, rather full, if anything, and seems a little tender; bowels confined two days; has not vomited during whole time of present illness."

On examination of chest no dulness was found anywhere—back or front, and with the stethoscope respiration appeared to be natural, and nothing abnormal was found but a little sonoro-sibilant rhonchus here and there over both lungs. During all the time of examination the child kept up a constant scream, and, when the examination was concluded, lay drowsily back with closed eyes in her mother's arms. A rhubarb and jalap powder was ordered, with a saline mixture.

On February 8 she was seen again. "Ever since last visit has been lying in a drowsy state, with eyes half closed and dim-looking, taking no notice of anything that passed. Yesterday, however, she cried once or twice to be nursed, which she had not done before. Is now (10 a.m.) lying in her mother's lap, apparently half stupefied. The eyes are half closed, showing the whites. When forehead is pressed she looks up, and appears as if about to cry, but closes her eyes again almost immediately. No very positive redness produced by pressure on forehead or cheek. Lips still twitch; and yesterday her mother thought the child was going to have a fit, she twitched so much. Is said sometimes to squeal out sharply. Does not squint. Vertigo still continues, for child seems to be occasionally afraid of falling. Temperature in axilla, 101°; pulse, 152, very intermittent in force and rhythm. Tongue thickly furred and rather dry. There has been no sickness, and the bowels act every day. The child has still a little cough, and seems rather hoarse." Unfortunately, the chest was not examined on this visit, principally from a wish not to disturb the child unnecessarily. She seemed exceedingly ill, and her case was strongly suspected, in spite of the contra-

dictory symptoms, to be one of tubercular meningitis. Some calomel powders were ordered.

On the next visit, February 11,—“Until yesterday the child ‘lay as if dying, with hands drawn up.’ Since then has been very fretful, asking constantly for drink, and moaning and whining. Eyes now bright. Pulse excessively intermittent in force and rhythm; can’t be counted from restlessness. Has not squinted. Bowels, since taking the powders, have been opened two or three times a day; motions green and slimy. Is hoarse; coughs a great deal; and there is some rattling of phlegm about the chest. Retches sometimes after the cough, but she has not vomited all through the course of the disease. Temperature in axilla before examination of chest, 99.4° . Percussion of the right back showed dulness from the supra-spinous fossa to half way between the spine and the lower angle of scapula; respiration there very bronchial and almost tubular; no crepitation; vocal resonance increased.” A draught containing ipecacuanha wine and sal volatile was ordered every four hours.

On February 15,—“Child has been improving since visit: has been more lively. Bowels have acted about every other day; motions dark and offensive. Is still hoarse, and requires much drink. At night is fretful, but in the daytime lies quietly enough if nursed. Appetite very bad. Temperature in axilla, 99° . Pulse can’t be counted on account of restlessness. There is still dulness at right back, as before. The child cries loudly all the time of examination, and the transmitted hoarse laryngeal breathing makes it impossible to tell the condition of the respiration.” Half a grain of quinine was ordered three times a day.

After this date the appetite became good, and the child very rapidly improved; on February 25 the signs of consolidation had almost entirely disappeared.

In the above case there was little to draw attention to

the chest. Instead of flushed cheeks, active nostrils, rapid breathing, and frequent hacking cough, the face was pale, the nares were motionless, the cough slight, and the breathing quiet; while the vertigo, the twitching, the marked drowsiness persisting for seven days and becoming intensified into something approaching to stupor, all seemed to point to the existence of cerebral lesion. In spite, however, of the symptoms, there were many points of distinction between this case and tubercular meningitis, for which it was almost mistaken. The short duration of the premonitory symptoms, in the first place, argued against such a supposition; for although tubercular meningitis occasionally begins with suddenness, yet it is usually preceded by gradual loss of flesh, colour, and spirits, with slight feverishness, constituting a *malaise* which persists for several weeks before any regular symptoms of the disease are noticed. Again, the disease was not ushered in by sickness, and the constipation was slight and easily removed by an ordinary aperient; the pulse was rapid, while during the earlier periods of tubercular meningitis it is generally slow; the abdominal wall was not retracted; the respiration was regular, without sighing or the occasional pauses which are so marked a feature in the cerebral lesion; there was no flushing on pressure of the skin, no squinting or irregularity of pupils; and the child, although lying in a state of apparent unconsciousness, could be easily roused.

Mere irregularity of pulse is of absolutely no value as a means of diagnosis. In any febrile attack the pulse of a young child may become irregular; indeed, in many cases the mere quickening of the heart's action seems to lessen the regularity of its contractions. But while a *quick* irregular pulse is of little consequence, a *slow* irregular pulse carries a far different meaning, and, if suspicions of tubercular meningitis have been already excited, will go far to convert those suspicions into certainty.

The degree of heat of the skin, which was moderate for

inflammation of the lung, told rather in favour of meningitis, but it must be remembered that the child was seen in the morning, when the temperature of pneumonia falls to a certain extent. If it had been taken in the evening the thermometer would no doubt have been found to rise as high as 103° or 104° . Again, the impatience of the child under examination rather opposed the notion of pulmonary lesion ; but in cases where the pneumonia attacks the apex, a child is usually found to be less tractable than when any other part of the lung is the seat of inflammation. In health the apices are less freely expanded in ordinary inspiration than are the other parts of the lungs, on account of the deficient movement of the chest-wall in the superior regions ; and therefore in disease the patient suffers less from the temporary incapacity of a part which takes naturally only a moderate share in the respiratory function.

Instead of a cerebral lesion we might have been inclined at the first to suspect the existence of suppressed exanthema ; but the continuance of the stupor, and the absence of any more special symptoms, soon excluded this idea.

In this case the apex of the lung was the part affected, and it is often taught that such severe symptoms are practically confined to cases where the inflammation occupies this region of the lung. My own experience, however, is not in accordance with this statement. I have met with nervous complications as often in pneumonia of the base as in pneumonia of the apex of the lung. Indeed, when the upper part of the organ is attacked by the inflammation, the course of the disease, as a rule, is exceptionally mild. Thus :—

CASE VII.—RAPID COURSE OF PNEUMONIA OF THE APEX.

Thomas F., aged 4 years, a strong-looking, fat boy, was taken ill on May 25. He complained that he felt tired, and wanted to lie down. He was thirsty, and vomited directly after drinking. In the night he was very feverish, and coughed a great deal.

He was seen on May 28. "Face flushed ; nares acting ; has frequent hacking cough ; tongue furred ; and bowels confined. Respiration 44 ; pulse 128, very irregular ; temperature in axilla, 104.2° . Examination of chest shows dulness at right apex, principally at the supra-spinous fossa, where the breathing is tubular (with the mouth open) ; vocal resonance is unusually strong, and there is a little subcrepitant rhonchus, but no true crepitation. In front there is only slight dulness at the supra-clavicular region, and the respiration is bronchial." In this case the temperature fell on May 30 (the fifth day) ; and on the following day,—“Skin quite cool ; respiration natural, and not quicker than ordinary. A little dulness left at right supra-spinous fossa, but respiration merely harsh, and there is a little subcrepitant rhonchus with both inspiration and expiration.”

In this case there is nothing unusual. It is a good example of the ordinary course of pneumonia of the apex, such as may be seen every day, during the changeable seasons of the year, in the out-patients' room of a children's hospital.

Besides the nervous symptoms which are so often a cause of perplexity at the beginning of an attack of pneumonia, other complications may arise and occasion some difficulty in the diagnosis. It may happen that an attack of gastro-intestinal catarrh accompanies the onset of the inflammation. In such a case we may suspect the outbreak of typhoid fever, and indeed at a first glance the resemblance to enteric fever may seem to be a close one.

CASE VIII.—PNEUMONIA COMPLICATED BY GASTRO- INTESTINAL CATARRH.

Caroline H., aged 11 years, was taken ill on the evening of December 26, with frontal headache and vomiting of bitter fluid. Her skin was hot, she was very thirsty, had

no appetite, and complained of feeling very ill and weak. On the following day (December 27) her bowels became relaxed, and acted four or five times before the evening. She remained in this state till December 30, when I first saw her.

“Complains of frontal headache; skin intensely hot; tongue moist and covered with whitish fur, through which red papillæ project. Is very thirsty, and complains much of weakness; wanders at night, and got out of bed last night while delirious. Bowels acted five times yesterday; motions dark-coloured and slimy. Says some tenderness when pressed in left iliac fossa, and there is gurgling there and on the right side. She coughs a good deal, but lungs on examination appear quite healthy.” Small doses of acetate of lead and opium were ordered.

On January 1,—The diarrhœa was better, although the bowels had acted twice previous to visit. “Pulse 128; skin still intensely hot; wandered less last night, but cough was very troublesome and prevented her sleeping. Complains of great pain in left hypochondrium. There appears to be some want of resonance at both bases posteriorly; respiratory sounds a little harsh on left side, and sonoro-sibilant rhonchi are heard over back.”

On January 3,—“Bowels have not acted since last visit. Pulse 120, full but compressible, regular. The patient seems a little delirious; she talks loudly, describing her symptoms, with a peculiar ringing lengthening of the termination of her words. Pupils dilated, eyes glistening. Complains of pain at the top of her head and in the left side of her chest, which last she compares to ‘knives and forks’ sticking into her. Tongue still furred; cough very troublesome; skin still very hot. She was very delirious last night; tried to get out of bed, and talked wildly. No tenderness about abdomen, which is not at all swollen, and there is no appearance of eruption either there or at the back. On examination of back there is dulness of left base

reaching up to the angle of scapula, with bronchial breathing, a little subcrepitant rhonchus accompanying both inspiration and expiration, and fine crepitus at the end of deep inspiration. Vocal fremitus is increased. In front of left side no physical signs are found."

On January 5 (the ninth day) the pulse had fallen to 80, and the heat of the skin, as tested by the hand, seemed very much diminished. She slept well, was not delirious, and altogether seemed very much better, although very weak. On the next day (January 6) the chest was covered with sudamina, and on January 7 redux crepitation was heard all over the affected lung. She then became rapidly convalescent.

In this case the frontal headache, the diarrhoea, and the tenderness of the abdomen, combined with fever and delirium, might have given rise to suspicions of typhoid fever; but the mode of beginning of the two diseases is not quite the same. In primary croupous pneumonia the onset is more abrupt and violent than is commonly seen in the case of typhoid fever in a young patient. Often in the latter disease the exact day of beginning is difficult to ascertain, or if, as sometimes happens, the fever began without warning, the early symptoms are usually mild. The child complains of headache, can eat no dinner, and is languid; but it is rare for him to appear at that time to be seriously ill. In the case of croupous pneumonia, whatever the symptoms may be, the bodily prostration from the first is usually well marked. Again, in the present case the girl was light-headed almost from the first, and on the second night after the attack attempted to leave her bed while in a state of delirium. In typhoid fever the delirium begins later, and is seldom found before the end of the first week. Besides these two very important points, the prominence of the cough from the first, and the altered relation between the pulse and respiration, would lead us to expect the occurrence of the characteristic physical signs of lobar pneumonia.

Later, the non-appearance of the spots, which are seldom absent in a child of 11 years old, and the early convalescence,—for in typhoid fever the temperature rarely falls before fourteen days have elapsed,—confirmed the diagnosis.

Although in a healthy child who has passed the age of infancy the occurrence of gastro-intestinal catarrh at the beginning of the inflammatory attack is often an accident of slight moment, in an infant the presence of this complication is a serious misfortune. As a rule, primary croupous pneumonia, when it attacks a baby of six months old or upwards, is a disease of little gravity. The symptoms are mild and the temperature falls early. But the illness has this inoffensive character only as long as it is primary and uncomplicated. If it occur in a child who is under-nourished, or already weakened by digestive disturbance, the consequences may be serious. So, also, if the onset of the inflammation be accompanied by any derangement which tends to reduce the patient's strength, there is room for great anxiety. Our apprehensions are especially justifiable when this derangement takes the form of diarrhoea; for in such a case, if the purging be at all severe, the child will probably die.

CASE IX.—PNEUMONIA OF THE LEFT LUNG IN AN
INFANT : GASTRO-INTESTINAL CATARRH.

Frank K., aged eleven months, without any teeth, was one of twins. There was no history of phthisis in the family, but the father had died of pneumonia. The mother was healthy, and although the birth of the twins had been preceded by five miscarriages, neither of the children had snuffled or shown any sign of syphilis. They had been suckled for the first three months, and had then been brought up on Swiss milk. The patient's appetite had been failing for a few weeks, and his bowels had been relaxed. A few days before admission he had begun to cough.

When first seen the child was thin and pale. There was no beading of his ribs or enlargement of the epiphyses of his long bones. His fontanelle was large and flattened, and he had a little thrush in the mouth. The child's breathing seemed rather oppressed, and a little recession of the epigastrium was noticed in inspiration. Resonance was diminished all over the left back, and the breath-sounds there were high-pitched and blowing. There was no crepitation. The laryngeal sounds were conducted very plainly to the base of the lung on the diseased side. The right lung appeared to be healthy. The heart-sounds were clear. The temperature varied in the twenty-four hours between 101° and 104° . The bowels were relaxed, acting four or five times in the day, and the purging, although not in itself of a dangerous character, seemed to be little influenced by treatment. The child took at first fifteen grains of bismuth with half a minim of laudanum every four hours; then oxide of zinc with opium. The purging, however, continued, and in spite of stimulants, the infant died quietly at the end of a week. Shortly before death his temperature was 100.4° .

In this case the attack of pneumonia occurred in a child who was no doubt under-nourished, for Swiss milk, unless largely supplemented by more substantial and nutritious food, forms a very inappropriate diet for a child of eleven months. In addition, a gastro-intestinal derangement had accompanied the onset of the inflammation. The purging, although obstinate, was not very profuse, and if the lungs had been sound, would probably have given rise to but little anxiety. So, also, inflammation of the lung, if limited to one side of the chest, might be expected in a healthy baby to undergo speedy resolution. Unfortunately, the co-existence of the two complaints caused an amount of depression which the patient's natural strength proved insufficient to overcome.

An infant or young child is more dependent than an

older subject upon a daily supply of nourishment, for the needs of growth and development have to be satisfied, as well as the ordinary wants of tissue-repair : indeed, at this period of childhood, impoverishment of the blood amounting to actual anæmia is a common consequence of any derangement which interferes, for however short a time, with the digestion and assimilation of food. Disturbance of the organs concerned in nutrition is therefore in early life one of the most serious of complications. At a time when the functions of life are already hampered, and the needs of the system are most urgent, the accustomed supply of nutritive material comes to a sudden standstill. Consequently, the enfeebling influences of the disease are allowed full play, and unless the digestive organs right themselves speedily, life comes to a close from simple exhaustion.

Although the fall of temperature usually takes place on the fifth or seventh day, it is occasionally, as has been said, deferred to a later date. This postponement of resolution is often found when a large tract of lung has been involved in the inflammation. Sometimes, however, the prolongation of the pyrexial period is due to the invasion of successive tracts of lung, and when this occurs, the temperature, instead of retaining nearly the same level through the disease, undergoes occasional exacerbations, each of which is followed by a slight fall, so that the pyrexia may appear to be intermittent. Sometimes a period of several days will pass between the first fall of temperature and its after-rise ; in these cases a relapse is said to take place when the new portion of lung becomes invaded, but this second attack seldom lasts so long as the first, and the temperature usually falls again on the third or fourth day.

It may happen that the extension of the inflammation from one part of the lung to another is associated with nervous symptoms such as are apt to accompany the onset of the disease ; indeed, I have known signs of nervous dis-

order, which had been absent at the beginning of the attack, to occur for the first time during such extension. Thus, a little boy, aged three years and a half, got wet through on February 25th, and in the evening was seized with shivering. The next day he coughed a great deal, and his face was noticed to flush. On the 28th, at the hospital it was discovered that he had pneumonia at the right base behind. On March 3rd, the lower half of the lung was clearing, but the upper half of the same side was found to be dull on percussion and the breath-sounds to be tubular. There was no crepitation. During this and the following day the child was very drowsy, and passed his water and motions in his cot. He only took food when roused to do so. Temperature 101° ; pulse 144; respirations 80. The nervous symptoms soon passed off, but the progress of the disease was unusually slow, and it was not until March 18th that all physical signs of consolidation had disappeared.

An attack of pneumonia, if the disease be uncomplicated and occur in a child whose nutrition is satisfactory, need seldom be a cause of grave uneasiness. Inflammation of the lung, indeed, may be said to be one of the varieties of illness which usually run a milder course in early life than they do in after years. Even in cases where the onset is accompanied by the most violent nervous symptoms, the danger, as a rule, is more apparent than real. I do not remember ever to have met with a case of simple primary croupous pneumonia, beginning with convulsions or other form of nervous disturbance, which ended otherwise than favourably. But while the uncomplicated disease is so little dangerous to life, the presence of any associated disorder may be a matter of very serious moment. In infancy the exhausting effects of a gastro-intestinal catarrh have been already referred to; and even in later childhood the existence of purging, if the drain be at all profuse, should be regarded with apprehension. An intestinal derangement complicating a case of pneumonia often resists treatment with some obstinacy; and its occur-

rence can never fail greatly to reduce the strength of the patient, and delay the period of convalescence.

In croupous inflammation of the lung the symptoms of general disturbance and the signs of local mischief bear no necessary proportion to one another. The disease may be ushered in with the most violent nervous commotion, although the area of lung affected by the inflammation is curiously restricted. Nor can the energy of the onset be taken as any guide to the probable duration of the fever ; for the temperature may fall and resolution be completed as early in cases where the initial symptoms are severe as in cases where they are mild. Thus, in the case of a girl, aged thirteen years, in whom the disease began with convulsions, followed by squinting and delirium, the temperature became normal on the seventh day, and the patient was convalescent at the end of a fortnight. Resolution in croupous pneumonia is seldom otherwise than complete ; and it is rare for an unabsorbed deposit to be left as a consequence of this form of inflammation of the lung. From the foregoing remarks it may be concluded that in the primary uncomplicated disease the prospects of the patient, both immediate and remote, are most favourable ; but the prognosis is more serious when the pneumonia occurs in a weakened or unhealthy child. A secondary pneumonia is a very grave disease, and often proves fatal. The weakened state of the patient increases the danger of failure of the heart's action, and in infants, especially, it is rare for the illness to be recovered from.

Croupous pneumonia, as a rule, is not difficult to recognise, even before physical signs have appeared. Head-ache, sickness, a high temperature, and, in the course of a few hours, delirium, form a very suspicious combination of symptoms. If, in addition, the nares act, the pulse-respiration ratio is perverted, and there is a hacking cough, we can have little doubt that signs of acute consolidation of the lungs are about to manifest themselves. Not unfrequently, however, cough is absent at the first, and the normal relation of the

pulse to the respiration may seem to be little altered. Still, even without these symptoms, we can often arrive at a correct diagnosis. A high temperature, with a dry parched skin and early delirium, occurring in a child whose health has been previously satisfactory, is very suggestive of pneumonia. Marked delirium is not common in early life ; for although a feverish child is often light-headed, he wanders quietly and exhibits no symptoms calculated to excite alarm. But the delirium of pneumonia is usually noisy and energetic, and causes much anxiety. In every case, then, where a child who is suddenly seized with a feverish illness shows early and marked delirium, we should at once think of pneumonia, and look closely for symptoms pointing more particularly to the lungs. It must be remembered that examination of the chest may not furnish corroborative evidence until several days have elapsed, for the physical signs of consolidation are often slow to appear ; also that a violent onset may be followed by signs of very limited solidification of the lung. Therefore we should subject every part of the chest to careful examination, and in the majority of cases we shall find some local signs at the end of the second, or, at the most the third, day from the beginning of the disease.

When the physical signs of consolidation are developed late in the illness, the delay usually occurs in cases where the nervous symptoms are especially pronounced. On this account the beginning of the disease in infants is often very puzzling to the practitioner, who is forced to reserve a positive opinion until some definite alteration in the physical signs can be discovered in the chest. The difficulty is increased by the frequent absence of cough in these young patients. If, then, the attack begins with drowsiness, passing into stupor, but without cough, without action of the nares, and without perversion of the pulse-respiration ratio, it is excusable in the absence of signs of pulmonary consolidation to feel great uncertainty as to the cause of the symptoms.

Still, the very fact that the drowsiness comes on abruptly in a child who had been previously in a state of good health is a suspicious one ; and if we notice that the temperature is high, with a dry skin, and that there are no signs of headache (for in pneumonia pain in the head rarely occurs when the stupor is pronounced), the true nature of the illness should suggest itself to our minds. In most cases an alteration in the physical signs will come sooner or later, to show that our suspicions are well-founded.

After the physical signs have appeared there is no excuse for misapprehension. The dulness, incomplete as a rule, with no very great increase in the sense of resistance ; the intensely tubular breathing ; and the resonance of the cry—these signs, heard principally posteriorly, and only in a very limited degree at the front of the chest, are very characteristic, especially when combined with the extremely elevated temperature and the altered relation between the pulse and the respiration.

No mention has been made of vocal vibration of the chest-wall—a sign which is so valuable in the adult—because in children the presence of chest fremitus is very uncertain, and cannot be relied upon. In some children it may be completely absent from both sides of the chest. In others it is absent from the sound side, but is felt faintly over the affected lung. In other cases, again, the vibration of the chest-wall has seemed to be actually diminished on the diseased side as compared with the opposite half of the chest. In cases where a fremitus is felt *only* over the diseased side, the sign is an important one, and points distinctly to consolidation.

There is a certain combination of symptoms which might prove embarrassing to a beginner. Thus, if a child be suffering from slight bronchial catarrh, and at the same time be feverish from cutting a tooth, the temperature may reach 104°, or even higher, and this, with hurried breathing and a quick pulse, might give rise to strong suspicions of pneu-

monia, particularly as on examination of the chest nothing might be found but a little dry, wheezing rhonchus. In such a case, however, the pulse-respiration ratio would be little perverted. The character of the cough, too, which has not the short hacking quality so characteristic of inflammation of the lung, would be a valuable distinctive sign, and one which should make us search further for a cause of the pyrexia. This would be at once discovered by an examination of the mouth, and the history of the attack would remove any remaining hesitation as to the nature of the ailment.

The diagnosis between pneumonia and pleurisy will be considered in treating of the latter disease.

An ordinary case of primary croupous pneumonia requires little medicine. The treatment should be directed rather to ward off disturbing influences than by injudicious meddling to interfere with the course of a disease the tendency of which is to recovery.

The patient should be placed in an airy, well-ventilated room, which must not be kept too hot. Some febrifuge medicine may be given, such as *liq. ammon. acetatis*, with citrate of potash, so as to favour the action of the skin, and this object is aided by the addition of five or ten drops of *ipecacuanha* or antimonial wine to each dose of the mixture. The affected side must be kept constantly covered with large hot linseed-meal poultices. These poultices must be made large, so as to overlie a sufficient extent of surface, and they should be thick, so as to retain the heat as long as possible. They can be changed every two hours. If there is much pain in the side, a weak mustard and linseed poultice (one part of mustard to five of linseed-meal) may be used, and in this strength can be retained in contact with the skin for at least six hours. Heat applied in other forms, as in the shape of bags of heated bran or salt, is very useful; and dry heat has the advantage of being more manageable than the ordinary poultice. The child should be kept quiet in bed, and

every effort should be made to anticipate his wants, especially in the matter of drink, so that he may be spared the necessity for speech. Most children, indeed, will endure much discomfort rather than interrupt the respiratory function by complaints, so that their wants, unless anticipated, are apt to go unrelieved. The patient may take simple fluids in small quantities at frequent intervals ; and it is advisable to pour about an ounce of thin barley water or pure filtered water into a feeding-cup, and hold it to his lips every hour. A thirsty child may require drink more frequently. It is only necessary to allow time for a first quantity to be absorbed before a second is taken.

The bowels must be watched, for constipation would tend still further to increase the heat of the body ; and when necessary, therefore, the bowels must be relieved by small doses of castor oil or syrup of senna.

The diet is very important. Until the consolidation is complete the food should be limited to small quantities of milk and broth. If the patient be an infant at the breast, it is well to limit the number of times he is allowed to suck ; and this he will bear readily if some fluid substitute, such as thin barley-water or plain water, be given him at intervals with a spoon, for his desire for the breast is the result of thirst and not of hunger. As soon as the establishment of bronchial breathing and the disappearance of crepitation show that effusion into the air-cells has ceased, the diet should be improved ; but great care must be exercised in this respect, for while the patient is in a state of high fever his digestive power is necessarily very limited. The alteration should consist more in the frequency of his meals than in the quality of his food. Strong beef-tea may be substituted for weak broth, and this and the milk should be given to him at short intervals during the day and night. The milk should be guarded by the addition of small quantities of the saccharated solution of lime in the proportion of fifteen drops to each small teacupful. The yolk of an egg

oaten up with milk may be given once in the day if it is found to agree.

It is necessary in cases of pneumonia to watch the temperature, so that the bodily heat may not be allowed to attain an undue elevation. If the thermometer mark a higher level than 105° , it will be advisable to reduce the temperature by frequent tepid sponging of the surface ; and if the patient be not distressed by the fever, this simple measure will usually do all that is required. In cases, however, where the nervous system is evidently suffering from the intensity of the pyrexia, and the temperature is little reduced by the application, or rises again immediately, it will be necessary to employ cold in a more decided form. The child should be placed in water of the temperature of 70° Fahr., and allowed to remain in the bath until the thermometer shows a sensible reduction of heat. In these cases it is necessary to guard against the tendency to syncope, which in pneumonia is one of the accidents to be averted. A stimulant should be given both before and after the bath ; and when the child is returned to his bed, his feet, if cold, should be warmed by the application of a hot bottle. Another way of employing cold is to wrap the child in a sheet wrung out of cold water, and cover with a dry sheet or thin blanket thrown loosely over him. This treatment is often very successful in cases where the initial convulsions are violent and persistent, and are accompanied by a high temperature. By this means the abnormal heat is quickly reduced, and, as the temperature sinks, the nervous symptoms subside, and the child falls into a quiet sleep. Cold, so employed, is the best form of anti-pyretic. Aconite and other depressing drugs are not desirable remedies in a disease such as pneumonia, where sudden failure of strength is already a danger to be feared. In this respect quinine is a more useful drug to employ, for in addition to its anti-pyretic action, which it exercises without any tendency to weaken the heart, it is thought to have some influence in checking

the spread of the inflammation over the lung. To be of service quinine must be given in full doses. In early life the drug is very well borne, and even infants can take it in good quantity without any indication of distress. For a child of three years old three grains of the alkaloid may be given in a little glycerine every three or four hours. In this quantity it is rare to find any sign of cinchonism follow the absorption of the remedy.

As a rule, little physic is required in cases where the disease is primary. If the pain be severe, a small dose of Dover's powder (two grains for a child of five years old) may be given at bed-time ; and when the pyrexia subsides, quinine may be administered in ordinary tonic doses. Although the muscular prostration is apparently severe, it is not wise in an ordinary case to resort to alcoholic stimulation. The disease, if left alone, runs a short, sharp course, and stimulants, while the fever is high, neither mitigate the present discomfort nor hasten the period of convalescence. If, when the disease is over, the child be left unusually weak and pallid, a little alcohol in the shape of *vinum ferri* may be ordered ; but even this mild stimulant is not often required, for when the temperature falls, strength and colour are usually very quickly regained.

In the case of a secondary pneumonia, the great necessity is to support the strength, which has usually been much reduced by previous illness. Stimulants are often required early, and ammonia and bark may be given from the beginning. If the child is constitutionally weakly, a great danger may arise from temporary attacks of faintness. These seizures come on suddenly : the colour of the face turns to an ashy grey, the lips are blue, and all lustre leaves the eyes. Directly these symptoms are noticed, energetic measures must be adopted to restore the vigour of the heart's action, for unless this can be done the child will die. Strong brandy and water should be poured down the throat, vigorous frictions be used to the chest, and hot applications be applied

to the feet. In some cases the attacks recur again and again, so that the life of the patient is kept in continual danger.

The diet of the child suffering from a secondary pneumonia is to be carefully arranged, so as to supply plenty of nourishment without overtaxing the digestive powers. If the patient be old enough to digest it, beef essence or strong beef-tea should be given in small quantities, and repeated at regular intervals. Strong meat jelly may be used, and plenty of milk may be allowed, care being taken that the formation of large lumps of curd is prevented by the addition of some thickening material, such as gelatine or pearl-barley jelly. A good stimulant in these cases is the brandy and egg mixture of the British Pharmacopœia.

CHAPTER IV.

PLEURISY.

PLEURISY a more common disease in young children than is often supposed

—Symptoms—Physical signs—their peculiarities.

Case X.—An ordinary attack of pleurisy—Comment.

Case XI.—Circumscribed pleurisy—Comment.

Case XII.—Empyema—Absorption of the fluid—Comment.

Symptoms of empyema—Temperature—Modes of termination.

Case XIII.—Empyema—Spontaneous evacuation.

Pleurisy with retraction.

Case XIV.—Great distortion of chest from pleurisy.

Case XV.—Empyema—Secondary acute tuberculosis—Comment.

Case XVI.—Empyema—Extension of inflammation to the peritoneum.

Diagnosis—Prognosis—Treatment—Paracentesis—Mode of operation—

Effects of puncturing lung.

Case XVII.—Traumatic pneumo-thorax from puncture of lung during paracentesis.

Slight danger of paracentesis—After treatment—Diet.

PLEURISY is a very common disease in early life, and may be met with in the child at any age. In spite, however, of its frequency, the disease is one about which mistakes in diagnosis are often made. The symptoms may be so obscure that attention is not directed to the chest at all; or the physical signs may be so similar to those of pneumonia that the case is mistaken for one of inflammation of the lung. If we approach a case of pleurisy in the child with knowledge derived solely from observation of disease in adult patients, we shall be apt to fall into error; especially as the normal absence of vocal vibration of the chest-wall in early life deprives us of a valuable means of distinguishing a case of liquid effusion in the pleura from one of solidification of the lung.

With regard to the frequency of pleurisy, I find from reference to the records of the East London Children's Hospital that during the three years ending April 30, 1885, 2724 in-patients (medical and surgical) were admitted. Of these, 50 were cases of pleurisy, 48 of empyema, and 31 of pleuro-pneumonia : in all, 138, or one case of inflammation of the pleura in every twenty in-patients. These, however, it must be remembered, are hospital statistics. They relate to children of whom many are cachectic ; most are ill-nourished ; and nearly all are brought up under conditions little favourable to healthy growth. In well-to-do families we find a very different state of things ; for where children are well cared for and habitually guarded from harm, inflammation of the pleura is by no means a common complaint. It rarely happens, indeed, that a well-nourished child of sound constitution is attacked by this disease. Robust children for the most part escape : those who suffer from pleurisy are the cachectic and the frail.

The inflammation may be limited to one side of the chest or may involve the covering of both lungs. In a case of primary disease the affection, as a rule, is one-sided.

The onset of pleurisy, like that of pneumonia, is sudden, but, unlike the latter disease, it is rare to find the illness begin with violent nervous symptoms if the case be uncomplicated. The first complaint is usually one of pain in the side. The seat of the pain is not, however, always in the chest. The lower intercostal nerves are often affected, and the sensation being referred to the ends of these nerves where they ramify on the abdominal wall, the pain is often seated in the belly. This is important, for in children abdominal pain is apt to be at once attributed to worms or intestinal irritation. It is well, therefore, to remember that it may also be a sign of pleurisy, and in every case where it occurs a careful examination of the chest should be immediately made. The pain varies very much in intensity : it may

be very slight in degree or may amount to intense suffering. It is increased by coughing or deep inspiration, and is accompanied by much tenderness. In cases where the symptom is not strongly marked it may often be elicited during percussion, the child being seen to wince at the blow of the finger. In very young children the presence of pain in the side is usually shown by violent fits of crying, lasting for several hours. They show a great disinclination to be nursed or moved about, and when pressure is made upon the chest, as in lifting them up, their cries are immediately renewed. If the pain be a marked symptom its presence in older children is usually at once recognized by the expression of the child's face, which is seen to exhibit great distress when, from coughing or other cause, the child has been forced to draw a deep breath.

The breathing is rather hurried, but unless the disease be complicated by pneumonia the rapidity of the respirations is very much less than is the case in inflammation of the lung. On account of the pain the respiratory movement is very shallow, and the shallowness of the breathing may be a sufficient cause to determine its rapidity; for in cases where the pain, as judged by the other signs, appears to be severe, the breathing is especially rapid. Sometimes the nares act slightly during respiration, but this is by no means so constant a symptom as is the case in pneumonia. The pulse is quick at the first, as a rule, but this rapidity is seldom long maintained, and in most cases by the third or fourth day it has returned to its ordinary velocity. The pulse-respiration ratio is little perverted, and seldom falls below three to one. This is important with regard to diagnosis.

Occasionally the onset of the attack is indicated by vomiting, but this is not so common as is the case in pneumonia, where it is a very frequent symptom. There is generally headache, furred tongue, thirst, and loss of appetite, as in all acute attacks. Purging is sometimes a

symptom of the beginning, and occasionally the disease is ushered in by rigors or a convulsive fit.

The cough generally begins a few hours after the occurrence of pain in the side ; it is short and dry, but much less frequent than in cases of inflammation of the lung, and may be sometimes so slight as to escape notice altogether. If the pain be severe, the cough causes great distress, and the child is then seen to do all in his power to repress the cough, so as to avoid the necessity of drawing a deep breath. The degree of fever varies considerably in different cases. There may be absolutely no rise of temperature at all, or the evening temperature may be so little raised that the slight increase of heat escapes notice unless a thermometer be used. Even if a temperature of 102° or 103° be reached at the first, it seldom persists for a longer period than three or four days, when it begins gradually to decline. This *gradual* subsidence of the febrile heat is very different from the sudden fall of temperature which is so noticeable a phenomenon in pneumonia. It may happen, however, that the temperature remains high after the end of the first week, and shows no sign of declining. Such an unusual persistence of the pyrexia is no evidence that the fluid contents of the chest have already become purulent, for I have known the symptom to be present at the end of the second week, when a puncture of the thoracic wall has withdrawn a syringeful of perfectly clear fluid. Continued high temperature may be due to natural irritability of the patient, and is more common in scrofulous subjects than in others. In many cases, however, it indicates the presence of some complication. This is often pericarditis. Therefore in all cases where the high temperature persists after the end of the first week the possible occurrence of this complication should be kept in mind.

The quality of the heat is also a point to be noted. In pneumonia it is of a pungent character, which is very striking, and the skin is at the same time dry. In pleurisy, on

the contrary, the heat has not this pungent character, and the skin is sometimes moist, even in the early days of the disease.

On account of this frequent absence of fever, the face is seldom seen to be flushed, as in pneumonia. Indeed, a very characteristic symptom in pleurisy is the pale, rather sallow colour of the cheeks, which often at once attracts attention.

Another symptom which forms a great contrast to what we know of pneumonia is the comparatively slight degree to which the strength is reduced. In cases of inflammation of the lung the prostration produced by the disease is, as a rule, very great, while in the case of pleurisy, although differences are of course to be found in this respect, it is not at all uncommon for a child with its pleura half full of fluid to walk without assistance into the out-patient room. It is, perhaps, on this account that the child seldom looks very ill, and, unless disturbed by pain, his face has rarely the distressed expression which is so common in many other acute diseases.

The physical signs of pleurisy may resemble those of pneumonia very closely, and it often requires much care to distinguish by this means between the two diseases. The movement of the affected side is less free than that of the other from the very first. In the beginning this is no doubt the result of an instinctive effort on the part of the patient for the avoidance of pain ; but as effusion takes place into the pleura, the expansion of the chest-wall over the seat of disease becomes more and more constrained, and the immobility of the affected side during the respiratory movement presents a marked contrast to the rather exaggerated activity of the other half of the chest. Still, although the affected side of the chest moves less than the other in respiration, the intercostal spaces are not always motionless. In thin children they may often be seen to sink in to an appreciable extent at each inspiration even in cases of great effusion ; and this not

only laterally but also in the front and at the back of the chest. In observing this sign it is important to distinguish between real and apparent recession ; for some of the lower intercostal spaces—the eighth, ninth, and tenth—may appear to sink in through the action of the serratus magnus muscle, the digitations of which, contracting at each breath, elevate the surface under which they lie, so that the intermediate parts appear to be depressed.

If the effusion be large, displacement of the soft parts takes place ; and this is a very important sign.* The heart and mediastinum are pushed to the right or to the left (according to the side affected), the liver and spleen are slightly depressed, and the intercostal spaces become bulged and slightly widened. In these cases the whole side may become dilated, but the dilatation is often difficult to estimate by the measuring-tape, for enlargement of one side is always accompanied by enlargement of the other. The sound side is enlarged partly on account of the increased size of that lung, which has extra labour thrown upon it, partly as an immediate consequence of the pressing forwards of the sternum by the effused fluid, this displacement necessarily carrying with it the ribs covering the unaffected pleura. A far more valuable symptom is the alteration in shape of the affected half of the chest : the chest-wall becomes flattened laterally, and rather prominent on the anterior face, so as to assume something of a square shape ; and this alteration, which can be accurately outlined by the cyrtometer, is a very characteristic feature of the disease.

* Displacement of the heart is sometimes prevented by adhesion between the pleura and pericardium. Therefore the absence of cardiac displacement is no proof that the physical signs are due to another cause than pleuritic effusion. Indeed in some cases of pleurisy the heart's apex is found beating to the left of its natural position. Thus in a child aged four there was complete dulness of the whole left side, with weak blowing breathing and absence of vocal fremitus, but the heart's apex was felt to beat one inch to the left of the nipple line.

Vocal vibration in very young children can, unfortunately, be seldom detected, and therefore its absence—a sign which is so valuable in the diagnosis of pleurisy in the adult—furnishes us with no information. Sometimes, in children of four or five years and upwards, we can detect a slight fremitus on the sound side as the child speaks, but very often no vibration at all is to be detected at any part of the chest; so that in young children we must be prepared to make our diagnosis without its assistance. On percussion we find dulness at the base of the chest on one side. The note is of a peculiar quality: it is a high-pitched, flat sound, which, if much fluid be present, conveys the idea of perfect airlessness, and is combined with great sense of resistance. In uncomplicated pneumonia the percussion-note is not completely dull, and the sense of resistance is little altered. To the practised ear and finger the peculiar quality of the percussion-note, combined with the increased resistance, suggests at once the presence of fluid.

But, besides the quality of the dulness, we derive great assistance from its distribution. In basic pneumonia the dulness is almost always limited at first to one region of the thorax, usually the posterior aspect. In pleurisy, on the contrary, the dulness passes round to the front also, although anteriorly it is more limited in extent. Dulness at the extreme base in front, combined with dulness behind, is strongly presumptive of the presence of fluid. As the fluid increases in quantity the level of the dulness rises both before and behind. It always remains higher behind than in front, although the line which marks its upper limit is sharper anteriorly than at the back, the percussion-note passing more abruptly from dulness to clearness. If the fluid rises higher than the angle of the scapula a tubular note is obtained under the clavicle in front: this is due to the relaxed state of the pulmonary tissue, which is in contact at that point with the chest-wall. The movability of the dulness, when present, is a sign of the utmost value as to the presence

of fluid. It depends upon the fact that the fluid will always, unless prevented by adhesion, gravitate to the most dependent part, so that the limits of the dulness will be altered according to the position of the patient. This symptom is, however, much more commonly to be discovered after absorption of the fluid has begun than during the stage of effusion.

The auscultatory signs of pleurisy in the child may differ very widely from those to which we are accustomed in the adult. In the first place, friction is often absent, or, if present at an early stage of the disease, it may exhibit very peculiar characters. Thus, in the early stages, it has seldom the grating or grazing character which is so distinctive of extra-pulmonary mechanism, but presents itself instead as a sharp crepitating sound, which simulates very closely the râles produced in the smaller bronchial tubes. It is important to be aware of this peculiarity, for otherwise the presence of crepitation, combined with dulness of the base of one side, might mislead us as to the nature of the disease. Crepitating friction is, however, larger than the true pncumonic crepitus, is more superficial, and has not the puffy character so characteristic of inflammation of the lung. It may be heard at the back, front, or side of the chest at the upper margin of the dull area, or even below the level of the fluid.

The breathing is at first very weak at the affected side, and this may be due to the presence of pain, which forces the child instinctively to spare the affected lung as much as possible. The presence of a feeble breath-sound at one base combined with the other symptoms is, as Dr. West has pointed out, a very valuable sign of pleurisy in the child, while the disease is as yet only beginning. As effusion increases the breathing may remain very weak at the base, but higher up (and especially about the angle of the scapula) it often becomes loud and bronchial, and sometimes assumes a tubular character which is indistinguishable from the tubu-

lar respiration of pneumonia. This quality of the respiration is not confined to cases complicated with pneumonia, for it may be present when the temperature is not materially increased ; nor is it limited to the inter-scapular regions or to the immediate vicinity of the spine, but it may be heard also in front of the chest and at the sides, below the level of the effusion. In the axillary and infra-axillary regions, indeed, tubular or even cavernous breathing is very common, and this character may be maintained even in cases of very copious effusion. Sometimes the tubular character of the breath-sound is heard even down to the extreme base. The vocal resonance is usually exaggerated and becomes bronchophonic or ægophonic ; but it is not necessarily so in all cases where the breathing is bronchial : over a spot where the respiration is markedly tubular the vocal resonance may be almost completely suppressed.

Occasionally the quality of the breathing is found, like the dulness, to vary according to position. Thus, in a rickety child of eighteen months old who suffered from pleurisy of the right side, and in whom the dulness reached upwards to the level of a line drawn round the body at about two inches above the nipple, the respiration was intensely tubular both in front and at the back to within a very short distance of the base. On placing the stethoscope over the nipple, marked tubular breathing was heard while the child sat upright ; but on placing him upon his back, the tubular character was lost, and the respiration became of ordinary bronchial quality.

The breathing is not bronchial in all cases. Sometimes we find the signs to agree very closely with those in the adult, the breath-sound being weak or suppressed except close to the spine. Generally, however, at some period of the effusion bronchial breathing can be detected in the child, for the signs seldom remain stationary ; and bronchial respiration heard at one visit may at the next be replaced by a breathing so weak as to be only caught with difficulty ;

while a weak and merely harsh breath-sound may suddenly assume a loud blowing type, without any apparent diminution in the amount of the effusion or the occurrence of any other symptom to account for the alteration.

A good example of an ordinary attack of pleurisy in the child is seen in the following case :—

CASE X.—ORDINARY ATTACK OF PLEURISY.

Maurice R., aged 9 years, was admitted into the East London Children's Hospital on December 3. Until a week before admission the boy had been quite well, but on November 26, a cold wet day, he ran about the streets with bare feet. A few hours afterwards he was seized with pain under the right nipple. He did not shiver or seem cold; on the contrary, his skin was very hot to the touch. He was not sick, but complained much of thirst, getting up several times in the night to drink water. A few days afterwards he began to cough.

State on Admission.—Does not look very ill. Eyes bright; lips pink; nostrils motionless. Generally lies on right side. Says he has now no pain. Seldom coughs. Right side of chest moves less than left during inspiration. A feeble vocal fremitus is felt on both sides, but very faintly towards right base. The heart's apex beats rather to the outer side of the left nipple-line, and is therefore displaced slightly to the left. Percussion is perfectly dull all over right half of chest, at the back, front, and side, with increased resistance. Immediately under the clavicle the note is tubular, and it becomes rather tympanitic about the nipple. Respiration in front is weaker at the right side than at the left, and is very high-pitched and bronchial from apex to base. Just below the right nipple a coarse but faint crepitating sound is heard with inspiration: it is superficial, is not puffy in character, is not limited to the close of the inspiratory act, and is not at all like pneu-

monic crepitation. In the right axilla the breathing is bronchial and moderately loud, but in the infra-axillary region it becomes very loud and intensely tubular. At this spot the vocal resonance is bronchophonic. In the supra-spinous fossa behind the respiration is bronchial, and remains so as far as the angle of the scapula; below that point to the base it is very loud and intensely tubular, and at the base the same superficial crepitating sound is heard as was noted in front. Vocal resonance is increased all over right back, and at base is markedly bronchophonic. No alteration either in the percussion or auscultatory signs to be discovered during changes in position of patient. All over left side resonance is normal, and the breathing loud and puerile. Temperature in evening, 104° ; pulse, 157; respiration, 30.

On December 6 the fluid had rather increased in quantity, for the heart's apex was felt beating half an inch to the outside of the left nipple-line. Vocal fremitus could only be felt at the upper third in front and at the supra-spinous fossa behind. Percussion and auscultation gave the same results as before. Temperature in morning, 99.6° ; pulse, 96; respiration, 24. In evening, temperature, 101.6° ; pulse, 84; respiration, 22. Temperature, yesterday evening, 100.8° ; pulse, 96; respiration, 24.

10th.—Absorption of the fluid has begun. The heart's apex is now just below the left nipple, very slightly to outer side. Intercostal spaces sink in on inspiration at the lower part of the chest. Vocal fremitus is again felt all over right side. On percussion the apex of the lung is now resonant, the level of the dulness being in front at the second rib, behind at the spine of the scapula. Respiration is obscured everywhere in front by a very loud grating friction-sound, and vocal resonance is only slightly increased at the base. At the supra-spinous fossa the breathing is harsh, with crepitating friction; from the spine of scapula to its lower angle very loud and bronchial; at base, bron-

chial, without any tubular quality, and accompanied by loud grating friction. No vocal resonance at back. Percussion still dull in axillary and infra-axillary regions, especially the latter; but on placing the boy upon his left side the dulness almost completely disappears, returning when the sitting posture is resumed. With inspiration a faint crepitating friction is heard in these regions, which becomes very much intensified by a change in position. The temperature has fallen since last visit, being seldom over 100° in the evening. Pulse 90 to 98; respiration 22 to 30.

13th.—The fluid has greatly diminished since last report. Percussion is now natural at right front and in axillary and infra-axillary regions, except just at the base. At the right back the note is unnaturally high-pitched from the spine of the scapula downwards, but is only completely dull at the base. Loud grating friction-sound is heard all over the right back and side, and also, but less loudly, about the right nipple, in which latter spot only can the breathing be heard. It has a slight bronchial quality. Heart's apex beats half an inch below the left nipple and half an inch to inner side. Temperature in morning, 99° ; pulse, 90; respiration, 18. In evening, temperature, 100.4° ; pulse, 104; respiration, 21.

The after progress of the case was very slow. The fluid became gradually absorbed, the dulness disappearing first in the front of the chest, but remaining for some time at the base behind. Even as late as February 4 absorption was not quite completed. On that date the following note was taken:—Boy fat, and looks well. No falling-in of the affected side of the chest—on the contrary, there is a slight bulging of the base, as estimated by the cyrtometer. The right side moves well with inspiration, except towards the base, where the movement seems rather hampered. In the upright position there is complete dulness at the right posterior base, with considerable resistance. The upper margin of this area is well defined, the dulness passing abruptly into

good resonance above. When the boy stoops forwards, so that his back becomes horizontal, the dulness is replaced by clearness. Air seems to enter the lung perfectly to the extreme base. Respiration there is loud and slightly harsh, and a feeble catch of friction is heard at the end of a deep inspiration. Vocal resonance is slightly increased, and vocal fremitus, felt everywhere else about the chest, is here hardly perceptible.

A few days afterwards the boy was discharged from the Hospital, but he returned on March 7 for examination. By this time all the physical signs had disappeared, with the exception of a feeble crepitating friction, which was caught towards the end of inspiration in the infra-axillary region.

The above example affords a very good illustration of the peculiar characters of the physical signs often met with in the pleurisy of childhood. In such a case the intensely tubular quality of the breathing, and the crepitating character of the friction-sound, might lead to error in diagnosis unless close attention were paid to these signs and to the general symptoms by which they are accompanied.

The history of the attack, which was stated to have begun suddenly with pain in the side preceding the cough, pointed distinctly to pleurisy, and the general symptoms noted on the boy's admission were sufficiently characteristic. He did not look very ill; the strength was not much reduced; the cough was rare; the nares motionless; and although the temperature was high, being 104° , yet the breathing was not hurried, and the pulse-respiration ratio consequently undisturbed.

On examination of the chest the dulness was found to be extreme, a dead, flat note being elicited by the finger, which is characteristic of the presence of fluid, while the sense of resistance was very much increased. The dulness was very extensive, and reached to the front of the chest, the whole of the right side being dull from apex to base. The tubular breathing was limited to the base at the side and back. If

the case had been one of pneumonia, tubular breathing would have been heard over the whole of the dull area. Besides these signs, which were very distinctive, the heart was displaced to the left—a change of position which is not seen in cases of mere consolidation of the lung; and vocal fremitus, which was well marked on the sound side, was felt but faintly at the back over the seat of effusion. The superficial crepitating friction-sound, too, which occupied the whole extent of the inspiration, was very different from the puff of fine crepitation heard at the end of a deep inspiration in pneumonia. Besides, this was noted only at the base of the chest, where the breathing was tubular. In pneumonia crepitation ceases in the part of the lung which has become sufficiently consolidated to allow of the production of tubular breathing.

As the case advanced, and the fluid began to be absorbed, the loud grating friction and the alteration of the percussion-note on change of position furnished additional evidence, had further proof been wanting, of the nature of the case.

Sometimes the fluid, instead of being free to occupy the whole of the pleural cavity, is confined to certain regions by bands of lymph which cause the pleural surfaces to adhere at particular spots so as to partition off the parts in which effusion takes place. In such cases the disease is less easy of diagnosis, and is still more likely to be mistaken for inflammation of the lung.

CASE XI.—LOCALISED PLEURISY.

Emma W., aged 5, a delicate-looking child, with small features, fair hair, and a thin transparent skin, had had a cough more or less for two years—ever since a severe attack of pertussis. There was a little dulness at the right supraspinous fossa, and the breathing there was high-pitched and harsh, with prolonged expiration. She was still under

treatment when (on May 30) she had an attack of measles. The disease run its usual course, but the cough was very severe. On June 7 she began to complain of pain in the right side : she still coughed a great deal, seemed languid, had no appetite, and was very hot at night, perspiring freely. She was seen on June 14 looking pale and weakly, but could walk well (she had, indeed, come on foot to the Hospital), and was not at all prostrated by her illness.

On examination of the chest the front of the right side was found to be completely dull from apex to base, with greatly increased resistance. The dulness did not extend into the axillary region, but ceased laterally rather abruptly at a line drawn downwards from the anterior fold of the armpit. The breathing above and just below the clavicle was bronchial ; over the lower two-thirds it was very loud and tubular, without any rhonchus. Over the area of dulness the resonance of the voice and cough was unnaturally strong. At the right back there was no dulness, with the exception of that at the supra-spinous fossa already referred to, but the respiration from apex to base was bronchial, with a fine subcrepitant rhonchus towards the end of inspiration. Temperature in axilla, 99.2° ; pulse, 136, unequal ; respiration, 50.

On June 21 the subcrepitant rhonchus had disappeared from the back, otherwise the signs remained the same as on the previous visit. The cough was very troublesome, the appetite poor, and she perspired very much at night. Temperature 99.6° ; pulse, 148 ; respiration, 42.

The child improved but slowly, and it was not until July 31 that any notable change was observed in the physical signs. By this date the child was fatter and looked pretty well, although pale, and she slept well at night, but was sometimes disturbed by her cough. The dulness was now less complete than before at the right front, and below the clavicle was rather tubular. The breathing was cavernous from apex to base, without rhonchus, and the vocal

resonance bronchophonic. At the back the supra-spinous fossa alone was dull, and the breathing there cavernous, as in front; lower down the respiration was bronchial, and at the base merely harsh. No rhonchus was heard at the back, and the vocal resonance, except at the apex, was natural.

After this the patient was not seen again for six months. By this time she had become fat, and looked very well. There was a little flattening of the right mammary and infra-mammary regions; the right nipple was a little lower than left; and the spine was slightly curved, with concavity to right. The chest measured at the level of nipple on the right side ten inches and one-eighth, on the left ten inches and a half. There was no dulness anywhere about the right side; the supra-spinous fossa even was quite resonant and the breathing everywhere healthy. About the right nipple a faint superficial crepitating sound was heard with the latter half of inspiration. This was no doubt pleuritic, and was the only indication left of the foregoing disease. The child was said to cough sometimes in the morning on first rising, but not afterwards in the day.

The treatment throughout consisted of quinine and cod-liver oil.

In this case the limitation of the principal physical signs to the front of the chest furnished a possible source of error, which it required a careful consideration of the general symptoms and of the whole aspect of the case to avoid. Dulness limited to the front on one side, combined with tubular breathing and increased resonance of the voice, suggested the presence of pneumonia, and this view was supported by the hurry of respiration, which on the first visit reached the number of fifty in the minute. The dead, flat character of the dulness was, however, very different from the incomplete dulness of pneumonia; the resistance to the finger was also greater, and the area of dulness was rather abruptly margined towards the axilla.

The entire absence of rhonchus of any kind from the front of the chest was a phenomenon rarely to be met with in a case of inflammation of the lung, and itself suggested pleurisy. Such a condition is, however, occasionally found where the pneumonic consolidation is very complete, but it rapidly passes away, seldom persisting longer than two or three days. In the present case rhonchus was absent throughout.

Besides the physical signs, the general symptoms were very decidedly opposed to the pneumonic view of the case. The child, although pale and delicate-looking, had not the aspect of one seriously ill, and she had been able to walk without assistance to the hospital. This point is of very great importance, and tells very strongly not only against pneumonia, but also directly in favour of pleurisy. That a child with caseous deposit at the apex of one lung, who had but just passed through an attack of measles, and who presented *recent* physical signs such as those described, should be able to walk from her home to the hospital without apparent suffering, could only be possible in a case of pleurisy. Had the disease been pneumonia, consolidation of such extent, especially when succeeding to measles, would have been attended by a degree of prostration sufficient to put such an effort out of the question. Besides this, the temperature of the body was little higher than natural (99.2°), and at night she perspired freely.

There can be little doubt, therefore, that the case was one of circumscribed pleurisy. The subcrepitant rhonchus heard at the back was due to accompanying bronchitis, so common during and after an attack of measles; and the bronchial breathing heard posteriorly was no doubt owing to compression of the lung by the fluid. The hurry of respiration must be attributed to the exertion she had just undergone.

The diagnosis was confirmed by the after progress of the case. The long persistence of the physical signs without notable change, and especially without the appearance of

rhonchi, was very significant ; and the moderate retraction of the side found six months after the attack afforded a final proof that a correct estimate had been formed of the nature of the disease.

The removal of the small consolidation of the right apex is a point to be noted in passing. This was probably of some standing, and may have dated from any time subsequent to the attack of pertussis two years before, during all which time the child had remained pale and thin, with a frequent cough. The deposit, however, was undergoing no active change, and while thus indolent there is always in young subjects the possibility that absorption may take place. Its removal appears to have been complete, for at the last examination no signs were left to indicate that the apex of the lung had ever been the seat of disease. Limitation of the liquid effusion is in most cases owing to the presence of old adhesions, the previous occurrence of a pleuritic attack having left the adjacent surfaces of the pleuræ adherent to one another in certain spots. In this case, however, there was no history of any previous attack, although such might have occurred and been overlooked.

The mode of termination of pleurisy is the same in the child as it is in the adult. Absorption of the fluid may take place with or without retraction, or the fluid may be purulent, and the case become one of *empyema*. In children, cases of empyema are very common, for the fluid becomes purulent at an earlier period with them than it does with the adult.

In the following case the fluid, although purulent, was being gradually absorbed, and some retraction of the side had taken place :—

CASE XII.—EMPYEMA—ABSORPTION OF FLUID.

John C., aged $3\frac{1}{2}$ years, was admitted into the East

London Children's Hospital on August 29. Has always been weakly and very subject to coughs. Five weeks ago the child was seized with a fit of convulsions. On recovery he complained of pain in the left side, and was very feverish and thirsty. He soon began to cough, and his breathing was very shallow and rapid. He was seen by a medical man, who treated him for "inflammation of the membranes of the brain." Since that time he has been very weak and low, and has been troubled with a frequent cough.

August 31.—Child pale and thin. Ends of long bones enlarged; ends of ribs nodular. Fingers distinctly clubbed; legs very small and thin. Can stand, but is hardly able to walk without assistance. Skin not particularly harsh. Spleen much enlarged; liver normal in size. Left shoulder and scapula lower than right; and spine slightly curved, with concavity to left side. Heart's impulse felt at the left side of ensiform cartilage. Left side of chest moves very little with inspiration. There is dulness over the whole of the left half of the chest, both back and front, with greatly increased resistance; in the first and second interspaces the note is rather tubular. The respiration is loud and coarse in left front as far as the nipple, becoming at that point rather tubular in quality. In the left axilla the respiration is blowing, and becomes tubular in the infra-axillary region. At the back the breathing is tubular at the base; higher up it is diffused blowing, and remains so as far as the supra-spinous fossa. No friction or rhonchus can be heard, and the vocal resonance is only slightly increased, not at all ægophonic. Temperature yesterday evening, $104\cdot4^{\circ}$; this morning at 9 a.m., $99\cdot4^{\circ}$. The chest was punctured posteriorly on the left side, about two inches below, and in a line with, the angle of the scapula; and a small quantity of thick purulent fluid was withdrawn with the pneumatic aspirator. The temperature the same evening was $101\cdot6^{\circ}$.

On September 3, five drops of tinct. ferri perchlor. were

ordered three times a day, and one ounce of wine was added to the diet, which consisted of milk and bread, with meat and broth on alternate days.

September 10.—Sweats very much about the head and neck; skin of those parts covered with miliaria. The child is fretful when touched, but does not seem in pain; eats heartily; coughs short and loose.

Percussion of the left half of the chest gives pretty much the same results as before. At the extreme base, both before, behind, and laterally, the note is tympanitic from transmission of the stomach note; elsewhere the percussion-sound is dull, except just below the clavicle and in the axilla, where it is tubular. The respiration at the back is now only bronchial, without any tubular or blowing quality. It is also bronchial in front, and in the axilla is cavernous. The vocal resonance is everywhere on the left side bronchophonic, and as heard at the angle of the scapula the quality of the voice is rather ægophonic. Temperature, 6 p.m., 102.2° ; pulse, 154; respiration, 58. Since August 31 the temperature has seldom been over 99° in the morning; in the evening it has varied from 101° to 102° .

The physical signs remained very stationary from this date until the end of the month, although the heart was gradually drawn back to its natural position.

On October 1, the chest measurement at the level of the nipple was, on the right side $9\frac{1}{2}$ inches, on the left $9\frac{7}{8}$ inches. Half a teaspoonful of ol. morrhuæ was ordered to be taken three times a day in addition to the iron mixture.

October 9.—Percussion is now natural in front of left side, and the left supra-spinous fossa is resonant. Over the scapula region the percussion-note is tubular, and at the posterior base it is dull. Respiration is bronchial at the base behind; superiorly it is harsh, with prolonged expiration. All over the upper half of left back is a peculiar clicking sound like very small bubbles, not at all superficial. In front the breathing is merely harsh; in the

axillary and infra-axillary regions it is slightly bronchial and is accompanied by a faint crepitating friction.

From this date the physical signs became gradually more and more healthy, and on November 5 the least possible dulness remained at the left base behind, but the breathing there and everywhere else was perfectly natural, without friction or any other abnormal sound. The child became quite fat and strong on his legs. The spleen, however, did not diminish in size, but remained very large, reaching as low as the level of the umbilicus.

The history of the attack in this case was very characteristic, and the distortion of the left side of the chest (lowering of the shoulder and scapula, with slight lateral curvature of the spine), inconsiderable though it was, combined with the physical signs, rendered the diagnosis of pleurisy an easy matter. Absorption of the effused fluid was evidently far advanced, but the elevation of temperature seemed to imply that the presence of purulent matter in the pleura was injuriously affecting the system; and it was therefore thought advisable to remove what little there might be left without delay. The after progress of the case, although slow, was satisfactory, and the speedy improvement which followed the administration of cod-liver oil is to be noted. The oil might indeed have been given with advantage at an earlier period. The child was the subject of rickets, and in such cases the influence of this drug in furthering nutrition and improving the general condition is very striking. The enlargement of the spleen, one of the consequences of the rickety state, did not appear to impede recovery, although the size of the organ remained unaltered. Such cases, indeed, generally do well.

The presence of pus in the pleural cavity is usually accompanied by signs that the nutrition of the patient is greatly interfered with. Still, all children do not suffer alike. In some there is rapid loss of flesh, and the patient becomes thinner and paler every day. He shows little desire for food,

has a distressed look in his face, is restless and irritable and sleeps badly at night. His circulation is feeble, his fingers become clubbed at the extremities, and his hands and feet are cold. In other cases the general symptoms are much less striking. The child, no doubt, grows very pale, and shows a distinct loss of flesh. His fingers are clubbed; his complexion is sallow, and his hands and feet tend to be cold; but his strength remains fairly good; he sleeps well at night and shows sufficient appreciation of his food. This difference in the degree to which the system is affected by the purulent contents of the thorax seems, in most cases, to depend upon the presence or absence of fever. Empyema does not always set up pyrexia; sometimes the temperature remains completely normal throughout the illness. It is especially in children of strumous constitution, and in others who, although not scrofulous, have a highly irritable nervous system that fever is so likely to be induced. In such subjects the temperature is almost always over 100° in the evening; rising sometimes to 102° or 103° , but sinking again to a lower level in the morning. The highest temperature is generally found at about 2 or 3 P.M., after which hours the fever abates to some extent.

If no operation is performed for the evacuation of the purulent fluid, this may become gradually absorbed, and the child may eventually recover. In other cases the fluid points at some part of the chest; a swelling appears at one of the interspaces, becomes red, and after a few days bursts, and the fluid slowly drains away. When this takes place the child may recover, the opening closing after several weeks; or the perforation may remain open, and the child die, worn out by long-continued suppuration. Instead of opening externally, a perforation may take place into a bronchial tube, and the fluid be evacuated in the form of profuse expectoration. These cases often end favourably. In a third class of cases the presence of purulent

fluid in the pleura seems to be a determining cause of a general formation of tubercle, and the patient dies from acute tuberculosis.

CASE XIII.—EMPYEMA—SPONTANEOUS EVACUATION
OF THE FLUID.

Anne B., aged 6, a thin child, rather small for her age, very pale and weakly-looking, had been suffering, it was said, from "low fever" for a considerable time. Her illness had begun in the middle of February with fever and cough, and she had ever since been getting paler and thinner, although the fever had in a great measure subsided. Her cough was still troublesome, but was loose, and gave her no pain.

When seen on June 4 she complained of pain in the left hypochondrium, and this part was said to swell at times. On examination, the left side of the chest was seen to be almost motionless in respiration. There was no fulness at the seat of pain, but that spot seemed tender on pressure. The whole left back was completely dull from apex to base, with increased resistance. The breathing was loud and almost cavernous over the whole dull area, and the vocal vibration was annulled except at the supra-spinous fossa, while the resonance of the voice was everywhere increased. In front of the same side percussion was dull and the breathing blowing, although weaker than at the back. The heart was displaced slightly to the right.

A few days afterwards a swelling the size of a hen's egg formed in the left nipple line, the centre of which was over the eighth interspace. This was very tender to the touch, and fluctuated freely. As the mother objected to any operative procedure, she was directed to apply a succession of hot linseed-meal poultices. In the course of the week a second swelling appeared higher up on the same side, just below the nipple. This also fluctuated. The superficial

veins of that side, both of the chest and belly, were very well marked. The skin over the abscesses became very red and tense, and on July 3 the lower one burst, and discharged a considerable quantity of thin purulent fluid. After this the upper swelling subsided without opening. The child seemed at once very much relieved; her appetite, which had been quite lost, returned; her spirits improved, and she began at once to regain flesh.

On July 26 the following note was taken:—Looking much fatter and stronger; runs about, and her spirits are good; eats heartily. Cough loose, and very troublesome at night. Perspires very much, especially when in bed. The abscess is still discharging, and about an ounce of thin watery fluid comes away from it in the day. There is much flattening under the left clavicle, and the left front of chest does not move in inspiration; laterally, however, the intercostal spaces sink in a little in inspiration, and the ribs move slightly. Spine curved laterally, with convexity to right. Shoulder and lower angle of scapula on the left side rather depressed. Percussion is dull all over left half of chest—both front and back. In front no respiration is heard at all, but over the whole left back it is loud and blowing. Vocal resonance there is slightly stronger than natural.

After this report the child was not seen again until the following February, six months afterwards. The improvement was then very remarkable. The abscess had continued to discharge for six weeks. It had then closed, and the child had been ever since constantly improving. She now coughed very little, and only in the morning or after exertion, and then sometimes expectorated a little thick yellow phlegm. She was fat, and looked strong and healthy.

On examination of the chest some slight retraction was seen at the lower part of the left side, but the deformity was scarcely noticeable; the left shoulder was not depressed, the spine was straight, and the nipples were on the same

level. The apex of the heart beat one inch below the nipple in the nipple line. No dulness was found at any part of the left side, and the respiration was generally healthy, although towards the upper part of the left back it was rather loud and slightly blowing. Vocal vibration was felt faintly as far as the extreme base.

This case is interesting from the fact that in empyema spontaneous evacuation of the fluid through the chest-wall is rarely followed by recovery. Indeed, if the case is left to nature alone, the child most commonly dies. The unfavourable termination under such conditions is due to the incompleteness with which the pleural contents are evacuated. The abscess usually points in one of the upper intercostal spaces—often in the fifth; and although a more or less constant discharge escapes from the opening, there is still retention of fluid in considerable quantity at the base of the chest. Therefore, unless measures are taken to promote free emptying of the pleural cavity, a cure is almost impossible. Fortunately in the above case the abscess pointed low down in the chest so that the purulent fluid was enabled to flow readily away. By this means the pleural cavity was completely emptied in the course of six or eight weeks, and the patient then rapidly regained her health.

The case also illustrates the extraordinary readjusting power possessed by childhood. The left lung had been exposed for a considerable time to great pressure from the effused fluid, but it had notwithstanding been able, on the removal of this fluid, again to expand, and to do this so effectually as almost entirely to remedy the deformity at first noticed.

Sometimes, instead of discharging itself through an intercostal space the purulent collection passes into a bronchus, and is coughed up and expectorated. Contrary to what might be supposed, evacuation of the pleural cavity by this route is, on the whole, a favourable symptom, and many such cases recover. It is curious that pneumonia is rarely induced

by the communication with the air-tube ; indeed, if death ensue, it is not easy—often it is impossible—on examination of the body to discover the opening through which the entrance of the fluid into the air-tube took place. When the fluid is thus brought up through the lung, its evacuation is effected by fits of coughing. But the ejection of pus is not continuous. Often a considerable interval is seen between the successive appearances of the purulent matter. Usually a discharge of pus occurs every few days. This may continue until the pleural cavity is completely emptied of its irritating contents, or after a certain quantity has been thus evacuated the fluid ceases to escape, by this channel, and the remainder points through the chest-wall, or is withdrawn by operation.

When fluid has existed for a considerable time and has exposed the lung to great pressure, its removal, however effected, is often followed by retraction of the diseased side of the chest. Serious deformity from this cause is, however, less frequently seen in the child than it is in the adult, probably on account of the tendency to rapid repair which has just been referred to. Still, instances of serious chest distortion are occasionally met with.

CASE XIV.—GREAT DISTORTION OF CHEST FROM PLEURISY.

Thos. MacE., aged fourteen, was brought to the East London Children's Hospital with much deformity of his chest, the result of an attack of pleurisy fifteen months before. The right side was much retracted. The shoulder and scapula were markedly depressed ; and the spine was curved laterally, with the concavity turned towards the affected side. The scapula, besides being depressed, was tilted backwards at the lower part, so that its inferior angle projected. In front the infra clavicular region was deeply hollowed ; the nipple was half an inch below the level of the

left ; and, at the lower part of the affected side, a special depression of the seventh rib, retracted below the level of the others, left an oblique diagonal groove passing transversely downwards towards the middle line. There was very little movement of this side during inspiration. The veins of the chest were unusually full, and the heart was drawn to the right so that its apex could be felt beating at the ensiform cartilage. The circumference of the thorax at the lower part of the sternum was, on the left side $13\frac{1}{8}$ in., on the right only $11\frac{1}{4}$ in. The percussion note on the right side was generally high pitched, without being anywhere completely dull. The breath sound, above, was loud and harsh ; below, it was much weaker, and was faintly bronchial. The vocal vibration was exaggerated, except at the base in front. The boy was in fair health, but was subject to cramp and to breathlessness after exertion.

This case is not given as an illustration of extreme retraction, but furnishes a fair example of the more serious form of chest distortion seen in children as a consequence of pleurisy. In such a case the tilting backwards of the angle of the scapula, the marked depression of the shoulder, and the absence of signs of excavation would distinguish the disease from fibroid induration of the lung—a complaint which also produces great retraction of the side. (See Chapter VII.)

When the purulent fluid is not evacuated, either spontaneously or by operation, its retention in the chest-cavity is always attended with considerable risk ; for not only is the general nutrition of the child seriously affected while such fluid remains, but there is also a direct danger that its presence may lead to the development of general tuberculosis, as in the following case :—

CASE XV.—EMPYEMA—SECONDARY ACUTE TUBERCULOSIS.

Charles B., a little boy, aged two years and nine months,

sixteen teeth, was admitted into the East London Children's Hospital on September 17.

Little information could be gained as to the history of his attack. The mother could only say that he had been healthy two or three months previously ; that he had then begun suddenly to cough and lose flesh, and that during the last month his bowels had been relaxed five or six times a day, causing him to waste very rapidly.

State on Admission.—Child greatly emaciated ; eyes hollow ; cheek-bones project ; complexion rather earthy ; head square-shaped ; fontanelle size of a shilling, depressed ; veins of forehead full. Abdomen rather swollen, and feels hot to hand ; abdominal veins full ; no enlarged glands felt in belly on deep pressure ; spleen and liver appear to be of normal size. Ends of long bones not large ; fingers slightly clubbed. Tongue clean and red. Stools yellow, slimy, and offensive. Has a frequent cough. Temperature in rectum 100.6° ; pulse 130 ; respiration 40.

Physical Signs.—Very little respiratory movement comparatively on right side. Intercostal spaces not bulged, and can be seen to sink in to a slight extent on that side at the lower front ; but on the left side they sink in deeply at each inspiration. Heart's apex beats on the left side in the fifth interspace in nipple-line. The chest measurement at the level of the nipple is on the right side $8\frac{7}{8}$ inches, and on the left $8\frac{5}{8}$ inches. No vocal fremitus felt either side. Percussion dull all over right side, back and front. Behind, the dulness is complete, with extreme sense of resistance. In front, the dulness is more tubular, especially under the clavicle, and does not reach quite to the middle line. Still, the percussion-note at the right edge of the sternum is not quite resonant. Respiration all over right lung is loud and bronchial, with intense resonance of voice and cry. Between the scapulæ the breathing is cavernous. Here and there dry rhonchi are heard, but there is no friction or crepitation. The left lung is apparently healthy.

A lead and opium mixture was ordered to check the diarrhœa, and ten drops of brandy every three hours. For his diet, bread and milk. The diarrhœa ceased at once, and the child's appetite begun to improve. He continued still very thirsty, and the temperature remained high, varying from 100° to 102° —generally one degree of difference between the morning and evening temperature.

September 24.—Child seems weaker. The bowels act twice a day; motions not offensive. Physical signs as before. The right side seems tender, and the pressure of the stethoscope causes pain. Cough loose and rather frequent; if the child coughs violently all the veins swell up, even down to the wrists; jugular veins then much distended. Temperature in evening 102.4° ; pulse 120; respiration 30. He was ordered *ol. morrhue* ʒss. *ter die*, with a mixture containing *tinct. ferri perchlor.* Some meat and broth on alternate days were added to his diet.

On October 4 the respiration on the right side in front was covered by a grating rhonchus, and at nipple a coarse crepitation was heard. These, however, had disappeared by October 9. On this date the two sides of the chest were measured with cyrtometer, when at the nipple level the right side was found to measure $8\frac{7}{8}$ inches, the left $9\frac{1}{8}$ inches. The temperature remained high at night, varying from 101° to 103° , or even 104° , and sinking to 100° in the morning; pulse 130 to 140; respiration 44 to 48.

22nd.—Child wasting more and more. Physical signs much the same on right side. Over lower part of right front fine bubbling is heard at the end of a deep inspiration. On the left side percussion is rather dull in the supra-spinous fossa, and generally over the left back is too high-pitched. Respiration in supra-spinous fossa is bronchial, below, harsh, with a little dry rhonchus. Over left front respiration is harsh, with a little sonorous rhonchus. Temperature in evening 104° ; pulse 120; respiration 48.

On October 25 the child had an attack of diarrhœa. The temperature then fell to 100° in the evening, and remained low till October 29, when the thermometer again marked 104° in the rectum.

On November 5 the percussion signs were unaltered, but under the right clavicle the loud bronchial respiration was accompanied by a large metallic bubbling rhonchus which was also heard downwards as far as the extreme base. At the nipple the breathing was blowing, almost tubular. The same bubbling rhonchus was heard also in the right infra-axillary region and at the back, but posteriorly the bubbles were smaller and less metallic. After a violent cough the rhonchus partially disappeared at the front and side. On the left side some fine crepitantrâles were heard at the end of inspiration. Temperature in the morning 102° ; pulse, 130; respiration 44. In the evening, temperature 103° ; pulse 132; respiration 56. On the previous evening (November 4) the temperature had been as high as 105° .

From November 5 to November 19 the temperature continued exceedingly high in the evening, the thermometer marking 104° to 105° , and on one occasion (November 15) 106° , in the rectum. There was always a great fall towards the morning, and the heat of the body was almost uniformly 99° at ten a.m. There was usually a difference of about twenty beats between the morning and evening pulse, and the respiration was also always higher at night, ranging at that time from 42 to 60. The subsidence of the fever in the morning was always accompanied by a marked change in the behaviour of the child. In the morning he would sit up in bed and play with his toys, eat a hearty breakfast, and sometimes cry out for cake. In the evening he lay with flushed face and parched lips, completely indifferent to all that passed around.

On November 18 the bubbling rhonchus was universal on the right side, and at the left back the breathing was harsh

and was accompanied by a rhonchus coarser than before, and mixed here and there with a finer crepitation.

On November 20 patches of dulness were discovered at the left back, and there was greater resistance to the finger; the breathing on that side was bronchial, and everywhere it was accompanied by a coarse bubbling rhonchus as on the right side. On this date the temperature suddenly became lowered; in the evening it reached no higher than 99.2° . On the following morning it was 97° , and fell in the evening to 95.2° . A few hours afterwards the child died. The temperatures were taken throughout the case in the rectum, and with great care.

Autopsy.—Body extremely emaciated. Right lung almost entirely adherent to chest-wall; the adhesions with difficulty broken down, especially at apex, where they are so firm that the lung is torn in bringing it away. A little purulent fluid—about one ounce—is found in right pleura at the base. Right lung smaller than left, tough and firm. Scattered through the substance, and forming nodules projecting on the surface under the pleura, are yellowish-white non-friable tubercular masses, varying in size from a pin's head to a pea. These have not the granular appearance of cheesy masses, and do not appear to be softening. With them are more pellucid small bodies, evidently grey granulations. In many places, where the tubercles are collected closely together, the intermediate lung tissue is the seat of quite recent broncho-pneumonia. Left lung not condensed, but presents the same masses. It was firmly adherent to the pericardium, which could not be detached from it. The same masses are found in the spleen, which is thickly studded with them. They are also present, although to a less amount, in the kidneys, and on the upper surface of the liver are some grey granulations. There are no granulations in the meninges.

The diagnosis in this case was not difficult. The complete dulness over the whole of the right half of the chest,

both at the back and at the front, combined with the extreme sense of resistance; the breathing, which was everywhere loud and piercingly bronchial; the intense bronchophonic resonance of the voice and cry; the absence of liquid rhonchus or crepitation,—these signs occurring, without any contraction of the side, at so late a period of the disease (for the child had been ill two or three months at the time of his admission) could only be the result of pleurisy. Inflammation seldom affects the whole of a lung from apex to base, and pneumonic consolidation, if extensive, does not long remain indolent. Small pneumonic deposits may remain unchanged in children for many months, and may eventually be removed; but large deposits soon begin to give evidence either of commencing absorption or of degeneration and decay, leading to destructive disease of the lung in which they are situated. Again, the sense of resistance to the finger on percussion is very much greater when fluid is present, or in the case of a lung solidified by pressure, than where the consolidation is owing to pneumonia; and this alone is a very important element in the diagnosis. The heart, again, was slightly displaced to the left, and the apex was found to beat in the fifth interspace in the nipple-line. It had been probably pushed further to the left, but was being drawn back into its normal position as the effused fluid became gradually absorbed. At the time of admission there was probably little fluid left in the diseased pleura, for the intercostal spaces on the affected side were not prominent, and were seen to sink in slightly at each inspiration.

In the after progress of the case the continued wasting, the high evening temperature, combined with the occurrence of dry and crepitating rhonchi on the diseased side, and eventually in the opposite lung, pointed to the occurrence of secondary tuberculosis.

The pleurisy was here evidently the primary disease, occurring, as it did, suddenly, while the child was in his

ordinary health ; and the tubercle was a secondary formation. In many cases, however, tuberculosis is the primary disorder, and itself gives rise to a secondary pleurisy. The pleurisy is then generally of the fibrinous variety, a plastic exudation being thrown out, which is seldom attended by liquid effusion, and gives rise to but few symptoms, pain in the side being as a rule the only thing complained of.

The ordinary serous pleurisy is always accompanied by a certain amount of fibrinous exudation, and when this is extensive it may lead to secondary results after the fluid which was first thrown out has become reabsorbed. These cases form a special variety of basic lung disease, to which, as occurring in the adult, attention has been directed by Sir Andrew Clark. The pathological alteration consists in an excessive formation in the base of the lung of fibroid tissue, which causes a loss of elasticity of the part, so that it no longer contracts and expands as in health. If present to an extreme degree, the part attacked shrinks, the bronchial tubes become dilated, and the heart may be drawn towards the affected side. At first, however, there is no contraction of the base of the chest, and this contraction may even be absent throughout if great dilatation of the bronchial tubes be present. (See Chapter VII.)

In every case of empyema as soon as the purulent nature of the fluid has been ascertained, we should at once adopt measures for emptying the cavity of the chest. There is little likelihood that the fluid will be absorbed, and this process, could it occur, would be necessarily a long one. Prompt evacuation is therefore essential to a speedy cure. Still, even when operative interference is had recourse to early, and a fistulous opening in a suitable position is established in the chest, recovery, although in the child the most probable termination to the illness, is yet far from certain. Sometimes even in these favourable conditions the pleural cavity continues to secrete a thin pus, the abdominal

organs become amyloid, and the patient wastes and dies. In other cases general tuberculosis follows as a secondary lesion. In others, again, the inflammation extends to the peritoneal cavity, and the child dies of peritonitis.

Extension of the inflammation from the chest to the belly is not a common accident. Even when it occurs the complication is not readily recognised; often its existence is not ascertained until the body is examined after death. In one case of the kind which came under my notice—a little boy of three years old—the extension of the inflammation was announced by a rise of temperature to $102\cdot6^{\circ}$, by looseness of the bowels and by vomiting. The child wasted very rapidly, soon became collapsed, and died in a few days. But the symptoms are often much less obvious, and the physical signs presented by the belly itself may be so little marked that the most careful examination fails to discover sufficient evidence of the inflammatory lesion.

CASE XVI.—EMPHYEMA, PARACENTESIS, EXTENSION OF
INFLAMMATION FROM THE CHEST TO THE BELLY.

Arthur S., aged twenty-one months, with sixteen teeth, was admitted into the East London Children's Hospital. The boy came of a healthy family, and had always been strong until a few months previously, when he had swallowed some spirits of hartshorn, and had thus set up a chronic inflammation of his larynx.

Five weeks before admission the child was taken ill with fever and lost his appetite. Some days afterwards he was noticed to cough, and for a week before coming into the hospital the cough had been very frequent, dry, hacking, and painful. The child during this time often complained of pain in his left side, and seemed to be relieved by gentle rubbing. He was said to have lost flesh considerably.

On admission, the percussion note was found to be completely dull all over the left side, both at the back and at the

front. Resistance was greatly increased. The respiration was weak and harsh generally, but was bronchial towards the apex behind, and was tubular under the clavicle. There was slight recession of both sides of the chest at the lower part with each inspiration, and also of the epigastrium. The jugular veins on both sides were full. The heart's apex was felt between the fifth and sixth ribs, half an inch to the inner side of the nipple (vertical) line. The child's water contained no albumen. The temperature was normal on the first night, but rose a day or two after admission, and remained for about a week between 100° and 102° in the evening.

The chest was punctured several times with the aspirator, on each occasion a small quantity of greenish healthy pus was withdrawn. Afterwards, as the physical signs continued to show the presence of fluid in the chest cavity, the chest was opened antiseptically and six ounces of thick ropy pus escaped. A double caoutchouc tube was then placed in the wound to allow of complete drainage of the pleura. After the operation the child seemed more comfortable, but irregular fever still persisted. A few days later the belly was noticed to be full, but there was no tenderness or increased tension of the parieties. To the touch the abdomen felt perfectly normal. The child, however, looked very ill; his face was slightly livid, and his nares worked as he breathed.

As the days passed the fulness of the belly continued undiminished, but there was no pain or increase of tension, nor any sensation of abnormal solidity of the contents. Repeated examination, indeed, failed to discover any reason for the distension of the belly, and its prominence was attributed to flatulence. The child's appetite now failed; some diarrhoea came on, and the boy, growing daily weaker and thinner, died seventeen days after the operation had been performed upon his chest.

The physical signs of the left side had improved after

withdrawal of the pus; but there remained considerable dulness and much resistance of the chest-wall to percussion on that side.

During the course of the illness there was irregular fever. Now and then for a few days the temperature would be normal, but it soon rose again, and the mercury often stood between 102° and 103° in the evening. The fever was not greater after the belly had been noticed to swell than it had been previously; but the child seemed much more seriously ill.

On examination of the body, no fluid was found in the left pleural cavity; but the lung of that side was much compressed, and was attached by dense adhesions to the wall. The pleura on that, and also on the opposite side, was much thickened. The abdominal cavity contained eight ounces of thick, turbid fluid; and much thick yellow lymph was found about the right lobe of the liver, the ascending colon and the caecum. The liver itself was slightly fatty. Nowhere about the body were grey granulations or cheesy masses to be observed.

This case well illustrates the difficulty sometimes met with in detecting the presence of acute peritonitis in the child. If it were not an actual fact familiar to most clinical physicians, it would seem incredible that inflammation of so sensitive a membrane as the peritoneum could exist without pain, tenderness, or any alteration of the physical signs of the abdomen appreciable by the touch. Ordinarily in cases of abdominal lesion, when pain and tenderness are absent, there remains increased tension of the parietes when pressure is made upon the belly, as if the muscular wall contracted instinctively to protect the inflamed part from injury; indeed, increased tension of the parietes on one side only of the belly is strongly suggestive of an internal lesion. But in the above case even this symptom was absent, and an unusual fulness of the belly was the sole appreciable deviation from a normal state. Of the cases of

latent peritonitis which have come under my notice, some have been, like the above, cases of secondary inflammation by extension. Others have been cases of tubercular peritonitis. In the tubercular form of the lesion, as will be mentioned hereafter, absence of all local symptoms is not rare during the early stage of the illness.

The importance of accurate diagnosis in a case of pleurisy need not be insisted upon, but common as is the disease, its presence in the child is often overlooked: at any rate in hospital practice it is not rare to meet with cases of pleurisy, the nature of which seems to have been previously misapprehended. "Low fever" or "debility" are names often applied to cases of pleuritic effusion; but here, in all probability, the chest had never been examined at all. In other instances, where an examination of the chest has been made, but no doubt very hurriedly and superficially, the disease is spoken of as "liver complaint" or "phthisis." An error in diagnosis is only excusable after we have exhausted all the methods at our command to arrive at a conclusion. In no case of disease in the child should an examination of the chest be omitted, and if there be any obscurity in the symptoms, such an examination is imperatively called for. Even when the examination is made with all care, an error may still be fallen into, and the disease be confounded with pneumonia. This mistake is hardly to be wondered at when we remember the resemblance between the physical signs of the two diseases—a resemblance which is by no means so generally recognised as its importance requires. If, however, we consider the whole aspect of the case, noting the manner of invasion of the disease, and comparing the physical signs with the general symptoms, this, with accurate observation of the character of the signs, will in most cases enable us to avoid an error.

The initial symptoms—pain in the side or belly, followed by cough, with usually fever, but only slight reduction of

the strength—should at once suggest pleurisy, and these symptoms are tolerably constant. There is little disturbance of the natural relation between the pulse and the respiration ; and the face, unless the pain be severe, seldom shows evidence of distress. In pneumonia the pain in the side, if present at all, is slight, and occurs subsequently to the cough, the pulse-respiration ratio is greatly perverted, the prostration is as a rule extreme, and the child has the aspect of one seriously ill.

When effusion has taken place, the dead, flat percussion-note with greatly increased resistance, the cavernous or blowing or merely weak bronchial breathing, and the increased resonance of the voice or cry—these signs discovered at the base, and *found at the front of the chest as well as at the back*, are very characteristic, and if the disease be an acute primary one, are diagnostic of pleurisy. If, however, there is a history of protracted ill-health, such physical signs may not be necessarily indicative of fluid in the chest.

In cases where the lung is the seat of fibroid thickening and the bronchi are dilated, a secondary pneumonia may spread with great rapidity, so as quickly to involve the whole thickness of the lung. Dulness with increased resistance would then be found all over one side of the chest both in front and behind : the breathing would be loud and cavernous, and the resonance of the voice and cry would be intensified. Still, in such a case we may be saved from error by the history of the case, and by remarking that there is no displacement of organs, or that displacement, if present, is found in a direction opposite to that which obtains in cases of pleurisy. Moreover, in the case of a secondary catarrhal pneumonia, abundant large, metallic rhonchi are heard with the breath-sounds. In pleurisy adventitious sounds are usually absent ; but if a little friction be present, it is heard close to the ear as a fine crackling sound, which has none of the metallic quality of bubbles generated in a rigid dilated tube.

Again, in pleurisy an ægophonic quality in the transmitted voice is common ; in pneumonia ægophony is never heard.

In a primary basic pneumonia the physical signs are limited at first to the back or the front of the chest, and are not found in both situations ; and besides, the puff of fine crepitation caught at the end of a deep breath is seldom absent in pneumonia, and is very different from the larger superficial crackling sound produced in an inflamed pleura.

The well-marked cavernous breathing often present in cases of pleurisy in the young subject may be a source of confusion ; but if it be borne in mind that cavernous or loud blowing respiration, audible even at the extreme base of the chest, is compatible with a large collection of fluid, we shall no longer be perplexed by the phenomenon.

On account of the feebleness of vocal vibration of the chest-walls in children, no information is to be derived from the absence of fremitus, unless this is to be felt over other parts of the parietes ; nor, on the other hand, does a strong fremitus necessarily exclude the presence of fluid, for vibration of the chest-wall may persist even where the effusion is copious. In a boy, aged 10 years, lately a patient in the East London Children's Hospital, and from whose chest ten ounces of fluid were removed by paracentesis, vocal vibration before the operation was well marked, and was little, if at all, weaker over the diseased side than over the healthy part of the chest. So, also, alterations in the voice or cry present in themselves nothing characteristic unless the quality is markedly ægophonic, in which case only may the sign be considered presumptive of the presence of fluid in the chest.

When the effusion reaches high up in the chest the heart is displaced and the intercostal spaces are obliterated, and these are very valuable diagnostic signs. In such a case, too, the complete absence of rhonchus over a dull area of large extent is much more common in pleurisy than in other forms of chest disease, and if present in a case of acute consolidation of the lung would not persist for many days

in succession. Again, the alteration in the shape of the affected side, which becomes somewhat square in outline, as evidenced by the cyrtometer, is a very characteristic feature of an extensive pleuritic effusion, and one which is not found in cases of ordinary consolidation.

The want of accordance between the physical signs and the general symptoms is another striking feature in the disease. A child is said to be languid, to have lost his appetite, and to be losing flesh ; but he coughs little if at all, and walks about as usual. On examination half or the whole of one side is found to be dull on percussion, with loud bronchial breathing, and the heart is displaced from its natural position. In such a case the slightness of the general symptoms would be in itself an important element in the diagnosis.

The colour of the face is also a characteristic symptom. In pleurisy the cheeks have often a uniform straw tint, which is unlike the complexion of any other disease.

Although the physical signs which result from solidification of lung differ considerably, as a rule, from those which are due to a collection of fluid in the pleural sac, in certain rare cases great difficulty is met with in making a distinction. There is a condition occasionally present in the child in which the whole of one lung is solidified and filled with dry caseous matter. Moreover, the bulk of the organ is increased so that it encroaches upon neighbouring parts. As a consequence, the walls of the chest are pressed outwards and the heart is displaced. It is this bulging of the chest and displacement of the heart which creates the chief difficulty ; and as, in addition, the percussion-note is completely dull and the resistance to the percussing finger is increased, it is no wonder that such cases are apt to be mistaken for empyema. Such complete filling of the lung with caseous matter may be combined with general tuberculosis, and Virchow has described very similar appearances as a consequence of inherited syphilis. Fortunately, as such cases are

rare, the difficulty in separating them from empyema cannot often arise.

The alteration in the percussion-note which results from change of position cannot always be relied upon to distinguish pleural effusion from solidification of lung. No doubt, if this sign be present, the existence of fluid may be considered to be proved ; but the pleural sac may be full of liquid although the percussion-note remains dull whatever be the position in which the patient is placed. It is in the stage of absorption that this sign can be discovered with the greatest certainty. A child is found to have some dulness at one posterior base, extending into the infra-axillary region : resistance is increased : the respiration is fairly loud, and is bronchial behind, blowing below in the armpit : some fine rather superficial-sounding crackling is caught at the end of respiration : the history is obscure. In such a case where it is important, on account of treatment, to ascertain the exact lesion present, careful percussion of the affected side while the patient is in different positions will often clear up any doubt that may have been entertained as to the presence of fluid.

When the pleurisy is *localised* so that the physical signs are limited to one aspect of the chest, the resemblance to pneumonia is very close, and students almost invariably mistake such a case for one of inflammation of the lung. But the distinction is not a difficult one to make. The percussion-note has the dull, toneless quality, characteristic of the presence of fluid ; the sense of resistance is extreme, and the breath-sound is unaccompanied by crepitation at the confines of the dull area. As a rule, in cases of localised pleurisy, the respiration is merely harsh ; sometimes, however, it is blowing, but it is invariably weak as if generated at a considerable distance from the ear. The continued absence of all crepitation or crackling in such a case is important, and the fact that the physical signs referred to continue for a considerable time unchanged in a case where

there has been recent inflammation in the chest, is a strong argument in favour of their pleuritic origin.

When we have satisfied ourselves that fluid is present, we have still to decide upon its nature. An accumulation of serous fluid in the pleural cavity—unless the quantity be sufficiently large to interfere with the function of neighbouring organs—does not call for operative interference, but if the fluid be purulent, it should be removed without delay. In early life the effused fluid often becomes purulent very quickly, and usually after the lapse of two or three weeks from the beginning of the illness, pus is found with the exploring syringe. But there are exceptions to this rule, and we cannot say positively that the fluid is purulent because the case is three weeks old. The temperature is useless as a guide, for fever depends less upon the nature of the pleural contents than upon the constitutional peculiarities of the patient. So, also, the physical signs are the same whether the fluid be thick or thin. Bacelli, indeed, has attempted to show that the clear conduction of the whispered voice to the chest-wall may be held to imply that the fluid is still serous ; but in practice this sign has not been found to be trustworthy. Even the straw tint of the face, although, as a rule, more pronounced when pus is present, may be well marked in the case of a simple serous effusion. Clubbing of the finger-ends, however, is a more trustworthy guide ; and my experience agrees with that of Dr. T. Barlow, that if this condition be present, we may infer with some confidence that the fluid has become purulent. In any case, if there be doubt, we possess in the hypodermic syringe a ready means of deciding the question by withdrawing a specimen of the fluid.

On account of the appearance of the patient, empyema is often mistaken for pulmonary phthisis, and, no doubt, in a case of some standing, many of the symptoms are common to both forms of illness. In either disease the child is apt to be pale, haggard-looking, and feeble ; his flesh is wasted,

and his skin harsh and dry. He is troubled with a hacking cough, and a little exertion makes him pant. His appetite is capricious and his digestion bad. Even the physical signs may seem at first sight to favour the suspicion excited by the general weakness, for an examination of the chest shows want of resonance over a more or less extended area with—in many cases—a blowing or even cavernous quality in the breath-sounds. A closer scrutiny, however, will discover many important differences. In pleurisy the illness begins suddenly, with pain in the side followed by cough. The dulness of the chest is limited strictly to one side of the body ; it is complete with extreme sense of resistance, and, unless the empyema be loculated, occupies the front and back of the chest, reaching down to the extreme base. Moreover, the heart's apex is usually displaced by the pressure of fluid, and if the amount of purulent matter be large, the chest-wall is expanded. In a chronic case, however, there is usually a certain amount of contraction of the affected side, so that the shoulder and angle of the scapula will be seen to be appreciably lowered, and the spine in consequence to be a little curved. In pulmonary phthisis, when the disease is sufficiently advanced to simulate empyema, the opposite side is rarely found to be healthy. Over the affected area, although resonance is diminished, the percussion note is less completely dull and the sense of resistance less pronounced. More or less rhonchus will be heard with the stethoscope, probably large in size and metallic in quality—a râle very different from the finer superficial crackling sound often heard in a case of pleuritic effusion. Lastly, in empyema the vocal resonance may have an ægophonic quality. This is never the case in simple phthisis.

Pleurisy is sometimes suspected when not actually present. An enlarged liver may press upon the diaphragm on the right side and cause dulness and weak breathing at the base of the chest on that side both in front and at the back. But

dulness due to an enlarged liver reaches to a higher level in front than behind, and the quality of the percussion-note is in no way altered by an alteration in the position of the patient. Moreover, the change from dulness to resonance is not abrupt, but there is a tubular (or tympanitic) note at the point of junction where the thin border of the lung overlaps the upper margin of the liver. In pleurisy the dulness is higher behind than in front, and changes abruptly to resonance at the upper margin of the fluid. In most cases, too, change of position has a distinct influence upon the percussion-note, and the normal resonance at the right base is restored when the child is placed upon his left side.

The diagnosis between pleurisy and hydrothorax seldom presents much difficulty. It is true that the physical signs are identical—being due in each case to a collection of fluid in the chest cavity, but hydrothorax is usually double, and there is no accession of fever when the effusion occurs, as is almost always the case at the onset of a secondary pleurisy. Moreover, in a passive exudation resulting from some impediment to the circulation, evidence is usually to be obtained of similar effusions taking place in other parts of the body.

Between pleurisy and collapse of the lung the distinguishing points have already been discussed in a previous chapter. (See page 38.)

The *prognosis* in pleurisy depends less upon the nature of the fluid than upon the age and constitutional condition of the patient. In childhood the nature of the fluid, taken by itself, has but little influence in determining the issue of the illness, for if a serous effusion, which remains serous, is absorbed without much difficulty, a purulent collection can be readily evacuated by a simple surgical procedure. In either case, if the constitution be sound, recovery follows quickly. In the case of a serous effusion, the only danger which can arise is that due to excess of fluid. If by this means the heart be sufficiently displaced to interfere with

the passage of blood through the large vessels, the child will die unless relieved. It has seemed to me that young infants are more oppressed than older children by comparatively small collections of fluid. At any rate, under the age of six months pleurisy is a very fatal disease : and often at this period death occurs so early that the fatal ending cannot be attributed to the mere quality of the pleural contents.

If the patient suffer from a diathetic tendency, special dangers may arise. In tubercular and scrofulous subjects an empyema may give rise to an attack of tuberculosis. In strumous children, too, there is usually considerable fever ; nutrition suffers greatly ; and however thoroughly evacuation of the chest-cavity be carried out, the inflamed membrane recovers but slowly, and it is a long time before the secretion of pus is entirely arrested. If an empyema open spontaneously through the chest-wall, the case seldom does well if left alone, for the opening is rarely so placed as to drain the chest completely. Evacuation of the pleura through a bronchus has a far more favourable termination, and many cases where this accident occurs recover without further help. Often, however, evacuation by this channel is imperfect and requires to be supplemented by a counter-opening in a suitable situation. In all cases the persistence of a high temperature after discharge of the purulent effusion by operation, and fœtor of the pus, must be looked upon as unfavourable symptoms. Still, in the latter case, free evacuation and irrigation of the chest will often be followed by the happiest results.

The *treatment* of pleurisy at first is simple, and consists in perfect quiet, warmth to the chest, a carefully ordered diet, and the administration of a simple saline. The severe measures which used to be at once resorted to in the treatment of every case of pleuritic effusion, are now set aside for simpler remedies. Bleeding need only be mentioned to be condemned. The only form in which it would be for a moment admissible is that of the application of a leech to

the side to relieve pain ; but this is now unnecessary, as we have in the hypodermic injecting syringe an instrument which enables us to effect the same object rapidly, and without the withdrawal of blood. In pleurisy a child wants all his strength, and to lower it is to do him harm. Mercury, again, which used to be the indispensable remedy at a time when its influence in promoting the absorption of all serous exudations was held in universal belief, is now practically abandoned in the treatment of pleurisy. In children the drug should never be given unless the indications for its employment are very decided, on account of the anæmia and depression which it is so apt to induce ; and the best authorities now concur in believing its value as an absorbent in cases of pleuritic effusion to be more than doubtful.

The child should be kept perfectly quiet in bed. Hot linseed-meal poultices, frequently renewed, should be applied to his chest ; or the upper part of the body may be swathed, not too tightly, in a broad flannel bandage wound round and round the chest and crossed over the shoulders. His diet should be carefully suited to his powers of digestion, and should consist principally of milk and light broth. For medicine he may take a citrate of potash mixture or a draught containing a solution of acetate of ammonia with sweet spirits of nitre. If the pain in the side be severe, the hypodermic injection of morphia, in the proportion of one-thirtieth of a grain for a child of twelve months old, will give very rapid relief.

In children who are old enough to describe their sensations, very severe pain referred to the epigastrium and accompanied by marked oppression of breathing is usually indicative of inflammation affecting the pleura overlying the diaphragm ; and in such cases the action of this muscle during respiration may cause great suffering. When this occurs the application of a bandage to the belly, so as considerably to restrict the action of the diaphragm, will often give immediate ease.

If the cough be distressing, small doses of opium should be given by the mouth. It is important to keep the bowels properly relieved, for any distension of the stomach would increase the uneasiness of the patient. For the same reason a careful watch should be kept over the digestive functions, and any signs of dyspeptic fermentation of food should at once cause us to reconsider the dietary, and restrict the quantity of farinaceous or other fermentable matter which is being taken. Violent purgation is to be avoided, for, as in all acute chest disorders, irritation of the bowels seems to be attended with anything but benefit to the patient.

When effusion into the pleura has ceased, measures must be at once adopted to hasten absorption of the fluid. Counter-irritation of the chest-wall has long been extensively employed to effect this object, and although the value of the practice has been called in question, in this disease it has always seemed to me to yield very satisfactory results. I have certainly met with many cases where fluid, which under treatment previously adopted had not perceptibly diminished in quantity, has begun at once to be absorbed immediately that counter-irritation was made use of, and this although the treatment remained in all other respects the same. The liniment of iodine is the most convenient form of counter-irritant. A spot about the size of the palm of the hand should be well painted over with the liniment twice a day until the skin begins to crack and peel. Another spot of the same size should then be selected, and the process be repeated in exactly the same manner.

At this time the child should be put upon as dry a diet as possible, for by diminishing the quantity of fluid taken into the body the blood is stinted of necessary liquid, and a stimulus is thus given to absorption. For medicines the iodide of potassium is very useful if given in full doses. To a child of five years old fifteen grains of the salt may be given with perfect safety every four hours.

Quinine, again, is useful if given in sufficient doses. A

child of five years old will take three grains every six hours without inconvenience if proper attention is paid to the bowels; and the effect of the remedy will often be seen in the rapid removal of fluid from the chest. Iron, in the form of the tincture of the perchloride, is also of service, and should be given in tolerably large doses—five to twenty drops, according to the age.

In ordinary cases, when the disease has occurred in a child of healthy constitution, the above measures will be found to induce an early improvement in the physical signs of the chest. If, however, our expectations of amendment are disappointed, it will be necessary to employ more direct means to relieve the patient. In early life a pleuritic effusion quickly becomes purulent. Therefore, if no signs of absorption have been noticed by the end of the third week from the beginning of the illness, it is best to make an exploratory puncture to ascertain the nature of the fluid. If the examination shows the presence of pus, steps should be taken at once to evacuate the pleural cavity. If, however, the fluid withdrawn by the test puncture be clear, it is well to wait before interfering further. In an ordinary case of serous effusion it is unnecessary to employ mechanical means for emptying the pleural cavity. In most cases the small quantity of fluid removed by the syringe proves a sufficient stimulus to absorption, so that the remainder rapidly disappears by the ordinary processes of nature. It is only when the quantity of serous fluid collected constitutes a danger to the child that operative interference is called for. Thus, if, without causing actual attacks of dyspnœa, the accumulation of fluid be so great as to press upon the opposite lung and hamper the breathing, or if there have been several fits of dyspnœa when the collection of fluid is large—in either case the operation should be performed without unnecessary delay.

If it be decided to empty the chest, that mode of proceeding is to be preferred by which the evacuation of the

pleural cavity can be most completely carried out. For this reason in hospital practice, when the presence of pus in the pleural cavity has been ascertained, I usually apply without loss of time to a surgical colleague, and beg him to fit a drainage-tube into the chest-wall. This operation is unattended with risk to the child and allows of complete escape of the effused fluid. In private, however, where others besides the patient have usually to be considered, it may be judicious in the first place to employ the aspirator, and empty the pleural sac of its fluid, more or less completely, by this means. It sometimes happens, indeed, that the use of this instrument is followed by the happiest results. I have met with several cases, in which, after removal of the fluid by the aspirator, no more pus has been thrown out, and the physical signs of the chest have quickly become normal. But these cases are unfortunately exceptional; in most instances the chest fills again as often as it is emptied, and in the end the drainage-tube has to be resorted to.

In using the aspirator to withdraw the fluid, or in searching for pus with the exploring needle, a spot should be selected where the dulness is complete and the resistance well marked: also, a point is to be preferred where the intercostal space is noticed to bulge, for here the false membranes are probably scanty and thin.

If a fine needle be used, the introduction of the instrument causes little pain, and anæsthetics are unnecessary. The chief discomfort during the operation arises from the cough, which often becomes very violent and distressing. This appears to be the result of the irritation of the pleura set up by the instrument. Trousseau's explanation, that it is a consequence of the irritation of air entering the disused lung, is scarcely a sufficient one, for the severity of the cough is not proportioned to the amount of effusion, and therefore to the degree of compression of the lung, but may be very distressing although the quantity of fluid in the

chest is scanty. The fluid has at first the ordinary appearance of purulent matter, but if the operation be performed by the aspirator, it may become after a time tinged with blood, and often at the end of the tapping seems to consist of pure blood. When this is the case, the needle should be at once withdrawn. The blood does not come from a puncture of the lung, but is probably, as Trousseau suggests, poured out by some ruptured vessels of the pleura, or of false membranes which were becoming organised. Even if the lung be punctured during the operation, and air be extravasated into the pleura, no apprehension need be excited by the accident, as traumatic pneumo-thorax thus produced appears to be accompanied by little inconvenience, and soon disappears.

CASE XVII.—TRAUMATIC PNEUMO-THORAX, FROM PUNCTURE OF THE LUNG DURING THE OPERATION OF PARACENTESIS.

Matilda H., aged 6 years, was admitted into the East London Children's Hospital on April 2, after suffering for a fortnight from cough and some fever. All over the left side there was complete dulness on percussion, with weak bronchial breathing, and absence of vocal resonance. The respiration was almost entirely abdominal, and the heart was pushed to the right. The child looked well, and had not the characteristic complexion of pleurisy, but her temperature was high in the evening, rising usually above 101° . On April 16, the thirtieth day of the disease, the child was punctured with the pneumatic aspirator, and eight ounces of thick greenish yellow inodorous pus were withdrawn. At the end of the operation there was a gush of air through the cannula, and on withdrawal of the instrument the following physical signs were noted :—"Percussion amphoric all left side ; respiration loud and amphoric, with metallic tinkle after cough ; heart's apex immediately

below nipple." The temperature that evening was 100° . One week afterwards the air had completely disappeared from the pleura, and except for some slight dulness at the base, and also at the supra-spinous fossa, with bronchial breathing, and superficial clicking sounds, the physical signs were natural. When the child left the hospital, on May 25, the physical signs were those of health, except that the breathing remained very weak on the affected side, and that there was some slight curvature of the spine, with the concavity towards the left. Her weight was found to have increased by three pounds. In this case there can be little doubt that the lung was punctured by the aspirating needle, and that traumatic pneumo-thorax was produced. Still no inconvenience to the patient seemed to be occasioned by the accident; the temperature that evening was only 100° , and in a week the air was completely re-absorbed.

In using the aspirator or exploring needle we must not be discouraged if, on the first trial, no fluid is withdrawn. There are many reasons why fluid, even when present in large quantity, may fail to appear. The fluid itself may be too thick to pass readily through the fine tube of the exploring needle, or may contain flakes and shreds which block up the channel; the membranes may be so tough that they are pushed before the point of the needle, so that the latter fails to penetrate into the sac: the empyema may be loculated, and at the point where the needle enters no fluid may be present: lastly, the effusion may be prevented from escaping by rigidity of the walls of the thorax.

In childhood the flexibility of the thoracic wall is much greater than it becomes in after life. There is also a readier power of expansion in the compressed lung-tissue in the child than in the adult. It is, indeed, to the existence of these two properties that the commonly favourable termination of empyema in the young subject is to be ascribed. If, however, from any cause the chest-wall cannot contract under the pressure of the atmosphere, while at the same

time the condensed lung is not able immediately to re-expand, no pressure is brought to bear upon the confined fluid to force it out of the chest. Under such circumstances no amount of suction force will be successful in withdrawing the purulent contents of the pleura, for, as my colleague, Mr. R. W. Parker, has reminded us, in emptying an empyema cavity the fluid is not drawn out but driven out, and the driving force is always the direct pressure of the external air. If, then, in a case where the existence of purulent fluid has been ascertained, a difficulty is met with in evacuating the pleura, it is advisable to make free incision in an intercostal space and insert a good sized drainage tube into the chest wall.

The operation of opening the chest by free incision is usually performed with all antiseptic precautions ; but I do not think that these can be indispensable to success. In the days when we were contented with fixing between the ribs a roomy silver cannula, through which the air whistled at each inspiration, and covering the opening with a large pad of tow to soak up the exuding discharge, the issue of the case was rarely otherwise than favourable. I do not know that in these days, with our improved appliances and scientific precautions, we can boast of any better results.

When it is intended to leave a drainage tube in the chest-wall, the point fixed upon for the opening is of importance. If it be too low down, the rising of the diaphragm, as the side contracts, may displace the tube or interfere with the escape of the fluid. The most convenient level is the ninth interspace, and the spot chosen should be at some little distance from the angle of the scapula, as otherwise the movements of the shoulder may retard the closing of the wound after the tube has been withdrawn. If the operation is performed early, and the evacuation of the pleural cavity is complete, a cure usually follows without great delay. The discharge from the cannula diminishes rapidly in quantity, so that after a few weeks the tube can be withdrawn and the

opening allowed to close. If the case do badly, the unfavourable issue is usually due either to constitutional weakness, such as a pronounced strumous diathesis; to the operation having been injudiciously delayed so that the lung has become bound down and cannot re-expand; or to an awkward position of the opening which hinders or prevents the escape of the fluid.

If there be any *foetor* of the pus the chest should be washed out daily with a very weak solution of iodine (3 ss. to 3 j. of water) or chloralum (one part in fifteen of water). In all these cases the utmost care should be taken to avoid a new chill. The patient should be anxiously guarded from sudden changes of temperature, and it is useful to keep the chest covered with a long flannel bandage.

The amount of deformity, even in cases where an opening in the chest has been maintained for a considerable time, is often very much less than would be expected, and the rapidity and completeness with which such deformities may be repaired is often very surprising. If the operation be performed sufficiently early, and the chest be completely relieved of its fluid contents, permanent deformity rarely ensues. Even a temporary contraction of the chest by this means is usually averted. It must be remembered that contraction of the side following upon a pleural effusion is no proof that the chest cavity no longer contains fluid. Indeed, the contrary is the fact; for, as Dr. Barlow has pointed out, the occurrence of deformity in such a case tends directly to suggest retention of purulent matter at the base of the lung or between the pulmonary lobes.

If the case be seen for the first time when the empyema is pointing, the swelling must be lanced, so as to allow of the escape of the fluid. If, however, the skin be not very red, the chest may be punctured behind, in the ordinary situation, and the threatened abscess disregarded. In such cases the pus soon disappears from under the skin and the swell-

ing subsides. The same plan may often be adopted with advantage, even if the empyema have already burst high up in the chest. Where this has occurred, the fluid being retained in the pleural cavity, and escaping only with difficulty, is often intensely offensive: the introduction of a drainage tube below efficiently drains the pleura, allows of thorough cleansing of the cavity, and permits the upper opening to close. Weak injections of iodine in these cases effectually control the foetor, and the patients often do very well.

In all cases of empyema, especially after the fluid has been evacuated, the diet should be as nutritious as possible. Meat, eggs, and milk should be given as freely as the child's powers of digestion will allow, but the stomach should not be overloaded with too large a quantity of food at one time. Especial care, indeed, should be taken not to *overfeed* the patient in our anxiety to supply material necessary to enable him to withstand the exhausting drain upon his system caused by the purulent discharge. For medicines, quinine and iron are especially indicated. Cod liver oil, too, is very useful, and I have found the natural light brown oil prepared by Mr. Kerr, of New Bond Street, to be more completely digested by weakly children than the over-purified colourless oils in common use. M. Duboué recommends the administration of tannin after evacuation of the fluid in empyema. He states that this drug acts as a tonic astringent, improving digestion and diminishing secretion. It may be given in doses of from three to five grains several times in the day. Small quantities of wine or brandy are also required in these cases.

The above method of treatment is also applicable to cases of secondary pleurisy accompanied by effusion of fluid. In dry pleurisies, such as are common in cases of phthisis, the pain is best relieved by mustard poultices, or by small blisters if the child be old enough.

CHAPTER V.

ACUTE CATARRHAL PNEUMONIA (BRONCHO-PNEUMONIA—LOBULAR-PNEUMONIA).

ACUTE catarrhal pneumonia especially the pneumonia of infancy—Reasons
—The disease differs from croupous pneumonia pathologically as well as clinically.

Pathology—Modes of production—Often preceded by collapse of the lung
—Symptoms and physical signs.

Case XVIII.—Ordinary acute catarrhal pneumonia in an infant.

Case XIX.—Acute catarrhal pneumonia secondary to collapse of the lung—Comment.

Case XX.—Subacute type of catarrhal pneumonia—Comment.

Favourable termination.

Case XXI.—Gradual subsidence of acute catarrhal pneumonia—Comment.

Bronchiectasis a common consequence of acute catarrhal pneumonia—may occasion difficulty in diagnosis.

Case XXII.—Acute catarrhal pneumonia with dilatation of bronchus—Comment.

Diagnosis—Prognosis—Treatment.

IN a preceding chapter we have seen that croupous (lobar) pneumonia may be either a primary or a secondary disease. It may either attack a perfectly healthy subject, or may occur as a complication in the course of some other illness. Catarrhal, or lobular, pneumonia is, on the contrary, nearly always a secondary lesion. In almost every case it is preceded by severe pulmonary catarrh, of which it is one of the direct consequences.

Catarrhal pneumonia is peculiarly the pneumonia of infancy. This, according to Rindfleisch, is the consequence of the greater irritability of the epithelial elements in the child's lung. The cells are larger, contain more protoplasm,

and adhere less firmly to the alveolar surface than they do in after life. In the young subject the catarrhal process extends very rapidly and readily from the minuter bronchi to the alveolæ; and therefore, on account of the proneness of the child to catarrhal diseases, especially to pulmonary catarrh, this form of pneumonia is of great frequency, and is much more often met with than the croupous form, which is the more common lesion in the adult.

The distinction between croupous and catarrhal pneumonia is one which lies not only in the clinical character of the two diseases, but also extends to the nature of the pathological process. In croupous pneumonia a highly fibrinous liquid escapes from the blood-vessels, accompanied by large quantities of both red and white corpuscles, and coagulates in the air-cells of the lung. The process appears to consist more in extravasation than in exudation; for although a certain proportion of the corpuscles may pass bodily through the walls of the vessels, as stated by Cohnheim, yet there is little doubt that actual rupture of the blood-vessels, and escape of their contents, is an important feature in the process. Whether the coagulable matter be merely blood plasma, or consist of that fluid altered in character during its passage from the blood; whether, again, the cells with which the alveoli are choked consist entirely of blood corpuscles, or are composed in part of a rapid multiplication of the cells and nuclear elements of the tissues themselves,—these are questions which, although interesting to the pathologist, are yet comparatively unimportant to the practical physician. The great point of interest to him lies in the fact that in croupous pneumonia the cellular elements filling up the vesicles are exuded or extravasated from the blood.

In catarrhal pneumonia, on the contrary, the cells which crowd the air-vesicles are not derived from the blood at all, but are produced in the cells themselves by a proliferation of the alveolar epithelium. These epithelial cells rapidly

multiply and enlarge, and are embedded in a semi-fluid mucoid material which contains little or no fibrin such as is found in the spontaneously coagulable matter which holds together the corpuscles in croupous pneumonia. The process, which is a catarrhal one, takes place in detached lobules scattered through the lung, and the appearance presented to the naked eye is that of reddish-grey nodules dispersed over the organ. The colour afterwards changes to yellowish-grey, for the cells, which rapidly enlarge and develop one or more nuclei, soon undergo a fatty metamorphosis. If the fatty change be completed, absorption takes place, and the consolidation is removed; if it remain incomplete, the cells atrophy, the little masses become caseous, and the disease passes into a chronic stage. But besides the changes in the air-vesicles, the bronchi themselves participate in the disease. The walls of the air-tubes become thickened and their calibre often dilated. Acute dilatations of the bronchi from this cause are very common in children, and the physical signs to which such dilatations give rise may occasion some difficulty in the diagnosis.

This form of pulmonary inflammation is often the result of an extension of the catarrhal process from the bronchi to the air-vesicles, but it may arise in another way. Collapse of the lung is a very common complication of bronchitis in children, and a collapsed state of the air-vesicles greatly favours a rapid multiplication and development of the alveolar epithelium cells. In part through the imperfect aëration of the blood, and in part through the absence of the ordinary respiratory movement in the affected tissue, a great impediment is created to the passage of the blood through the capillaries of the collapsed portion of the lung, and a passive congestion is set up. This is soon followed by œdema. Inflammation then occurs at separate points, which may remain isolated, being divided by the congested and œdematous tissue, or may coalesce until a large portion of the lung becomes solidified.

From the above short and imperfect sketch of the pathological processes which give rise to the disease, it is evident that catarrhal pneumonia must almost always be a secondary inflammation. It seems probable that the occurrence of this form of pneumonia is favoured by anything which reduces the strength, or which diminishes the respiratory muscular force of the child. Therefore, bad feeding and hygiene, all diseases where catarrh is a prominent symptom, and in fact all diseases which require a recumbent position to be long maintained, and so cause congestion of the posterior parts of the lungs, are predisposing causes which render more probable the occurrence of this complication.

The differences in the method of production of catarrhal pneumonia—whether or not it is preceded by collapse—have an important bearing in a clinical point of view. The combination with collapse is usually found in cases of pertussis, and the disease is then much more insidious in its invasion, and often runs a subacute and very tedious course. The complaints in which catarrhal pneumonia is especially liable to occur are bronchitis, measles, and pertussis. Severe capillary bronchitis is very rarely left uncomplicated by catarrhal inflammation, and the pneumonia usually in such cases begins early, especially if rickets be present; for this disease, if at all severe, is always accompanied by a certain amount of collapse of the lung. The softening of the ribs, which is so prominent a symptom in severe rickets, offers a great impediment to the entrance of air even if the lungs be healthy; but if the passage of air be further hindered by the addition of pulmonary catarrh, a full inspiration becomes impossible. The patient is, therefore, unable to take in a sufficient supply of air to furnish the means for a successful cough, and the mucus, not being expelled, accumulates in the tubes and causes collapse. The mechanism of collapse and the causes which predispose to it have been already fully considered in a previous chapter.

The course of catarrhal pneumonia is as follows:—After

pulmonary catarrh has existed for some little time, a sudden change takes place in the character of the symptoms. The temperature rises, but not to such a height as is found in the croupous form of the disease; the thermometer is usually found to mark about 102° . The breathing, which was before hurried, becomes very rapid and laborious, and there is intense dyspnoea. The efforts made to draw in the air are shown by the widely-dilated nares and the violent action of all the accessory muscles; while the insufficient distension of the lungs is evidenced by the great recession of the lower part of the chest-walls and by the sinking in of the intercostal spaces. In fact, although the breathing is so laborious, it is yet shallow. The inspirations are short and imperfect, the expirations noisy and prolonged. This rapidity in the breathing produces the same alteration in the ratio between the pulse and the respiration which is seen in lobar pneumonia. A ratio of 1 to 2, or even of 1 to 1.5, is not at all uncommon; and this varies at different times in the day, for the dyspnoea is sometimes paroxysmal, and occasionally becomes intensified to such a degree as to appear to threaten immediate suffocation. The cough also alters its character. Instead of the loose prolonged cough of bronchitis, we get the short hack of pneumonia. It often occurs in paroxysms, each expiration being accompanied by a short dry cough, which is evidently painful from the distress visible in the face of the patient. At the same time the secretion from the bronchi is diminished, for the rattling of mucus in the chest at each breath, which is so common a feature in bronchitis, can no longer be detected. The pulse is rapid, but small and feeble. Its weakness is no doubt partly due to comparative emptiness of the arteries, on account of the impediment to the passage of the blood through the lungs; and the same explanation applies to the fulness of the superficial veins which is so often observed in the disease.

Examination of the chest often yields no very positive

results. As long as the pneumonia remains confined to isolated patches there is seldom any dulness on percussion, but sometimes we can detect a general want of healthy pulmonary tone over the back, and this is more marked on employing broad percussion, striking with three fingers upon three fingers placed upon the chest-wall as pleximetres. Occasionally we find a localised spot of dulness with very limited bronchial breathing, and sometimes the breathing may even approach to the tubular character when the affected portion of lung is of some size from coalescence of several inflamed patches.

The following case well illustrates these symptoms :—

CASE XVIII.—ORDINARY CASE OF ACUTE CATARRHAL
PNEUMONIA IN AN INFANT.

L. W., a little boy, aged five months, had had a bad cough, and was feverish for a fortnight. On July 6 he suddenly became very much worse. He was seen on July 7. Skin very hot ; respirations very rapid ; face livid ; nares acting ; cough almost constant, short and hacking. There was great recession of the chest-walls at the lower part, and the intercostal spaces of both sides sank in deeply at each inspiration. No dulness was found on percussion of the back at either side, and the breathing seemed natural. In the left infra-axillary region, a little below the level of the nipple, a spot of dulness could be detected, and the respiration there was bronchial, approaching to tubular. Everywhere else the breath-sound was natural.

A saline mixture, containing a few drops of antimonial wine, was prescribed, and the whole chest was ordered to be kept covered with linseed-meal poultices.

The child went on well for a few days, when early on the morning of July 14 he was taken suddenly worse. Breathing excessively rapid ; nares acting ; much lividity of lips and eyelids, and a general straw-tint all over face ; eyes

wide open and turned upwards; fontanelle rather elevated and tense; pulse 160, excessively weak; temperature in rectum 102°. Great recession of chest-walls and sinking in of intercostal spaces. The child had been a little convulsed, but had had no regular fit. On examination of the chest no dulness was detected, but the respiration was heard to be tubular in spots about both sides of back. Convulsions came on violently in the evening, and the child died.

It is especially in cases where the pneumonia arises from an extension of the inflammation from the bronchi to the air-vesicles, or where the inflammation occurs in patches of lobular collapse, that the absence of positive dulness is to be noted. When an extensively collapsed portion of lung becomes inflamed, there are usually signs of considerable consolidation, and the dulness is apt to take a form which is very characteristic, presenting itself as a narrow strip along the side of the vertebral column, reaching upwards a variable distance from the base. As the collapse is usually symmetrical, this pyramidal form of dulness is found on both sides. To detect it, it is necessary to percuss lightly.

Over the parts of the lung unaffected by the pneumonia the physical signs are those of the bronchitis which is present; the respiration is exaggerated and coarse, and is accompanied by sonoro-sibilant and bubbling rhonchi. As the stethoscope is moved towards the consolidated spots the bubbling rhonchus becomes finer and drier in character, and may even assume the fine puffy character of true pneumonic crepitus if the consolidated part is of any magnitude.

CASE XIX.—ACUTE CATARRHAL PNEUMONIA SECONDARY TO COLLAPSE OF THE LUNG.

George C., aged one year and nine months, a delicate child, who had been troubled for some weeks with looseness of the bowels, was seized with measles on November 21. The cough was very violent, and continued after the erup-

tion had subsided. On December 5 the boy seemed so ill that he was brought to the hospital, and was at once admitted as an in-patient.

On admission, the child was very quiet, and seemed exceedingly ill. His face was pale, with some lividity round the mouth and beneath the eyes; the nares dilated widely at each breath, and the respiration was rather laborious. There was some recession at the epigastrium, and the intercostal spaces of both sides sank in deeply in inspiration. The fontanelle, which was the size of a shilling, was depressed, and the pulse was small and feeble. The child had been lately sick; the bowels were still relaxed; and his tongue was thickly furred. Temperature 100.4° . On examination of the chest, dulness was discovered at the left back reaching from the base as far as the lower angle of the scapula; the respiration there was tubular, and much fine subcrepitant rhonchus was heard. There was also dulness at the extreme base on the right back; the breathing was bronchial, and fine subcrepitant rhonchus accompanied both expiration and inspiration. In front there was no dulness on either side, and ordinary dry and large bubbling rhonchi alone were heard. The child was perfectly quiet during the examination of his chest, and allowed himself to be held in any position without complaint.

On the following day, December 6, the dulness reached rather higher at the left back, and the crepitation there was finer and more puffy in character; the vocal resonance also was increased. At the right back the dulness reached as high as the inferior angle of the scapula; the respiration was bronchial, and inspiration was accompanied by a fine puffy crepitation. Temperature in morning 101.8° , pulse 152, respirations 44; temperature in evening 102.6° .

On the evening of December 9 the temperature was as high as 103.4° , and rose to 105° on the following morning. A few hours afterwards the child died.

At the post-mortem examination there was found solidi-

fication of the lower lobe of each lung, extending upwards in a pyramidal form from the base, so that the narrowest part of the consolidation was at the upper border of the lobe. The solidification did not extend the whole thickness of the lung, but occupied only the posterior surface ; in front the lobe appeared healthy, and crepitated on pressure. The consolidated parts were of a purplish-red colour, but were not depressed. The section was deep-red in colour, with small greyish patches scattered over the surface—evidently isolated spots of catarrhal pneumonia. The tissue was not tough, but broke down pretty easily under the finger. Much purulent bloody fluid escaped on pressure.

The catarrhal pneumonia from which this child suffered was no doubt secondary to collapse of the lung. The boy was in a weakly state from diarrhœa when he was attacked by measles. The accompanying catarrh was severe, and induced symmetrical pulmonary collapse, which afterwards became the seat of catarrhal pneumonia. The consolidation in this case was of some extent ; but the presence of bronchitic râles over the lungs generally, the character of the crepitation heard over the seat of consolidation, and the fact that this and bronchial or even blowing breathing were heard in the same spot, marked out the case very distinctly from ordinary croupous pneumonia. Again, in the diagnosis of catarrhal pneumonia the exact nature of the lesion may be determined by attention as much to the general as to the local features of the disease. In the present case the face was pale instead of being flushed, the two lungs were attacked simultaneously, the breathing was laborious, and the temperature until a few hours before death reached no very great elevation. In all these points, therefore, the disease differed from an ordinary case of croupous inflammation. Besides, the age of the child was of importance ; for although croupous pneumonia may attack infants of even a few months old, yet inflammation of the lung is much more commonly catarrhal than croupous under the age of two

years. Lastly, the form of pneumonia which occurs secondarily to measles is in the large majority of cases catarrhal. For these reasons the case was looked upon as one of catarrhal pneumonia, and the diagnosis was confirmed by a post-mortem examination of the body.

Children suffering from a severe form of this disease are singularly impassive and regardless of what they would under ordinary circumstances highly resent. They seem not so much stupid as unaffected by ordinary sources of annoyance. Thus, in the child whose case has just been referred to, there was no lack of intelligence in the expression of the eye, but the apathy he showed under examination, and the indifference with which he allowed himself to be shifted from one position into another to suit the convenience of the physician, was a striking feature in the case. When this apathy is carried to such an extreme degree, the prognosis is very unfavourable.

The course of catarrhal pneumonia is not always so acute as in the cases above described. The inflammation often progresses much more slowly, and, indeed, when the disease does not prove fatal within the first few days it usually takes on a subacute course, which may continue for several weeks before reaching its termination. In this form of the complaint the general symptoms are less severe; the skin, although hot, is often moist, and the physical signs may undergo little change for days or even weeks together, the pulmonary lesion appearing to be almost stationary. The child, however, gets weaker and thinner, paler, and more and more apathetic, and death may be preceded by extreme emaciation.

CASE XX.—SUBACUTE TYPE OF CATARRHAL PNEUMONIA.

Jane S., aged fifteen months, two teeth, was admitted into the East London Children's Hospital on March 16.

There was consumptive history on the mother's side, and one other child was said to have died from phthisis. The patient had been wasting more or less since birth, and had been subject to frequent attacks of diarrhoea ; her bowels, indeed, were habitually loose, with slimy offensive motions. Her diet had consisted principally of corn-flour, but she had been fed besides with portions of the food of her parents. She had never walked. For five weeks she had had a cough, which latterly had become worse, and she had begun to be feverish. When admitted, the patient was pale, thin, and miserable-looking, but the expression of the face was not actually distressed. Her limbs were excessively small and flabby. The joints were not nodular, but the ends of the ribs were beaded. Her chest was of normal shape, without distortion ; there was considerable recession of the lower parts on inspiration, and the intercostal spaces sank in deeply. There was no lividity of the face. Fontanelle size of a sixpence, depressed. Circulation feeble ; hands and feet cold, and pulse weak and small. Temperature in rectum $103\cdot4^{\circ}$. On examination of the chest some dulness was found, with slightly increased resistance at each base behind, but more marked and more extensive on the left side than on the right. The respiration was blowing, but not tubular, and was accompanied by a fine crepitating rhonchus. The resonance of the cough was stronger than natural.

On March 17, in the morning, the temperature was $101\cdot8^{\circ}$, pulse 112, respirations 36, and during the following days the temperature usually varied between 101° and 102° , although it sometimes rose to nearly 103° in the evening. The pulse, too, altered little in rapidity from day to day, varying between 120 and 140, but the respirations increased progressively in frequency, rising gradually to between 50 and 60 in the minute. The physical signs underwent but little alteration : the dulness became slightly more pronounced, and crept up rather higher at each back ; the

respiration assumed more of a sniffling quality, but never became perfectly tubular; the rhonchus lost much of its crepitating character, and became more bubbling, and the vocal resonance was broncophonic.

On March 24 some want of tone on percussion was noted at each supra-spinous fossa, especially the right, and the respiration at these spots was high-pitched and wheezing, with some subcrepitant rhonchus. At the base the physical signs remained the same, except that the respiration in each infra-axillary region had become slightly bronchial, especially on the right side, where there was some fine bubbling rhonchus, which disappeared after a cough; the resonance of the cough, too, was increased. There was, however, no dulness laterally, the alteration in the percussion-note ceasing on each side at the level of a line drawn perpendicularly downwards from the posterior fold of the axilla. In front, with the exception of some dry râles, there were no physical signs throughout the course of the illness.

The child got very thin, and towards the end her bowels became relaxed and she wasted very rapidly; her restlessness changed to dulness and apathy, and she gradually passed into a somnolent state, in which she died on March 26. Temperature shortly before death, 102.2° ; pulse 156; respiration 56.

On examination of the body, the lower parts of both lungs were found the seat of catarrhal pneumonia. The inflamed patches had in many places become caseous, and in some had softened down into a thick purulent liquid: the tissue between them was collapsed in many spots. Much thick purulent secretion was found in the bronchi. The upper lobes were emphysematous. There were no grey granulations anywhere about the body, and all the other organs appeared to be healthy.

In this case the child was a weakly, unhealthy-looking infant, the subject of rickets, and the only difficulty was to

decide upon the presence or absence of grey tubercle. There was a history of phthisis in the family, and therefore a constitutional tendency to caseous degeneration, but this did not necessarily imply any special proclivity to true tubercular disease, although indirectly it might be the cause of such a condition, for acute tuberculosis appears to be in many cases a secondary formation induced by general infection of the system through softening cheesy matter. There was, however, ample cause in the history of the present patient to account for the miserable condition into which she had fallen. She had been fed improperly almost from her birth ; that is to say, her system had been stinted of suitable nourishment on account of the indigestible form in which food had been supplied to her ; and, in addition, she had suffered from an habitual looseness of the bowels, which had still further lowered her strength and reduced a system ill-prepared to support so exhausting a derangement.

In such a state a child falls an easy victim to secondary acute diseases, and the condition of the patient was one especially favourable to the development of catarrhal pneumonia. The respiratory muscles necessarily shared in the weakness of the whole muscular system, and her enfeebled state necessitated an almost constantly recumbent position, which tended to promote congestion of the posterior parts of the lungs.

The disease in this case was evidently an acute one, for the child had only coughed for five weeks previous to her admission into the hospital ; and the consolidation was quite recent, for when the patient was first seen the dulness was not extensive, and it spread gradually after she had come under observation. The chief distinction between this case and one of ordinary croupous pneumonia lay in the character of the physical signs. The breathing was never tubular ; the rhonchus differed much from true pneumonic crepitation, and, besides, remained audible over the

seat of consolidation, even in spots where the breathing was most markedly blowing. The disease was also bilateral, affecting corresponding parts of both lungs ; the temperature was lower than is common in croupous pneumonia ; and, finally, the subacute course of the disease was very characteristic of its catarrhal origin.

When first seen the case might perhaps have been mistaken for one of croupous pneumonia undergoing resolution ; but a reference to the pyrexia would at once have shown that this could not be the case, for in croupous pneumonia the critical fall of temperature precedes by several hours any improvement in the physical signs. Afterwards, the gradual spreading of the signs of consolidation and the persistent elevation of temperature pointed to the true nature of the illness.

It would probably have been impossible in this case positively to exclude tuberculosis, for the catarrhal pneumonia might very well have occurred secondarily to the grey granulation, and there was sufficient evidence in the family history and the personal experience of the child to warrant such a conclusion. Still, the theory of tuberculosis was not necessary to the full explanation of the symptoms ; and partly for this reason, partly on account of the absence of the haggard, wearied expression so characteristic of the constitutional disease, and partly also on account of the seat of the consolidation—which was limited to the posterior base of the lung at each side—it was thought probable that this case was one of uncomplicated catarrhal pneumonia following upon a long period of malnutrition.

When catarrhal pneumonia ends favourably, improvement is gradual, and there is no marked crisis such as we see in croupous inflammation of the lung. In the croupous form of pneumonia the general symptoms improve before the local, and a sudden fall in the temperature is the earliest sign of approaching convalescence. In catarrhal pneu-

monia, on the contrary, the local symptoms are the first to show signs of amendment; the breathing becomes less hurried; the cough less harrassing; the superficial veins less prominent; the pulse fuller; and the expression more placid; while all the time the pyrexia continues, and it may be many days before the evening temperature sinks to the natural level of health. The physical signs also improve, but often very slowly, and in cases where the disease has run a very subacute course, weeks, or even months, may elapse before the consolidation is completely removed. This form of pneumonia is especially apt to leave behind it caseous deposits, which, after remaining inactive for a variable time, may begin to soften, and lead to ulcerative destruction of the pulmonary tissue (pulmonary phthisis). The following case illustrates the very gradual improvement which takes place when the disease terminates favourably :—

CASE XXI.—FAVOURABLE ENDING OF ACUTE
CATARRHAL PNEUMONIA.

George C., aged six years, was admitted into the East London Children's Hospital on April 21. Two years before he had had an attack of measles, and for twelve months had been suffering from cough following an attack of pertussis. He had long ceased to whoop, but for a week before admission had been troubled with sickness after his fits of coughing. For four days he had seemed very ill, and his breathing had been difficult. When he came into the hospital the dyspnœa was so distressing that he could not lie down in bed. On examination the chest was found to be very much retracted at each base, showing that the child had been subject to attacks of bronchitis, which had produced some permanent collapse at the bases of his lungs. Superiorly, the chest was rather expanded, and the upper part of the sternum projected forwards. His breathing was very laborious, his

face livid, and his nares acted widely. At each inspiration there was much recession of the lower part of the chest. The boy sweated profusely, especially about the head, and the cough occurred in paroxysms like those of whooping-cough, during which his face became much congested. The paroxysm terminated in a slight whoop, but there was no vomiting. On examination of the chest, dulness was found at each base behind, rather pyramidal in form, and here the breathing was bronchial, with a large dry crepitation. Elsewhere about the chest the percussion-note was extra-resonant, and bronchitic râles were heard. On the evening of admission the boy's temperature was 103.4° . On the following evening, however, it was no more than 101.8° , and for a week the temperature alternated between 101° in the morning and 102° or higher at night. His pulse during this time was 130 to 150, and his respirations 40 to 60. During the first week the physical signs and the general symptoms underwent little alteration.

On April 28 the lividity was noticed to have disappeared, but the respiration remained very laborious, and at each breath there was much recession of the lower parts of the chest-walls. Still the child perspired much less profusely, and his condition was evidently improved. The dulness remained at each base behind, with slightly increased resonance of the voice, but with little augmentation of the vocal vibration. The crepitation had become more bubbling in character, and was heard over the lower half of each side posteriorly and to a less extent in front. 9 a.m.: Temperature 100.6° ; pulse 130; respirations 48. 6 p.m.: Temperature, 104.6° .

On May 5 the boy was looking very much better. His respirations had diminished in frequency, but still became very hurried after a fit of coughing. There was still dulness at each base behind, although less marked than before. A fine bubbling rhonchus was heard generally over each back, and this over the area of dulness was drier and more

crackling in character. Temperature : 9 a.m., 100° ; 6 p.m., $102\cdot4^{\circ}$.

On May 8 the physical signs remained much the same, but the general symptoms had greatly improved. The boy could lie down comfortably in bed ; his appetite was good ; he looked very much better, but still coughed a great deal in paroxysms like pertussis, and sometimes whooped faintly. His temperature still remained elevated. 9 a.m. : Temperature, $100\cdot8^{\circ}$; pulse, 118 ; respirations 48. On the previous evening the temperature had been $101\cdot6^{\circ}$.

By May 12 the cough had become much less frequent, although it still occurred in violent paroxysms. There was still some amount of dulness posteriorly, but the rhonchus had very much diminished, and was now limited to a crackling sound heard towards the end of inspiration, and to a more or less general sibilant wheeze. The temperature still continued higher than natural, being over 100° every evening, and rather lower in the morning ; pulse about 120 ; respirations about 40.

On May 18 the boy left the hospital. His temperature was still about the same—*i.e.*, a fraction of a degree over 100° both morning and evening, although sometimes for a few hours it would fall to 99° . His breathing was, however, slower (34), and his general health had very greatly improved. There was still deficient resonance at the left base behind. This was especially marked at one spot near the spinal column, where the respiration was very metallic and blowing, and the vocal resonance bronchophonic. No rhonchus was heard in this spot, but immediately around it a metallic crackling rhonchus occupied the end of inspiration. The percussion-note over the right back and the chest generally was hyper-resonant.

In this boy—an old bronchitic subject—the lungs generally were the seat of emphysema, but at the bases was a certain amount of permanent collapse, the result of frequent and severe attacks of pulmonary catarrh. This condition of the

lungs was partly the cause of the high rate of breathing which persisted after the symptoms generally had so much improved, for this lesion is always accompanied by more frequent respirations, and also by a more laboured manner of breathing than is natural to a healthy child. Catarrhal pneumonia, supervening upon such a condition of lung, is very apt to run a subacute course; and in this case the improvement in the symptoms was very gradual, and little alteration in the physical signs was noted from day to day. When the boy left the hospital there was still some unabsorbed consolidation at the left back, and the temperature had not returned to its natural level. While such symptoms persist there is always danger lest the case drift into one of chronic pulmonary phthisis. A deposit may, however, remain unabsorbed for a longer period, without danger, in the child than in the adult; but while it remains there is cause for constant anxiety lest any fresh inflammation be set up in the diseased part to lead to a further extension of the lesion.

It is especially in cases where the disease has assumed a subacute type that the bronchi are apt to dilate. In cases of this kind recovery, when it occurs, is slow, and if the child be not seen until re-absorption of the consolidating material has begun, some perplexity may arise as to the nature of the case.

CASE XXII.—ACUTE CATARRHAL PNEUMONIA WITH DILATATION OF BRONCHUS.

Rose H., aged six years, was admitted into the East London Children's Hospital on April 30. The child had been ill six weeks. The disease had begun suddenly with fever and a little cough, which, however, was not at first very distressing. After a few days it became more severe, and was accompanied by some pain in the side. Sometimes she expectorated with the cough; and she lost flesh steadily.

The girl had had whooping-cough some years previously, but no other illness except an attack similar to the present about twelve months before coming under notice.

On admission the child looked ill. She was thin: her eyes were sunken and her cheeks hollow (on May 4, she weighed 32 lbs. 10 oz.). Her temperature was 102.8° at 6 p.m. Examination of the front of the chest discovered nothing but a little fine crepitation at the right base. At the right base behind there was dulness on percussion, with weak tubular breathing and a high-pitched metallic large bubbling rhonchus. The vocal resonance was bronchophonic. The dulness did not extend into the infra-axillary region, and the breath-sound was there very weak, with a fine distant crepitating sound accompanying the inspiration.

On May 1 the temperature was 100.2° both morning and evening. Pulse, 104; respiration, 30. After the girl had been a few days in the hospital her general condition improved greatly, but the temperature remained high at night, being generally between 102° and 103° . At the right posterior base the physical signs did not improve; on the contrary, the respiration assumed a cavernous quality, and was accompanied by a gurgling rhonchus, while the resonance of the voice continued to be bronchophonic. By May 18, however, the cavernous quality of respiration was replaced by a diffused blowing sound much less metallic, and from this date the dulness gradually diminished, all the physical signs improved, and the temperature became more natural. On June 4 no dulness was left at the base of the right lung; the respiration was harsh without rhonchus, and the vocal resonance was normal. The girl gained persistently in weight during her residence in the hospital, and left on June 4 weighing 37 lbs. 5 oz. She was treated with quinine, followed by 15 drop doses of *Liq. ferri perchlor.*

In this case the actual physical signs presented by the child on her admission were those of a cavity occupying the base of the right lung at the posterior aspect. The

problem was to determine the nature of the cavity, and to estimate the chances of recovery. The history pointed to an acute illness of six weeks' standing which had set in while the child was in health. This was evidently pneumonic, for the dulness was limited to the base behind, and the tubular breathing and crepitating rhonchus were sufficiently characteristic of a solidification of the lung. Had the disease been the remains of a pleurisy the dulness would have been chiefly in the infra-axillary region; the respiration would not have been tubular at so late a period; and any crepitating sounds which might have been present would have been due to friction of the pleural surfaces, and would therefore have been larger and more superficial in character.

What was the nature of the cavity? It might have been due to breaking down of lung tissue, or to dilatation of a bronchus. The child was very thin; there was considerable pyrexia; and although the base is a less usual seat of ulcerative destruction than other parts of the lung, yet such a lesion was quite possible. There was, however, one serious objection to the ulcerative origin of the cavity: the disease in this case had only existed six weeks, and although caseous deposits may be formed, may soften, and may break down even in a less period of time, yet it is excessively unusual for a *solitary* pneumonic deposit in a child to undergo such rapid softening. In almost all cases of acute phthisis—at any rate in young subjects—the deposits are numerous and are scattered over the lung. For this reason the cavity was thought to be due to a dilated bronchus, and the favourable progress of the case confirmed the diagnosis.

Catarrhal pneumonia is a very common complication of tuberculous bronchitis. In such cases the disease runs a rapid course, and the symptoms and physical signs differ in no appreciable respect from those already described as characteristic of the most acute form of the disorder. The

presence of the grey granulation is therefore a matter of inference to be determined by the history of the case, and by the occurrence of new symptoms arising from other and distant organs which have become implicated in the general complaint. This subject will be discussed in a later chapter.

On account of the absence of physical signs in the early period of catarrhal pneumonia, we must be prepared to make our diagnosis of the disease at that stage from a study of the general and local symptoms alone, without aid from a physical examination. The history of the case is usually that a child who had been suffering for some days or weeks with a very bad cough had suddenly become very much worse, with high fever, rapid laborious breathing, and an almost constant hacking cough. We find the patient with a distressed expression and a pale, more or less livid face. His nares act widely; he breathes quickly, but with much effort; and there is great recession of the soft parts of the chest-walls during inspiration. The pulse is small; the superficial veins are prominent; and the pulse-respiration ratio assumes the type so characteristic of pneumonia. His temperature is found to be 102° . Examination of the chest gives little positive information; there is no dulness on percussion, and the stethoscope merely detects general moist râles of varying sizes over the chest and back. The symptoms are those of capillary bronchitis, with this difference—viz., that the temperature is higher than is found in an uncomplicated case of that disease; the respirations are more laborious and are rapid out of proportion to the pulse; and the cough is the cough of pneumonia.

These differences are of the utmost importance. If in a case of bronchitis the cough *suddenly* changes its character and gets short and hacking, while the pulse-respiration ratio becomes greatly perverted, and the temperature rises to 102° or higher, we may safely conclude that the original complaint is complicated by an attack of catarrhal pneumonia.

In cases where, in addition to the physical signs of bronchitis, some local dulness is discovered at any part of the chest-wall, with bronchial or blowing breathing, and a more or less dry crepitating rhonchus, this must be taken as additional evidence in favour of the pneumonic complication.

Lobular collapse of the lung is a disease which presents many points of resemblance with the one we are considering. Both are secondary to bronchitis. In both the lividity is great, the cough frequent, short, and hacking, the pulse small, and the breathing hurried. The natural relation between the pulse and the respiration is altered in both diseases—indeed, in lobular collapse this perversion often reaches its extremest degree—and in both a physical examination may reveal nothing but the signs of bronchitis. Two important points of distinction are, however, found in the temperature of the body and the character of the breathing. In catarrhal pneumonia the temperature is high and the breathing quick and laborious. In lobular collapse, on the contrary, the temperature is normal or even below the natural level, and although the respirations are rapid, the breath is drawn quietly and without effort.

When catarrhal pneumonia affects a considerable area of lung surface, so as to produce extensive consolidation, it presents a combination of the physical characteristics of bronchitis and pneumonia. The signs, however, differ in a very perceptible degree from those of croupous pneumonia even over the actual seat of solidification. As in every case of lung consolidation, there is dulness on percussion, with increased resonance of the cough, and perhaps a perceptible vocal vibration of the chest-wall (although this sign, as in the case of croupous pneumonia in children, is very untrustworthy), but the breathing is often merely bronchial, or, if blowing, usually wants the intense sniffing character so characteristic of true tubular respiration. The crepitation, again, is seldom so fine or dry as in croupous pneumonia ;

it occupies, also, a greater extent of the breath-sound, and is not limited to a fine puff at the end of inspiration. Often it is more a dry sub-crepitant rhonchus than a true crepitation. One of the principal points of difference, however, consists in the fact that while in croupous pneumonia the crepitation is limited to the confines of the consolidated portion of the lung, and is not heard where tubular breathing exists, in catarrhal pneumonia the crepitation becomes finer the nearer the stethoscope is moved to the centre of the consolidation, and is heard in the same spot with the most metallic blowing respiration.

The presence of the accompanying bronchitis is another feature which is characteristic of catarrhal, as distinguished from croupous pneumonia. In the latter the physical signs are limited to the part of the lung which is the actual seat of inflammation ; or if rhonchus be heard at any other part of the chest, it is merely of the sonoro-sibilant variety, and is trifling in amount. In the catarrhal form, on the contrary, dry and moist râles are heard all over both lungs.

In acute catarrhal pneumonia the two lungs are often attacked simultaneously, and a characteristic form for the consolidation to assume is that of a pyramidal strip on each side of the spine, reaching upwards from the base. Diffused collapse of the lung is very apt, also, to assume the same shape, and is also generally bilateral : it is, however, distinguished by the low temperature, and the absence of laborious breathing.

The prognosis in catarrhal pneumonia is very grave, and this variety of pulmonary inflammation is much more to be dreaded than the croupous form. Its greater severity may be partly accounted for by the fact that the latter disease is in young children nearly always a primary complaint, while in the case of the former the strength has been already reduced by the bronchitis which preceded it. The age of the child and the degree of vigour possessed by the patient at the time of the attack exercise a considerable influence upon

the ultimate issue of the illness ; for weakly children and infants, with their feeble resisting power, and tendency to rapid prostration of strength, have little prospect of passing safely through so formidable a complaint. The danger in ordinary cases is in proportion to the acuteness of the disease and the intensity of the general symptoms. When the inflammation runs a sub-acute course, as it is apt to do after whooping-cough, or in cases where it supervenes upon chronic bronchitis in an emphysematous subject, the chances in favour of ultimate recovery are very much increased.

In severe cases, symptoms of unfavourable augury are—lividity in the face and blueness of the lips ; a greatly perverted pulse-respiration ratio ; loss of brightness of the eye ; and marked apathy or somnolence. If a young child shows complete indifference to the discomforts of an examination, and allows himself to be disturbed, not only without complaint, but even without manifesting annoyance by any change of expression, the most serious apprehensions should be entertained as to the probable termination of his illness. On the other hand, if the lividity diminish, the breathing become easier, and the cough less constant and distressing, a more favourable view may be indulged in, even although the temperature show no sign of falling.

The treatment of catarrhal pneumonia differs little from that of the pulmonary catarrh by which it is preceded. “A cold on the chest” of a child should never be neglected, for while it lasts the child’s natural susceptibility to changes of temperature is greatly aggravated, and a fresh chill might set up a severe bronchitis, which would in all probability lead to an attack of catarrhal pneumonia.

In the early stage a pulmonary catarrh may be readily checked by confinement to bed, the repeated application of weak mustard poultices to the chest and back, and the administration of frequent small doses of ipecacuanha wine in a simple saline draught. If the bronchitis be a severe one, counter-irritation is very important, but strong applications

are not required. It is better to act slowly upon a large surface of the skin than rapidly upon a small surface. For an infant of twelve months old one part of flour of mustard may be added to six parts of linseed meal. The poultice should be made with warm, but not boiling, water, and should be large enough to cover the whole of either the chest or back. It may be kept in contact with the skin for six, seven, or eight hours, or as long as the patient is able to bear it. For older children a larger proportion of mustard may be used, but it should always be sufficiently diluted with linseed meal to allow of the application being kept upon the skin for several hours in succession without producing too severe an irritation.

In giving medicines at this time care should be taken to adapt the drug to the stage the catarrh has reached, for much mischief may be done by heedless dosing. The rule cannot be too rigidly followed, that in an early stage of pulmonary catarrh, while the bronchial mucous membrane is much swollen, and before secretion has become free, all stimulating expectorants are injurious. At this stage, ammonia, squill, tolu, and the other bronchial stimulants, instead of promoting secretion, only make the cough harder and the chest tighter, and aggravate the irritation which it is our object to allay. In an early stage of the disease, when the oppression of the chest is great and the dyspnoea troublesome, such treatment only increases the distress of the patient, and promotes the extension of the bronchitic inflammation to the air-cells. Many a case of catarrhal pneumonia has owed its origin to no other cause than injudicious treatment of a pulmonary catarrh. The greatest and most immediate benefit is obtained by medicines which excite the action of the skin and encourage free secretion from the bronchial tubes. If we can do this without at the same time depressing the patient's strength, we adopt the readiest means to relieve his suffering and hasten his ultimate recovery. For a child of one year old five drops each of

ipecacuanha and antimonial wines given every three hours in a simple saline draught will have the effect desired. If any depression seem to be induced by the mixture, a few drops of spirits of chloroform may be added to each dose, or a few drops of brandy may be given occasionally in addition. An emetic dose of ipecacuanha given early in the disease is useful in loosening the cough and relieving the tightness of the chest. This treatment, combined with confinement to one room, kept at an even temperature, and the application of weak mustard poultices, as before described, will soon produce an improvement in the symptoms ; afterwards, when secretion is free and the oppression has been removed, a mixture containing ammonia and squill, with small doses of pectoric, will prove of material assistance in completing the cure. On account of their tendency to collapse of the lung, very young infants suffering from pulmonary catarrh should be carefully watched, so that any unfavourable change in the symptoms may be immediately noted. Thus, if the breathing become shallow, and the face at all livid or dusky, they should be at once aroused if asleep, and forced to exert their inspiratory power. When an infant cries, his inspirations are much deeper than when he is quiet. A cold wet sponge may be suddenly applied to the naked chest, or vigorous friction may be used to the thorax. An emetic is also useful to clear away the phlegm from the tubes and promote deep inspiration.

Older children should be confined strictly to bed if the catarrh be at all severe ; for by this means not only do we protect the patient from a fresh chill, but we also insure a necessary amount of rest. A sick or feverish child requires *repose* ; his mind should be kept unexcited and his body quiet ; and to put him to bed is the most effectual way of accomplishing this object. Violent purgation is more injurious than beneficial in these cases, but care should be taken that the bowels are properly relieved. It is well to promote the flow of bile by an occasional dose of grey

powder. The diet should be simple, and may wisely be limited to milk and broth so long as the cough continues hard and the breathing much oppressed. In proportion, however, as the cough loosens, the diet may be improved, and stimulants may be given if required. Before this they should not be made use of unless the condition of the child imperatively calls for their administration.

When a case of bronchitis is complicated by the occurrence of catarrhal pneumonia, the treatment differs from that already described only by the more energetic use of counter-irritation. The poultices should be large enough to reach completely round the chest, and should be kept in contact with the body long enough to produce very deep redness of the skin. In bad cases, even if some slight blistering of the surface be produced, the effect will not be injurious.

The medicines at first should consist of the ipecacuanha and antimonial mixture, as recommended for bronchitis; but directly any signs of depression are noticed, this must be supplemented by a necessary quantity of stimulant, or, if thought desirable, must be omitted altogether and be replaced by ammonia and ether given with an alkali. Food must be administered in small quantities at frequent intervals, and should consist of milk with "Mellin's food" for infants not at the breast, and for older children of milk and broths. If necessary the milk may be guarded by a few drops of the saccharated solution of lime. As the strength fails it must be supported by strong beef-tea, brandy, and the brandy-and-egg mixture of the "Pharmacopœia."

Increased rapidity of respiration, combined with a fall in the pulse and a dusky tint of the face, is a sign that the right ventricle is becoming overtaxed. It is important, therefore, to afford some relief to the labouring heart or the child will die. This relief is best given by the abstraction of blood. In the case of an infant, three or four leeches

may be applied to the precordial region, and the number can be increased for an older child. In case of great urgency a little blood can be taken from the arm ; but it is wiser to adopt the milder course in all cases where it can be done without risk to the patient. Improvement is usually immediate. The breathing at once becomes slower and less shallow ; the pulse increases in frequency, and the dusky colour of the face gives place to a less ghastly tint.

It is necessary to keep a watchful eye upon the temperature. If the thermometer register 104° or 105° , it is advisable to reduce the bodily heat by the same measures which have been already recommended in the case of croupous pneumonia, taking the same precautions to obviate sudden collapse.

In cases of recovery care should be taken during convalescence not to be too anxious to return to the diet and habits of health. In particular, the patient should not be overloaded with large quantities of nourishing food, in the hope of more quickly restoring his exhausted powers. Recovery, when it happily occurs in catarrhal pneumonia, is always gradual ; and considering the severity of the disease, and the danger escaped from, we must be contented that it should be so.

CHAPTER VI.

CHRONIC CATARRHAL PNEUMONIA AND UNABSORBED PNEUMONIC DEPOSITS—PNEUMONIC PHTHISIS.

CHRONIC catarrhal pneumonia produces the common variety of consumptive disease—Pathology—Modes of formation of caseous masses—changes which they undergo—formation of cavity—Calcification—Fibroid formation—Secondary changes in other organs.

Symptoms and physical signs—Peculiarities of pulmonary disease in children.

Case XXIII.—Chronic catarrhal pneumonia—Rapid absorption of the deposit.

Physical signs of excavation—may be produced by dilated bronchi—Method of distinction.

Case XXIV.—Softening and breaking down of a caseous mass—Comment.

Case XXV.—Caseous consolidation with dilatation of bronchus resulting from an attack of acute catarrhal pneumonia—Comment.

Case XXVI.—Chronic catarrhal pneumonia, with secondary acute tuberculosis—Comment.

Diagnosis—Prognosis—Treatment.

THE principal form of chronic lung disease in the child arises, as it does in the adult, from the presence in the organ of cheesy deposits—the result of a modified process of fatty degeneration taking place in the products of inflammation. These cheesy deposits may be the remains of an ordinary croupous pneumonia, the consolidation having failed to be completely absorbed. This, however, is rare. It is more common to find an acute catarrhal pneumonia, leaving behind it an unabsorbed mass which undergoes the process of caseation, and leads to pulmonary phthisis. The most common form of consumptive disease, however, is the consequence of a chronic catarrhal process

—the catarrhal inflammation running a tedious course, stuffing the alveoli with newly-formed cells, plugging the smaller bronchi, and establishing more or less solid nodules in the tissue of the lung. These in most cases extend themselves by a repetition of the same process, and, by the changes which they undergo, lead to the most serious consequences.

In chronic catarrhal pneumonia, the mucous membrane secretes a peculiarly viscid and concentrated fluid, which is loaded with corpuscular elements. In the smaller air-tubes, the channel is often completely plugged by the quantity of secretion; the basement membrane disappears, and the corpuscles permeate the walls of the tubes. In the vesicles, the epithelial cells swell, become detached from the wall, and separate one from another. Their nuclei then divide, an active corpuscular proliferation is set up, and the alveoli become filled with large spheroidal cells, which contain large vesicular nuclei. These changes are accompanied by certain other alterations in the bronchi and pulmonary tissue. The peribronchial connective-tissue becomes increased, so that the walls of the finer air-tubes are thickened in places. The plugging of the smaller bronchi causes collapse of the air-vesicles to which they lead. As a consequence, there is a greater rush of air to the parts of the lung which still remain pervious, and this causes a cylindrical dilatation of many of the minuter bronchi, with marked stretching and thinning of their walls. The collapsed state of the vesicles induces congestion. This is soon followed by effusion of fluid into the air-cells, so that the part becomes œdematous. At this period the affected portions of lung are seen as dark reddish-brown patches, moist, smooth, and of doughy consistence. The consolidation is still capable of removal by absorption of the newly-formed cells, in which case nothing would be left but some thickening of the pulmonary tissue at the spot, with a little compensatory emphysema around it.

If absorption do not take place, the contents of the vesicles next undergo caseation. The newly-formed cells accumulate in the alveoli, and become closely packed together. They atrophy and undergo a partial fatty change, which, under the microscope, gives a cloudy appearance to the cell, and is indicated to the naked eye by a yellowish-white sharply-circumscribed opacity in the centre of the deposit. Within the area of the caseating mass the pulmonary tissue is destroyed, with the exception of the elastic fibres and coats of the larger arteries, which still remain entire.

At first the inflammation and its accompanying infiltration are limited to the circumference of the smaller bronchi; and on section of the lung we find a varying number of grey transparent nodules, whitish at the centre, varying in size from a poppy-grain to a millet-seed. Each of these is situated in the middle of a lobule. The intervening lung tissue may appear normal to the eye, but it is always more or less congested and collapsed, or is œdematous, or merely emphysematous and pale. Immediately surrounding the nodule is often a zone containing a reddish-grey glutinous infiltration. Gradually the grey nodules increase in size, and become distinctly cheesy; adjoining nodules unite; and finally the whole lobe may become infiltrated and caseous. Sometimes the inflammatory change spreads more abruptly from a large bronchus to the whole extent of lung which it supplies, so that the whole or greater part of a lobe may be at once invaded by the catarrhal and caseating process. This form is the old "tubercular infiltration" of Laennec.

It is then from a catarrhal process affecting the air-cells and finer tubes that pulmonary cheesy masses usually owe their origin. Sometimes, however, as has been said, they are formed in a different way, and a cheesy mass is left as a consequence of an attack of ordinary croupous pneumonia. In all cases of croupous inflammation of the lung, the stage

of yellow hepatisation is accompanied by a certain amount of catarrh of the alveolar walls. In fact, the principal cause of the change in colour of the hepatised part from dark red to reddish-yellow is stated by Rindfleisch to be the hindrance to the passage of blood by the infiltration of new cells in the interstices between the vessels. In ordinary cases this proliferation of epithelium is limited in amount, but sometimes the production of corpuscular elements is so great that the fibrinous pneumonic exudation becomes enclosed in a thick mass of new cells, and the whole undergoes caseation at the same time. By the post-mortem inspection of an affected lung, it is easy to discover what were the chances of a successful termination to the disease. Thus, if the part attacked by pneumonia be found to be sunken, dense, greyish, and slightly granular on section, allowing little fluid to be squeezed out on pressure, and showing under the microscope merely small dwindled cells without any excess of fat-globules, or signs of fatty degenerative metamorphosis, we may be sure that absorption was impossible, and that the chances of recovery in the case had been very small indeed.

After the process of caseation is completed, the cheesy mass may remain without alteration for a considerable time ; but, sooner or later, certain other changes take place in it. The period during which a cheesy deposit may remain indolent, varies in different cases ; but it may be stated generally that the larger the size of the mass, and the more quickly it has been formed, the more imminent is the risk of softening at its centre. Nodules slowly formed by the aggregation of smaller masses seldom soften before they reach the size of a hazel-nut ; while rapidly developed caseating deposits, even of single lobules, may undergo rapid softening. This is found in cases of "acute phthisis," where cheesy deposits of all sizes, and all undergoing a simultaneous softening, may be seen scattered over the lung. In children softening is less common than it is in

the adult, and cavities in the lung are more rarely seen. The softening process begins in the centre of the nodules, and is excited by the mere addition of fluid to the mass. The dead, dwindled cells are thus loosened, together with the molecular *débris* in which they lie. The softening is usually preceded by ulceration of the wall of the afferent bronchus—that, namely, which supplies the part, and lies in the centre of the nodule. The cheesy matter, when softened and converted into a pus-like pulp, is evacuated through an opening in the air-tube. Certain putrefactive changes then set in, and small fragments of cheesy matter separate from the newly formed cavity, and are expectorated. These, on examination, are found to contain elastic tissue and shrunk cells. Their appearance in the sputum is a sure sign that the cavity is continuing to extend itself. As a rule, when softening begins in a lung containing many cheesy deposits, more than one cavity is formed, and these often open into one another. Sometimes, after the cheesy matter has been removed, the cavity may contract by the formation of fibroid tissue in its wall, and become almost—or, if originally small, quite—closed, the sides cohering as a thick fibroid cicatrix. This is often combined with much proliferation of connective tissue in the lung, so that the part becomes indurated and its bronchi dilated.

Although the changes above described are those most commonly met with in cheesy deposits, yet the masses do not necessarily soften and break down. In exceptional cases a different transformation may take place. Thus, if the nodules are of small size, they may become calcified by the deposition within them of phosphate and carbonate of lime. Afterwards the petrified mass may become encapsuled, and remain embedded in the lung; or it may set up suppuration in the tissue around, separate from it, and be expelled through a perforation in a bronchus. Sometimes the whole caseous mass becomes converted into fibroid tissue, and then a condition is left such as is described in

the next chapter under the name of fibroid induration of lung.

The presence in the lung of the caseous products of catarrhal pneumonia is often the cause of secondary changes of a like character occurring in other organs. Thus, the disease is generally confined at first to one side of the chest, but when the softening process invades a caseous pulmonary mass, secondary deposits soon begin to be formed in the opposite lung which had been previously healthy. Again, ulceration of the lung is often found combined with the same lesion in the mucous membrane of the intestine. The parts affected are those in which ulceration also occurs in typhoid fever—viz., Peyer's patches and the solitary follicles in the neighbourhood of the ileo-cæcal valve. The follicles become enlarged by a proliferation within them of their corpuscular elements. A cheesy degeneration then takes place, and the caseous follicles soften from without inwards. In this way a sharply circumscribed ulcer is left, which is surrounded by a zone of connective tissue swollen by corpuscular infiltration. The ulcer extends its boundaries by a caseation and softening of the infiltrated tissues; and in this way neighbouring ulcers unite and form large oval lesions of surface, of which the long axis is always placed transversely across the direction of the bowel.

Another result of a softening cheesy mass is to set up a true secondary tuberculosis; and in this way the grey granulation may be found after death disseminated over the body, mixed up with the other pathological products proper to the disease.

So long as a small deposit remains completely inactive in the lung, it need give rise to no symptoms at all. If it be near the surface, a physical examination will detect its presence; but unless it causes pyrexia there is no interference with nutrition, and, if it excite no irritation in the tissue around, there is no cough. In the present chapter the symptoms produced by cheesy deposits and the changes

occurring in them will alone be discussed. Tuberculosis and the consequences arising from the formation of the grey granulation in the lung will be described later (*see* Chapter VIII.).

A caseous pneumonic deposit is the ordinary form of chronic pulmonary consolidation, and the changes which occur in it as the softening process invades the mass constitute pulmonary phthisis—pneumonic phthisis, as it has been called, from its origin—the common form of consumption of the lungs. Children in whom this form of pulmonary disease is found do not necessarily present in their physical conformation any special peculiarities of build. They are not all, or nearly all of them, narrow-shouldered, or pigeon-breasted, or shallow in the chest. On the contrary, in children as in adults, we constantly meet with an unabsorbed pneumonic deposit in cases where the thorax is perfectly formed, and where the general structure and appearance are suggestive of anything but “consumption.” Still it cannot be denied that a constitutional predisposition to catarrhal changes in the lungs, and to rapid caseation of pulmonary deposits, is often expressed in the physical conformation. In pronounced cases the lungs, as Sir William Jenner has pointed out, are singularly small, and this peculiarity influences the shape of the thorax which becomes elongated, with unusual obliquity of ribs and abnormal arching of the diaphragm.

Much has been written about the peculiarities of chronic pulmonary disease in children, and the difficulties of its diagnosis, but these difficulties have been greatly overrated. It is true that the thorax is of small size, and that sounds from the throat are readily transmitted to the chest, but if the child’s mouth be open during the examination, the transmission of sounds is in a great measure prevented. It is also true that little trust is to be placed in the amount of vibration of the chest-wall, for in young children vocal fremitus is always weak, and is generally absent altogether.

But these are comparatively slight disadvantages. On the other hand, the resonance of the voice furnishes as important evidence in the child as it does in the adult. Most children will say "No" when desired. It is a word they understand, a sonorous word, and one which can be pronounced without trouble. From it we can form as sufficient an idea of the state of vocal resonance as we can possibly desire. Again, by gentle percussion with two fingers, the most minute variations in the sonoriety of the chest can be detected without difficulty. The great secret of examining the chest in children is to have the patient raised up to a convenient height. To bend down to a child as he sits on his mother's lap, or in a low cot, is to examine him under very disadvantageous conditions. The child should be always seated upon a table or high stool, so as to bring his body nearer to the ear of the observer.

There are two sources of error which it is important to avoid. In the first place, it is necessary to remember that the breath-sound in a child is naturally loud and coarse (puerile respiration), and that therefore mere loudness and harshness are not the results of disease. In a child, elevation of pitch in the breath-sound is the most valuable indication of alteration from a healthy state. The change is first noticed in the expiration, but it very soon spreads to the inspiration as well, and we then have what is called bronchial breathing—that is to say, both inspiration and expiration lose their breezy quality, and become harder, drier, and higher pitched, and the expiration gets louder and more prolonged. In fact, in young subjects, bronchial breathing seems in many cases to be the first alteration from a healthy state of breath-sound. Mere prolonged breathing is very common in children at all parts of the chest, especially at the apices; and it is of importance to be aware (as a second source of error) that enlarged bronchial glands assist greatly in conducting sounds from the throat and trachea; so that in cases where the enlargement is consider-

able, blowing breathing with very long expiration may be heard at the apices of the lungs, although the lungs are themselves quite healthy. In children who suffer from enlarged tonsils, and even in children in whom no morbid condition of the fauces can be discovered, the same peculiarity is often noticed, especially when the lips are closed. On this account it is important to have the child's mouth open during examination of the chest. There is often great irregularity in the breathing of a child. The respirations may vary greatly in depth from time to time; and even in comparing the two sides, the intensity of the breath-sound is often found to be greater on one side than it is on the other. On this account, mere weak breathing in a child, however local it may appear to be, does not necessarily indicate disease of the lung. Sometimes, however, the beginning of pleurisy is marked by weak breathing at one base, but even in this case the respiration is usually also high-pitched or otherwise altered from a state of health. When the breathing becomes bronchial as well as weak, it must be looked upon as a sign of disease.

The symptoms and signs of chronic catarrhal pneumonia are those of a slowly growing consolidation of the lung, which gives rise to pyrexia, and, by the consequent interference with nutrition, occasions some loss of flesh, and eventually, perhaps extreme emaciation. Until, however, the disease is far advanced, the general symptoms are comparatively mild unless there be much pyrexia, and are far overbalanced by the local symptoms and by the intensity of the physical signs. A child may have a cough for a long time without appearing to suffer in any other way; but if, in addition to the cough, we find elevation of temperature at night, followed by slight loss of flesh, we have great reason to fear that the catarrhal condition of the mucous membrane of the finer tubes has spread to the alveoli, and that destructive changes have begun. When thus chronic from the first, the seat of the pneumonia is the same in the child as it is in

the adult—viz., almost invariably at the apex of one lung. In a very short time after the symptoms have been noticed, we find the percussion-note at the apex on one side to be high-pitched: the breathing becomes faintly bronchial, and a click or two is heard at the end of a deep inspiration. The alteration in the percussion-note is usually observed first at the supra-spinous fossa. There is, however, here a source of fallacy which it is well to be aware of. On account of the inequality of breathing in early life, and the deficiency of expansion of the upper part of the chest, young children are liable to slight collapse of the apices of the lungs. Such collapses are usually seated at the posterior aspect of the lung, and give rise to slight dulness at the supra-spinous fossa, with weak bronchial breathing, and often a little cre-pitation at the end of inspiration. If, therefore, the temperature at the time of examination be not elevated, we should hesitate about inferring, from such signs alone, the existence of a caseous deposit, but should wait to correct our first impressions by the results of a second investigation. In the case of collapse the physical signs seldom persist longer than two or three days, and if the child be old enough to regulate his breathing as required, a few deep inspirations will usually cause the dulness to disappear at once.

The caseous deposit is at first limited strictly to one apex, the other being perfectly healthy. The temperature at this time rises to about 101° in the evening, and generally falls again to below 100° in the morning. The appetite is often impaired, and the bowels are very apt to be relaxed, for children suffering from catarrh in any form are easily affected by the slightest chill, and therefore the sensitive intestinal mucous membrane is very frequently involved in the derangement. The presence of the caseous deposit in the lung seems to keep up an irritable condition of the whole mucous membrane, so that bronchitic râles of the dry variety are very frequently to be heard, and these often persist for a long

time, although the general symptoms are undergoing progressive improvement.

CASE XXIII.—CHRONIC CATARRHAL PNEUMONIA.
RAPID ABSORPTION OF THE DEPOSIT.

William T., aged nine years, was born of a consumptive family on the father's side; the mother's family was healthy. The boy had had whooping-cough and measles, but recovered completely, and remained in good health until Christmas, when he began to cough, especially at night. Towards the end of January, after a severe wetting, the cough got worse, and he was thirsty and feverish; his appetite, too, began to fail, and he complained of pains in the chest and limbs. From that time he seemed to grow daily worse, until February 19, when he was admitted into the East London Children's Hospital.

On admission the boy was pale and rather thin. His sternum was a little prominent, and there was slight flattening in each infra-mammary region. The two sides expanded pretty equally. The heart's impulse was in natural site. Some want of resonance was found at the right apex, both before and behind, and the respiration was high-pitched and rather bronchial (with the mouth open). Sonoro-sibilant rhonchi were heard not only here but all over the right side, both at the back and the front, and also—although to a less extent—on the left side. With this exception, the physical signs generally were natural. Heart's sounds were healthy. The boy's temperature was always over 100° at night, and it sometimes rose to 101°. In the morning, after the first three days, it generally fell to 99°. During the first week the pulse varied between 102 and 108, but afterwards it became slower. The respirations were 28 to 30.

After the boy had been in the hospital a few days he began to perspire copiously at night, but the sweating was

greatly diminished by half-drachm doses of tinct. belladonnæ given at bed-time. The physical signs gradually improved, and on March 18 the right apex was resonant, except at the supra-spinous fossa, where the percussion-note was slightly high-pitched. Respiration, however, was natural. The dry râles still persisted over both lungs. The patient gained in weight. On March 3 he weighed 3 st. 11 $\frac{1}{4}$ lbs., and on March 18, 4 st. 1 lb. After March 3 the temperature was always under 100° at night, and the respirations never rose higher than 26 in the minute. The boy left the hospital on March 28.

The case was one of chronic catarrhal pneumonia affecting the apex of the right lung. After the boy's admission into the hospital, absorption very quickly took place, and on his discharge there was left merely a little thickening of the tissue of the lung at the seat of disease. In this case, on account of the rapidity of its removal, it is unlikely that caseation of the deposit had taken place; for although it is probable that a cheesy mass may undergo absorption under favourable conditions (especially in a child), yet the process of removal must necessarily be slow.

When the catarrhal pneumonia is chronic from the first, the resulting consolidation is, as has been said, almost invariably seated at the apex of the lung. In other cases, however, where the solidification is the consequence of an acute inflammatory attack, whether of the croupous or catarrhal variety, other portions of the lung are frequently affected, and the base is very often the seat of disease. In this way any part of the lung may come to contain an unabsorbed deposit, which may set up phthisical changes, and lead to extensive disorganisation.

When softening begins in a pulmonary cheesy mass, the change is indicated by the moister character of the crepitating or crackling sounds which accompany the respiration. Very soon the rhonchus assumes a metallic quality which is very noticeable, for pulmonary softening, when it has once

begun in a child, usually proceeds with rapidity. The breath-sounds quickly become blowing and metallic, or even cavernous, while the resonance of the voice and cry is intensified and becomes broncophonic. If the case have been from the first under observation, and the changes in the physical signs have been noted from day to day, the occurrence of signs such as the above, coming on gradually, and not succeeding to an acute inflammatory attack, indicates the progress of softening and ulcerative destruction of pulmonary tissue ; but unless the successive steps of the process have been followed, as in the case supposed, the presence of dulness with cavernous breathing and gurgling rhonchus does not *necessarily* imply so serious a lesion of the lung. Acute and subacute catarrhal pneumonia frequently lead to considerable dilatation of bronchi ; and a dilated bronchus, surrounded by solidified tissue, will give rise to exactly such physical signs as have been described. It is important in our diagnosis to distinguish clearly between these two conditions, for the ultimate issue of the case is likely to be very different, according as to whether actual breaking-down of the lung be or be not in progress. In children the bronchi dilate with very great ease ; and as in such subjects fibroid induration is readily set up, the dilatation is often a permanent one. Cases of prolonged phthisis with cavernous signs, where the child falls away during the winter, but begins to improve again and gain flesh on the return of more genial weather, are usually cases of this description. Such children often live for many years, dying eventually of pneumonia or other acute pulmonary attack.

The seat of the physical signs is an important element in the diagnosis. Thus, when occurring as a consequence of catarrhal pneumonia* dilated bronchi are much more

* The statement in the text refers merely to cases of catarrhal pneumonia, and not to dilatations of the bronchi from other causes, such as fibroid induration of the lung.

often found towards the base of the lung than elsewhere, while ulcerating cavities are more frequently situated near the apex. Again, the general symptoms afford very valuable information. If the appetite is good, and the child is gaining flesh and strength, while the temperature is normal or only slightly elevated, we have good grounds for inferring that the cavernous signs are not due to ulcerative destruction of the lung. Besides, the tendency of ulcerating cavities is to extend themselves, that of dilated bronchi to contract. If, therefore, we find, as the days pass, that the cavernous sounds become less and less intense, and less widely audible, we may conclude that it is with the less serious lesion that we have to deal. The most conclusive evidence, however, is furnished by a careful microscopic examination of the expectoration. If this be found to contain areolar fragments of elastic tissue, the existence of ulcerative excavation of the lung is at once established. Unfortunately, in children, who generally swallow all sputa, this evidence is not often within our reach.

Cavities are not always indicated by such positive physical signs as have been described above. If they are seated near the surface of the lung, they probably always give rise to cavernous respiration, metallic bubbling rhonchus, and bronchophony; but in cases where they are more deeply seated, and are separated from the surface by tissue not completely consolidated, the evidences of their presence become much less distinct. Indeed, it is questionable whether a small cavity, seated in the middle of the lung, and not communicating with a bronchus of some size, can be discovered by any means of physical investigation at our command. It is certainly the case that a cavity may be present without giving rise to other signs than slight dulness, faintly bronchial breathing, a small bubbling rhonchus, and increased intensity of vocal resonance,—signs which are distinctive of nothing more than mere consolidation of the lung.

Again, cavernous breathing of the most intense kind may be present, although there is no cavity. This phenomenon is sometimes to be found in cases of pleuritic effusion in the child. The diagnosis is made by noting the existence of the other signs of pleurisy. Where the lung is really the seat of excavation, effusion is rarely seen, for although pleurisy is common in such cases, it is almost always of the plastic variety, in which little or no fluid is poured out. Enlarged bronchial glands, by their conducting power, will give rise to loud blowing breathing at the apices of the lungs, and this may even acquire something of a cavernous quality. In such cases, however, the rhonchus, if present, is not metallic, and when the mouth is open the cavernous character of the respiration is much diminished. It must not be forgotten that enlarged bronchial glands may co-exist with cavities at the apices of the lungs.

The process of softening in a cheesy mass is not *necessarily* accompanied by any special severity in the general symptoms. The cough is often not much more frequent; the appetite may remain good; and even the weight of the body may be but little diminished. Usually, however, the process is accompanied by a very evident impairment of the general health. The beginning of softening is often indicated by high fever, flushings of the face, and morning perspirations; and in most cases the appetite fails, the temperature rises, the languor increases, the patient sweats profusely at night, and loses flesh unmistakably. Often at this time diarrhoea is set up, and the rapidity of the wasting is increased. Children seldom expectorate unless the cough causes vomiting; as a rule they swallow the sputum as it reaches the mouth. For the same reason hæmoptysis is rare* in the child, the blood being swallowed instead of

* It is, however, far from uncommon for children to eject blood from the mouth, but this is seldom the result of hæmoptysis. Children are very subject to attacks of epistaxis, and, according to the seat of the hæmorrhage in the nasal cavity, the blood flows forward through the

expectorated. The digestion is deranged in most cases. There is usually considerable irritability of the stomach, and vomiting is easily excited.

The following case furnishes a good illustration of the symptoms and physical signs attending the breaking-down of a cheesy mass :—

CASE XXIV.—CHRONIC CATARRHAL PNEUMONIA—
SOFTENING AND EXCAVATION.

William P., aged two years, was attacked by measles on New Year's Day. About a week after the disappearance of the rash, his cough was noticed to be very violent ; this was especially the case at night, when he would become feverish and get flushed in the face. After coughing he sometimes retched, but did not vomit. His appetite remained fairly good, but he was noticed to be losing flesh. On February 14 the child was admitted into the East London Children's Hospital.

On admission, the right side was found to be completely dull in front from the third rib to the base, but the dulness was less marked outside the nipple-line than between that and the sternum. Above, the natural resonance was rather diminished as far as the apex. The respiration just below the clavicle was high-pitched and slightly bronchial, but over the area of dulness it became markedly bronchial and rather metallic in quality, although not actually blowing. No crepitation or rhonchus was heard during quiet breathing, but a double metallic click was caught at the end of a deep inspiration. Elsewhere on the right side there was

anterior nares, or backwards into the pharynx. If the latter, coughing is often produced, and large quantities of dark-coloured blood are ejected, to the great alarm of the parents. In such cases we generally find, on inquiry, that the nose has been bleeding, and blood may often be seen trickling down the back of the throat from the posterior nares. Sometimes the blood is swallowed, and is afterwards vomited.

little remarkable about the physical signs: at the back the percussion-note seemed generally high-pitched, but the respiration was natural and there was no rhonchus. The left side was healthy. The temperature after the first evening was always under 100° , usually only 99° ; pulse 108 to 126; respirations 30 to 54.

The boy went out on February 21, with no alteration in his physical signs, but he was re-admitted on July 11 with severe diarrhoea. At this time an examination of the chest showed dulness at the right apex, both before and behind, with increased resistance. At the right back, from the spine of the scapula downwards, the note was rather tympanic; respiration at the upper part was bronchial with moist clicking rhonchi. Over the rest of the right side there was some clicking and wheezing rhonchus, both in front and at the side. At the left back the percussion-note was generally high-pitched, but the respiration was natural, and no rhonchus was noted except at the base, where occasional bubbles were heard. At the left front the physical signs were natural. The child was now thinner and paler, and looked much worse than before. His temperature was always over 100° at night, often rising as high as 102° , and on one occasion to 103.8° ; respirations 42 to 60; pulse 102 to 138. The cough was not violent, and was loose. The diarrhoea soon ceased under the influence of lead and opium, and the omission of milk from the diet. The cough, however, continued, and the child daily lost in flesh and strength.

On July 29 the dulness at the right apex was increased, and the breathing had become blowing in quality, with a large, almost gurgling, rhonchus. This rhonchus was heard all over the right side, both before and behind. At the right supra-spinous fossa the respiration was cavernous. All over the left side a coarse, crepitating rhonchus was heard, which disappeared to a considerable extent after a cough.

On the evening of July 29 the diarrhoea returned with some severity. Temperature, 7 p.m., 101.8° ; pulse 138;

inspirations 48. On the following morning the temperature was 98°, and the child died in the course of the afternoon.

On examination of the body, the right lung was found to be much diseased. The upper lobe, adherent in places to the chest wall, was solid and of a dark purple colour. Its substance was dense, and could not be broken down by the finger. The denseness was due to a large excess of fibroid tissue, for the divisions between the lobules were seen as greyish bands, and the walls of the bronchioles were thickened, although their channels were not dilated. Small cheesy masses were scattered through the substance of the lobe, and at the upper part was a cavity, the size of a walnut, whose outer wall was formed merely by thickened and greyish-coloured pleura. At this point the lung had been adherent to the wall of the chest. The middle lobe was healthy. The lower lobe crepitated imperfectly, and was red in colour. At the anterior border of the base there was a patch of consolidation, with two small cavities containing softened cheesy matter. The left lung was the seat of recent catarrhal pneumonia; old patches of cheesy matter were dispersed through the substance; no grey granulations were anywhere seen. The bronchial and mesenteric glands were enlarged, and the intestines—both large and small—were studded with transverse ulcers.

In this case the child had an attack of acute catarrhal pneumonia secondary to measles. As so often happens under these circumstances, absorption was carried on very slowly, and on the admission of the child into the hospital, between four and five weeks after the beginning of the inflammatory attack, there was still considerable consolidation of the right lung. This consolidation was seated more at the base than elsewhere, but the incomplete resonance at the apex, together with the high-pitched and rather bronchial breathing heard below the clavicle, indicated that the upper part of the lung was not altogether untouched by the disease. The history of the case—for, as far as

could be discovered, the child had been healthy before the attack of measles—showed clearly the nature of the illness. Catarrhal pneumonia is a frequent sequel of measles; and the presence of acute tuberculosis, the other common consequence of rubeola, was excluded by the strict limitation of the pulmonary physical signs to one side of the chest, for catarrhal pneumonia occurring secondarily to acute tuberculosis would certainly give rise to symptoms in both lungs. The temperature at this time was not high, being usually below 100° in the evening, and under favourable conditions the case might not improbably have done well. The child's friends, however, removed him from the hospital, and on his readmission, five months afterwards, the disease was found to have made great progress. Absorption of the deposit had taken place to a certain extent towards the base of the lung, but at the upper part chronic catarrhal changes had evidently been going on, so that there were distinct evidences of consolidation and softening at the right apex. The temperature was now high, and there was diarrhoea, which, by its depressing influence upon general nutrition, still further diminished any chances of recovery that might have remained. The case now went on rapidly from bad to worse. In a few days signs of excavation became apparent: the diarrhoea, which had been arrested, returned; and the child sank and died.

It would have been difficult in this case to exclude secondary tuberculosis at the time of the child's readmission to the hospital in July. The character of the physical signs made it evident that the opposite lung had become affected, and there was nothing in the case at this period incompatible with the presence of the grey granulation. Still, the formation of secondary deposits in different parts of the body, and of ulcerations in the bowels, may occur in such cases quite independently of tuberculosis, and there was therefore no need to resort to this explanation of the general symptoms. It is a point worthy of notice in passing, that a diarrhoea

resulting from so serious a lesion of the bowels could be arrested for a time by treatment, although there is no reason to suppose that any attempt at repair had taken place in the intestinal ulcers.

In the following case the disease was the direct result of an acute attack of catarrhal pneumonia, the deposit failing to be completely absorbed. The cavernous signs were not, however, produced, as in the preceding instance, by ulcerative destruction of lung, but were the consequences of a much less serious lesion. The case furnishes a good illustration of a condition which is far from being uncommon, and is one in which accuracy of diagnosis is of the utmost importance in estimating the probabilities of a favourable termination to the illness.

CASE XXV.—CASEOUS CONSOLIDATION WITH DILATATION OF BRONCHUS.

William C., a thin, pale boy, aged fourteen years, but looking very much younger, was admitted into the East London Children's Hospital on November 20.

On stripping the boy for examination, the sternum was seen to be prominent, especially at the lower part, and the chest was contracted laterally at the base, showing the existence of a certain amount of permanent collapse of the bases of the lungs from successive catarrhs. The respiration was rather laborious, but the two sides of the chest moved equally, and the intercostal spaces of each side sank in. The percussion-note was high-pitched at the right supra-spinous fossa, but the breath-sounds there and all over both sides were quite concealed by a profuse crepitating rhonchus, which on the right side was hard, metallic, and ringing; and this quality became more marked as the stethoscope was moved towards the base. The resonance of the voice was natural. In front there was no dulness on either side, except a slight want of resonance above the right clavicle;

but a bubbling rhonchus was heard more or less copiously on both sides. The vocal fremitus was everywhere strong; temperature, 104.8° . On the following morning—temperature 101.8° ; pulse 120; respirations 46.

On November 24, in addition to the want of resonance at the right supra-spinous fossa, there was slight dulness at the right posterior base, and here the respiration was tubular. The same crepitating rhonchus was heard all over both sides posteriorly, changing at the right base to loud metallic gurgling. Over the upper part of the right back the respiration was bronchial. Temperature (8 a.m.) 103.6° ; pulse 120; respirations 48; temperature (9 p.m.) 102.6° .

After this report the dulness at the left posterior base still further increased. On December 1 the want of resonance extended upwards as high as the spine of the scapula. Over the dull area the respiration was cavernous, with fine metallic bubbling rhonchus and bronchophonic resonance of the voice. The signs of general bronchitis persisted, for there was still much fine subcrepitant rhonchus at the left back. The signs of consolidation of the right apex remained unaltered. By this date the fever had become much reduced, and the temperature was only 99.6° both morning and evening; pulse 120; respirations 36.

The temperature remained between 99° and 100° until the evening of December 9, when it rose suddenly to 104.2° , and on the following evening was 105.2° . After this date, however, it again sank, but very gradually, and on the evening of December 21 was 98.8° . The elevation of temperature, which was no doubt due to a return of the catarrhal inflammation, was accompanied by an increase in the amount of subcrepitant rhonchus, but with no extension of the dulness or other alteration in the physical signs. There was still the same evidence of slight consolidation at the right apex, and at the base the dulness still reached as high as the angle of the scapula. The respiration was here cavernous as before, with metallic bubbling and intensely bronchophonic

resonance of the voice. The dulness did not extend into the infra-axillary region. All over the right front a crepitating rhonchus could be heard.

The boy's general health began now greatly to improve. His appetite became very good, and he grew rapidly fatter. The temperature remained little higher than normal, rising sometimes to 100° , but, as a rule, maintaining a regular level of a little over 99° . His cough was loose, and not distressing. Towards the end of December he began to perspire profusely at night, but the sweats were controlled by a draught containing tincture of belladonna given at bedtime.

Little alteration was noticed from day to day in the physical signs. The rhonchus varied somewhat in quantity, and the respiration at the right base behind was sometimes more markedly cavernous than at others, owing, no doubt, to the varying quantities of secretion in the lung at that spot. Alterations in the vocal fremitus and in the resonance of the voice seemed to be dependent upon the same conditions. Towards the end of January the dulness at the right back began to diminish in extent. Instead of reaching upwards to the angle of the scapula, it became limited to the extreme base. Thus, on February 12 the physical signs were the following:—Percussion-note still high-pitched at right supra-spinous fossa, and dull at extreme base; but the dulness is, as before, limited to the back. All over the right back a loud crepitating rhonchus is heard, which at the base becomes larger, harder, more ringing, and more superficial. Respiration at the base is cavernous, but not very loud, and at the supra-spinous fossa is bronchial. Ordinary bronchitic râles are heard at the left back and in front on the right side. Vocal fremitus to-day rather diminished at right posterior base. 8 a.m.: Temperature 98.2° . 6 p.m.: Temperature 99.2° ; pulse 96; respirations 22.

The boy left the hospital on March 8, looking fat and well. The high-pitched percussion-note still persisted at the

right supra-spinous fossa, and there was still dulness at the right base behind. The auscultatory signs were exactly the same as at the last report. Temperature (7 p.m. March 7), 99.6°.

On April 30 the boy came to the hospital for examination. The physical signs were now very much the same as before. Both sides expanded fairly well and equally. There was still evidence of consolidation at the right apex, and at the base the dulness persisted, and the respiration was cavernous, with a sharp crepitating rhonchus not removed by coughing. Vocal resonance at base bronchophonic. The crepitation was heard more faintly in the infra-axillary region, but the respiration there was excessively weak. Very little rhonchus was heard elsewhere, and the left side appeared to be quite healthy. The boy looked very well, and was fat. He coughed occasionally, but expectorated nothing. The finger ends were not clubbed.

This patient on his admission was evidently suffering from a chronic consolidation at the apex of the right lung, with secondary acute catarrhal pneumonia. There was slight dulness at the right supra-spinous fossa; but the breath-sound was not tubular, as it would have been in a boy of his age if the consolidation had been recent. The high temperature showed that the crepitating rhonchi heard about the back were not due merely to bronchitis, and although there was no loss of resonance over the back generally, yet the metallic quality of the râles on the right side—a quality which became still more marked towards the base—indicated the presence of deposits in the lung. The pneumonic process continued, and a few days afterwards evident dulness could be detected at the right posterior base, with all the signs of pneumonic consolidation. Cavernous signs soon began to manifest themselves, and in a short time it became evident, from the loud cavernous respiration, the metallic gurgling, and the bronchophony, that the base of the right lung was the seat of a cavity of some size. It was

necessary to determine the nature of this cavity, for the prognosis depended greatly upon an accurate estimate of the pathological condition of the lung. The temperature and general symptoms here furnished important information. At the very time when the cavernous signs became fully developed the temperature fell considerably, and at the same time the respirations, which had been 48 in the minute, sank to 36. A change so favourable was sufficient to exclude softening and breaking down of a recent deposit, for such rapid destruction of lung-tissue, instead of an amelioration, would have caused a serious increase in the severity of the general symptoms. Besides, the cavernous signs were seated at the base of the lung—a comparatively unusual situation for a cavity, although common for a dilated bronchus; and the signs had supervened upon an attack of catarrhal pneumonia where the development of a bronchiectasis is so often met with. These reasons were sufficient for concluding that the signs were due to a cavity formed by the dilatation of a bronchus. The after-progress of the case confirmed the conclusion arrived at; the condition of the boy improved greatly, the appetite returned, he gained flesh fast, and rapidly resumed the appearance of health, although the physical signs suffered little alteration, and some considerable consolidation, with all the signs of a cavity, remained as a persistent condition.

It is remarkable in this case how little change was induced in the chronic consolidation of the right apex by this attack of catarrhal pneumonia. A certain amount of bronchitic rhonchus was heard there as in other parts of the chest, but no extension of the consolidation took place at this spot, and no active change in it was induced by the presence of the catarrhal condition. Such a consolidation forms a good illustration of the inactive cheesy deposit which reveals itself by physical signs, but does not affect the general condition of the patient, and gives rise to no constitutional symptoms.

A caseous deposit may remain a long time unabsorbed, and yet may finally disappear; indeed, in children it is difficult to fix a limit beyond which its removal by natural processes can no longer be looked for. In the adult a deposit rarely becomes absorbed after the expiration of three months, but in young subjects this term must be greatly extended, and, provided the deposit remain completely inactive, a much longer time must elapse before absorption can be pronounced to be impossible. The occurrence of pleurisy with effusion on the diseased side appears to favour a healthy removal of caseous consolidations; and a case has already been related in a previous chapter where the physical signs of consolidation of some standing completely disappeared on the absorption of the effused fluid (see case of Emma W., p. 85). This, however, is only true of cases where the deposit is perfectly inert, without any active change taking place in it. If the mass excite irritation in the tissue around, or if softening have begun to invade the deposit, the occurrence of pleurisy appears to exercise a very injurious influence upon the disease.

The presence of caseous masses in the lung often leads to the formation of the true grey granulation in the body, and cases frequently terminate in an outbreak of acute general tuberculosis.

CASE XXVI.—CHRONIC CATARRHAL PNEUMONIA WITH PLASTIC PLEURISY AND SECONDARY ACUTE TUBERCULOSIS.

Edwin S., aged two years, was admitted into the East London Children's Hospital on March 8. He had been ill for eight weeks. The illness began with an eruption like urticaria and he was feverish at night. The next day he began to cough and his bowels were relaxed. He was seen by a doctor, who said the boy had inflammation of the lung. Since then he had been poorly, very restless, and

had slept badly at night. Breathing sometimes laboured : not much cough. Motions have sometimes contained thread-worms.

State on Admission.—A small child, pale and thin ; features large and coarse ; complexion thick ; eyelashes long ; eyebrows well marked ; a little eczema of nostrils ; no enlarged cervical glands ; and no appearance of hair on shoulders or along spine. Hands rather cold, and nails livid. Finger-ends slightly clubbed. Lips a little dry. A little dulness at right supra-spinous fossa ; respiration there bronchial, with a little crepitating rhonchus heard during inspiration. Bowels slightly relaxed, and motions offensive. Temperature at night 101·4°.

The dulness at the right apex gradually increased after the child's admission ; the respiration became loud and blowing, and the physical signs were noticed in the front as well as at the back. On March 19 some clicking sounds were heard at the end of inspiration, from the clavicle as far down as the nipple, and also in the axilla. By the 23rd the percussion-note had become high-pitched over all this area. During the next few days the resonance still further diminished, and on March 30 the percussion-note was completely dull from apex to base at the right front, and the respiration was cavernous, without rhonchus. Over the rest of the right side the natural resonance was impaired ; the breath-sound was loud and cavernous in the axilla, and bronchial behind. At the posterior base an occasional click was caught at the end of inspiration. The pulse-respiration ratio was little altered at this time. On March 18, pulse 130, respirations 30 ; on March 23, pulse 104, respirations 32 ; on March 29, pulse 120, respirations 32 ; and on April 1, pulse 128, respirations 36. The child could not be persuaded to speak, and therefore the state of vocal resonance and the presence or absence of vibration of the chest-wall could not be ascertained. Heart's apex was immediately below and to the left of the left nipple. On

the left side the physical signs seemed everywhere perfectly natural.

The child was evidently getting weaker and thinner. On March 15 he weighed 17 lbs., and on March 30 16 lbs. 1 oz. His face was colourless, with an expression of distress. The bowels were occasionally relaxed, and his appetite was poor. The temperature was rarely under 100° , and in the evening usually rose to 101° or higher.

On March 29, temperature at 8 a.m. 100.2° , 7 p.m. 101.8° ; pulse 120; respirations 32.

After this report the physical signs continued much the same. The dulness remained limited to the front of the chest on the right side, and the resonance laterally and at the back was little diminished. Posteriorly, the only marked dulness was at the supra-spinous fossa. Everywhere over the upper half of the chest the breath-sounds were loud and cavernous, without rhonchus even after a cough.

On April 2 the chest was punctured with the hypodermic syringe, but no fluid escaped.

On April 6 the signs were unaltered, except that the percussion-note in front had become rather tubular in quality, and a fine crepitation was noticed at the end of a deep breath immediately below the clavicle, and also at other parts of the chest. At the right supra-spinous fossa some large metallic bubbling was heard, and the respiration there and all over the upper half of the right side was loud and cavernous. Nothing abnormal was discovered on the left side of the chest. The temperature remained high, as before.

After this date the only alteration in the physical signs was a greater intensity in the cavernous respiration. The crepitation varied in quantity, and was sometimes absent altogether; when heard it was always fine. There was no approach to metallic gurgling rhonchus, and the only large bubbling heard was at the right supra-spinous fossa. The

left side of the chest never at any time gave evidence of disease.

The child did not waste persistently. On April 21 his weight was 17 lbs. 13 oz., being a gain of nearly 2 lbs. during the previous three weeks; on April 29 he weighed 18 lbs.; and on May 6 18 lbs. 6 oz. On this day, however, a severe attack of diarrhoea came on; rapid wasting then set in, and the boy died on May 12.

On examination of the body, the right lung was found attached by easily separable recent adhesions to the chest-wall. The two upper lobes were solidified; the lower, although firmer than natural, still crepitated to a certain extent. The surface of the whole lung beneath the pleura was dotted over with minute pearly specks, evidently grey granulations. On section, the upper lobe was found to be infiltrated with softening cheesy matter, and contained several small cavities, of which one at the apex was as large as a nutmeg. The middle lobe consisted of small yellow nodules, separated by firm inflamed tissue. The lower lobe (red, and more solid and friable than natural) contained no cheesy masses; many small greyish yellow points (grey granulations) were sprinkled over the cut surface. The left lung, partly adherent at the base, contained many yellow nodules, very closely aggregated in the upper lobe, none of them softening. Grey granulations were numerous in this lung. They were also seen on the surface of the liver and spleen. The intestines contained no ulcers.

In this case the acute tuberculosis was in all probability secondary to the consumptive disease. The illness began suddenly with fever and cough, and the medical practitioner who was called in pronounced the case to be one of inflammation of the lung. The child presented many characteristics of the so-called scrofulous diathesis: the large, coarse features, and thick complexion; the long eyelashes and strongly-marked eyebrows; the eczematous

eruption about the nares,—all indicated a state of constitution favourable to caseous degeneration, and therefore one in which the absorption of a pneumonic deposit might be expected to be slow and incomplete. On the child's admission there was still some evidence of consolidation at the apex of the right lung, and this during the next few days became more marked. The temperature at this time was higher than the normal level, showing that the inflammatory process was not yet at an end. Shortly afterwards there was a rapid extension of the physical signs, and in the course of a few days all resonance completely disappeared from the front of the affected half of the chest, and even the resonance laterally and at the back was appreciably diminished. The occurrence of dulness had been preceded by clicking sounds, very superficial, and appearing to be generated immediately underneath the thoracic wall. This superficial character of the sounds, together with the sudden cessation of all rhonchus when the dulness was established; the rapidity with which resonance was lost; and the manner in which the change occurred—the dulness not travelling from the apex downwards, but involving at once the whole right front of the chest down to the extreme base—these facts, combined with the character of the percussion-note, which was completely dull, and the little alteration in the ratio between the pulse and the respiration, seemed to indicate that the diminution in resonance was not due to any sudden extension of the pulmonary consolidation, but resulted from the presence of pleuritic exudation. The pleurisy was of the plastic variety, as was shown by the limited distribution of the complete dulness. If fluid had been present, the loss of resonance would have been as complete at the back as at the front, unless, indeed, the pleurisy had been localised and the gravitation of fluid had been prevented by pleuritic adhesions. That the pleurisy was general, and not local, was indicated by the perceptible alteration of the percussion-note at the back, although

the dulness was much less complete posteriorly than in the front. In order that no doubt might remain as to the absence of fluid, the chest was punctured with the hypodermic syringe, but nothing escaped.

The softening of the cheesy consolidation at the apex of the lung went on rapidly ; the breath-sounds soon became cavernous, and large metallic bubbling rhonchi were heard at the supra-spinous fossa. The final attack of diarrhoea occurred rather unexpectedly, and carried off the patient before the symptoms of the secondary tuberculosis had become sufficiently developed to allow of the diagnosis of the complication. The grey granulations found on inspection of the body were probably of very recent origin ; and, no doubt, had the child lived a few days longer, some evidence of their presence would have been discovered in the left lung. In this lung some unsoftened cheesy nodules, and many fine grey granulations were found, but no irritation appeared to have been excited by their presence in the organ. It is remarkable that this child, although the subject of so exhausting a disease, and although suffering from so much pyrexia, did not lose flesh. On the contrary, the weight of his body increased from 16 lbs. 3 oz. on his admission, to 18 lbs. 6 oz. on April 29, when the attack of diarrhoea occurred which put an end to his life.

The diagnosis of chronic catarrhal pneumonia is more difficult in the child than it is in the adult. This arises partly from the physical peculiarities of children before referred to, partly from the frequent occurrence of wasting in early life independent of organic disease, and partly from the readiness with which a slight cough may be induced in young subjects by irritation of distant organs,—so that this symptom becomes a common complication of other complaints. A child may waste for a considerable time, and be troubled with a teasing cough, without being the subject of chronic pulmonary disease ; and, although pulmonary

consumption is always feared in such cases, the most careful examination of the chest may reveal nothing to encourage our suspicions. In the condition which I have elsewhere described under the name of "mucous disease," * a hacking cough is a common symptom, and this may continue for months without being followed by disease in the lung. Elevation of temperature, combined with the above symptoms, is a very important addition, and, if persistent, at once marks the case off from the ordinary forms of wasting disease. Whenever fever is present it is necessary to assure ourselves of the absence of common sources of irritation which might give rise to pyrexia. Thus, the mouth should be inspected for signs of aphthæ or inflammation of the gums. Dentition produces a rise in temperature, which may persist for weeks together; but here the pyrexia is very irregular, and the temperature rises much higher than is common at the beginning of chronic catarrhal pneumonia, being often as high as 104° or 105° in the morning. In children between five and six years old, such a cause of fever is often seen, and at this age wasting from mucous derangement of the bowels is most frequently met with. In these cases, therefore, the gums should be looked at, and will often be found to be tense and swollen over the position of the back molars, which are the first teeth of the second crop to make their appearance. Still, it must not be forgotten that any of these causes of pyrexia may be complications of the pulmonary complaint.

Whenever catarrhal pneumonia is suspected, the chest should be examined repeatedly for signs of disease. Such signs, to be trustworthy, must be positive. The degree of expansion in a child's lung is so variable, especially at the apex, that a distinct dulness must be required; and this dulness must be found to persist for several days in succes-

* See "Wasting Diseases of Infants and Children," 4th edition.

sion before we can be sure that we are not deceived by insufficient expansion or temporary collapse.

It is often difficult to distinguish between the beginning of catarrhal pneumonia, when the inflammation spreads gradually from the smaller tubes to the terminal vesicles, and cases of acute tuberculosis. In the latter case the loss of flesh is much greater than in the former, and on careful inquiry the pyrexia and wasting will be found to have preceded the cough, the general symptoms appearing before the local. In catarrhal pneumonia the local symptoms are the first to appear, and, unless the case have been complicated by some other form of wasting disease, there is a history of cough, and perhaps of shortness of breath, for some time before the general nutrition of the body began to suffer. Besides, in acute tuberculosis the course of the disease is rapid, the face at an early period wears a pinched, haggard expression, and the child has the appearance of one seriously ill.

In a more advanced stage of catarrhal pneumonia, when the signs of consolidation are pronounced, the diagnosis is easy. Dulness, bronchial breathing, and crepitating râles confined to the upper part of the lung on one side of the chest; the other lung being perfectly free from disease;—these signs found in a child the subject of a chronic cough, and in whom general nutrition, although impaired, is still fairly performed, may be safely attributed to chronic caseous consolidation. If, instead of the apex, the consolidation be seated at the base of the lung, the case has to be distinguished from empyema. It is rare for a chronic consolidation to be mistaken for empyema, but empyema, when accompanied (as it often is) by great emaciation in children, is constantly mistaken for “phthisis.” The diagnosis is made by referring to the history of the case—pleurisy almost invariably beginning with sudden and severe pain in the side, followed after some hours by a cough,—and by noting the character of the physical signs. If the dulness

be due to fluid, it is found at the front of the chest as well as at the back. The percussion-note is generally tubular under the clavicle of the affected side, and the respiration is either very weak, or is bronchial, blowing, or cavernous, without rhonchus. In addition, the outline of the diseased half of the chest, as taken by the cyrtometer, is found to have become more or less square-shaped, and the heart is displaced. If there be any vocal vibration on the healthy side, it is lost over the seat of disease. For a fuller discussion of the diagnostic characters of empyema, the reader is referred to a previous chapter. (See p. 113.)

When the case has advanced to the stage of ulceration and cavities have formed in the lung, we have still to satisfy ourselves that the cavities are really the result of destruction of tissue, and are not mere dilatations of bronchi. The means by which this distinction is arrived at have been already pointed out. (See p. 177.)

The probabilities of absorption in a case of a pulmonary cheesy mass are not easy to estimate. Much will depend upon the mode of formation of the deposit, upon the constitutional condition of the patient, and upon the general influences which can be brought to bear upon the complaint. As a rule, the more rapidly a deposit has formed, the greater is the probability of its safe removal, and therefore, in cases where the consolidation has resulted from an acute attack, the prognosis is more favourable than in cases where the disease has run a slow course from the first. The longer, however, the deposit remains unabsorbed, the fainter becomes the prospect of its disappearance; and although, in children, absorption may occur at a later period than is common in the adult, yet if months pass by without any alteration in the physical signs, it is unlikely that the affected lung will ever return to a completely healthy condition. The influence of the so-called scrofulous diathesis is no doubt to favour the occurrence of catarrhal processes, and also to determine an early caseation of pul-

monary deposits. Therefore, in children who present the characteristic marks of this type of constitution, any delay in the resolution of a pneumonic consolidation should be regarded with especial anxiety. So, also, in children with narrow contracted chests and feeble resisting power, the liability to pulmonary catarrh and the tendency of such catarrhs to become obstinate should not be lost sight of.

When the softening has begun in the deposit, the case assumes a very grave aspect, and as at this time the opposite lung is apt to become affected, recovery now can hardly be looked for. Still, under favourable conditions, the further progress of the disease may be greatly delayed. Persistent elevation of the temperature (other causes of pyrexia being excluded) is a very unfavourable sign, especially if combined with steady loss of flesh ; but even these cases do not always run an uninterrupted downward course, and there are often intervals of improvement, or, at any rate, periods during which the disease seems to be stationary. The occurrence of acute attacks of catarrhal pneumonia is necessarily dangerous, but the immediate effects are not always so serious as might be supposed, although the inflammatory attack seldom subsides without leaving behind it some addition to the original disease.

The occurrence of fibroid induration is to be welcomed as one of the favourable modes of ending of a cheesy deposit, for this change, if not always equivalent to a cure, is yet one which, with proper care, will greatly prolong the patient's life.

In cases of bronchial catarrh, when we have reason to believe that the inflammation is extending from the smaller tubes to the air-cells, treatment should be prompt and active. The child must be at once confined to bed, hot linseed-meal poultices should be kept applied to the chest, talking must be forbidden, and saline remedies should be ordered, with small doses of antimonial wine to promote

free secretion from the bronchial tubes. By this means the disease is often at once arrested, and after a few days, the pyrexia having subsided, the patient is able to return to his usual habits.

If the child be first seen after the formation of a pulmonary deposit, we may hope that the cheesy mass, so long as it remains inactive, may yet be absorbed. To favour its removal we should endeavour as much as possible to alter the conditions of existence under which nature had failed to restore the lung to a healthy state. If it can be done, we should send the patient into the country, choosing some elevated spot where the soil is porous and the air dry and bracing. Dryness of the air is a very important point, and, therefore, an inland district is more to be recommended than the seaside. The spot selected should, if possible, be sheltered from the north and east. The child should be warmly dressed, and should pass as much time as possible out of doors. In the house every care should be taken to keep the air sweet and pure ; in fact, a liberal supply of pure, dry air is one of the most important curative agents we have at our command.

The diet of the child should be regulated with great care ; he should take plenty of meat and milk, but starchy food, sugar, and all such fermentable articles of diet, are to be allowed sparingly. It is of extreme importance in these cases to keep the digestive organs in the highest state of efficiency, and, therefore, the tendency to acid fermentation of food, which is one of the peculiarities of early life, must not be encouraged by overloading the stomach with farinaceous food and saccharine matters. For the same reason, summer fruits are to be given with caution, and should not be allowed late in the day. It is important that the child take plenty of liquid, so as to favour local absorption ; and in order that the digestion of food may not be disturbed, it is wise to accustom the child to drink freely of some harmless fluid at an interval between the meals.

A careful watch should be kept over the bowels, and the earliest appearance of looseness should be met by suitable treatment. The best medicines are the alkalies, and they may be usefully combined with iron and arsenic. Thus, the bicarbonate of potash or soda may be given with the citrate of iron and ammonia, or of iron and quinine, and a few drops of Fowler's solution of arsenic may be added. Arsenic, it must be remembered, agrees well with children and between the ages of five or six and twelve it may be given to them in larger doses than are readily borne by the adult. Cod-liver oil—especially Kerr's natural light brown oil—can also be given. In cases where the appetite is bad, and great difficulty is found in persuading the child to take food, alkalies are again of service. Ten grains of bicarbonate of soda given three times a day with two drops of dilute hydrocyanic acid (P.B.) in a dessert-spoonful of some bitter infusion, such as chiretta, will often quickly restore the appetite after dilute nitric acid and quinine have been tried without any improvement. As a local application much benefit results from painting the chest over the seat of disease with linimentum iodi, and this should on no account be omitted.

When the occurrence of pyrexia announces that irritation is set up in the system by the presence of the pulmonary deposit, the child should be put to bed, or at any rate confined to one room : his chest should be well poulticed, and at the same times efforts should be made to reduce the temperature by suitable medicines. Arsenic, alone or in combination with quinine, is very valuable for this purpose, and a mixture containing quinine, digitalis, and opium, on the principle of the pill recommended by Niemeyer, can often be given with very striking results. While the pyrexia continues, the child should be kept as quiet as possible. He need not be confined to bed after the first few days, but every care should be taken to avoid chills, and to keep him as much as possible in an equable temperature. The hypo-

phosphites, especially the hypophosphite of lime, are often very useful in these cases, while the pyrexia is high ; and when softening has begun in the deposit the influence of these remedies is often very remarkable. It is at this time, too, that cod-liver oil is of especial value, and should always be given if it can be borne.

If the cough is troublesome, special medicines must be used to control it, but care should be taken not to derange the digestion by opiates, or to irritate the bronchial mucous membrane by the untimely use of stimulating expectorants. The principles which govern the treatment of an ordinary bronchial catarrh, apply equally to cases of chronic pulmonary disease. A hard cough is only made harder by ammonia and squill ; and opium, although it may for a time deaden the desire to cough, will not relieve the irritation of a congested and swollen mucous membrane. In such cases, alkalies, with small doses of antimonial and ipecacuanha wines, should be prescribed to relieve the loaded vessels ; and it is only when the cough has become perfectly loose, and all oppression of chest has passed away, that bronchial stimulants and small doses of morphia can be permitted.

In the latter stages of the disease, when our only object is to relieve painful symptoms, we must watch carefully over the digestive organs and the state of the bowels. The distressing sickness, which often occurs at the end of a paroxysmal cough, is best controlled by solution of arsenic in one-drop doses, combined with pernitrate of iron, and small quantities of morphia ; or, failing that, by the application of a very small blister under the clavicle of the side principally affected. If hæmoptysis occurs, the liquid extract of ergot should be given at once, with (if the bowels are not ulcerated) laxative doses of Epsom salts ; and perfect quiet should be enforced. In all cases of hæmoptysis the quantity of fluid taken at one time should be considerably restricted, so that the vessels may not be exposed to

any sudden strain from the rapid absorption of large quantities of liquid. Night-sweats are best subdued by belladonna given at bedtime, and in prescribing this medicine the remarkable tolerance of children for this drug must not be forgotten. For a child four years old the night draught should not contain less than twenty drops of the tincture.

Intercurrent attacks of acute catarrhal pneumonia should be treated by confining the child to bed, enveloping the chest in hot poultices, and keeping up the strength by the judicious administration of stimulants. For a child the best stimulant is the brandy-and-egg mixture of the Pharmacopœia.

CHAPTER VII.

FIBROID INDURATION OF THE LUNG (CIRRHOSIS).

PATHOLOGY—The disease always a secondary one—usually a consequence of catarrhal pneumonia or pleurisy.

Fibroid phthisis.

Case XXVII.—Fibroid thickening of lung following an attack of catarrhal pneumonia—Comment.

Symptoms and physical signs.

Cases XXVIII. and XXIX.—Fibroid induration of apex of lung with contraction of chest—Comment.

Case XXX.—Extreme fibroid induration and contraction—Comment.

Case XXXI.—Extreme fibroid induration with great dilatation of bronchi, but without contraction of the chest—Comment.

Case XXXII.—Fibroid induration with extensive excavation of lung—Comment.

Diagnosis of fibroid induration—Prognosis—Treatment.

FIBROID induration of the lung is a disease which is frequently met with in the adult, and has been often described. In children, being less common, it has received comparatively little notice, but it is not the less interesting and important.

The disease consists in a gradual thickening of the partitions between the air-cells by a dense fibroid tissue, which forms an interlacing network, spreading in all directions through the alveolar walls. Sometimes fibro-nucleated growths can be detected, but these are often absent, so that the induration has been described as depending upon the infiltration of an amorphous substance between the interstices of the alveoli. After a time the new material shrinks, contracting the lung and dilating the bronchi. In the

densest parts the air-vesicles may be completely obliterated by the new fibre-formation and its accompanying contraction ; and the cut surface of the lung is then seen to be smooth, shining, firm, hard, and non-granular. It is traversed by whitish bands, which are the thickened interlobular septa, and by thickened bronchi, whose channels are sometimes dilated, sometimes nearly or even quite obliterated by the new fibrous growth. In this stage the varying amounts of black pigment give to the surface a grey, marbled appearance, which is very peculiar and characteristic. The indurating process is not confined to the interlobular septa, but attacks also the walls of the vesicles in the interior of the lobules. It is associated with an accumulation within the vesicles of nucleated and epithelial cells, granular corpuscles, and granules—the result of an accompanying inflammatory process. Some pathologists are disposed to look upon the new fibroid tissue as a modification of the grey granulation. The complete dissimilarity in the clinical characters of the two diseases is, however, hostile to the ready acceptance of such a theory. Fibroid induration in young subjects seems to be always a secondary lung affection ; and the diseases to which, in the large majority of cases, it is found to succeed, are catarrhal pneumonia and pleurisy.

Catarrhal pneumonia in children is apt to assume a sub-acute character, especially when it occurs in the course of measles and whooping-cough. Bronchial dilatation is also a common accompaniment of this form of inflammation of the lung ; and the presence of bronchiectasis seems to have the effect of prolonging the course of the inflammation and delaying its resolution. It seems probable, too, that dilatations of the air-tubes continuing after the subsidence of the catarrhal disease, tend to excite a recurrence of the inflammatory process, and to increase the fibroid thickening. The dilatations become filled with secretion, which they are unable to expel. The retained secretion putrefies, and excites

great irritation of the surrounding tissue. A localised catarrhal pneumonia is set up, and the inflammatory process runs a tedious course on account of the continued irritation produced by the decomposing contents of the tubes. By this means the fibroid thickening is increased, and frequent recurrence of these attacks may lead to extensive induration of the lung. In some cases the irritation excited by the retained and fetid sputa may be so great that ulcerative destruction of tissue is set up, starting from the bronchial tubes. The disease in this form has been named "fibroid phthisis" by Sir Andrew Clark.

Pleurisy is the other cause to which many of these cases of fibroid lung disease may be attributed. In such cases the lymph appears, as it were, to creep into the lung, and the induration extends inwards from the pleura.

The following case offers an illustration of an early stage of the disease, and shows the influence of catarrhal pneumonia in producing thickening of the fibrous tissue of the lung.

CASE XXVII.—FIBROID THICKENING OF LUNG, FOLLOWING AN ATTACK OF CATARRHAL PNEUMONIA.

Charles S., aged six years, was admitted into the East London Children's Hospital on November 18. He came of a healthy family, and all his brothers and sisters were alive and well, except one who had died of diarrhoea. The boy had never had pertussis or measles, but had been for three years subject to a frequent cough. Two years previously, he had had an attack of scarlatina. Since that time his cough had been more troublesome, and he had occasionally expectorated mucus tinged with blood. During the last six months he had been losing flesh, and for a month before admission had complained of pains in the left breast and epigastrium, which were increased by taking food.

On admission there were no very positive physical signs.

The percussion-note over the whole of the right back was thought to be too high-pitched, and the same was noticed in the left axillary region, but there was no positive dulness. Elsewhere the note was healthy. The respiration everywhere was normal, and dry rhonchi were heard generally about the back.

Very shortly after admission the boy began to improve. His cough diminished, and he was allowed to leave his bed. The temperature varied in the evening from 100° to 101° , and his breathing was uniformly high, ranging from 40 to 48; but the appetite was good; he seemed cheerful, and did not appear to suffer much inconvenience from his illness.

On December 18 he caught cold. His cough became more severe, and the temperature rose in the evening to 102° . The pulse also became quicker, and the breathing accelerated. On the next evening (December 19) his temperature was 102.6° , the pulse 126, and the respirations 54; and from that date until December 27, when my attention was first directed to him by the house-surgeon, his evening temperature varied from 103° to 104.8° , his pulse from 126 to 156, and his respirations from 48 to 66.

On December 27 the following note was taken:—Boy rather small for age, but in tolerably good condition. Chest well formed, limbs straight, fingers distinctly clubbed. Lies in bed with head low; face flushed; nares acting; pulse 140; respirations 68, but respiratory movements not at all laboured, and no signs of dyspnoea; temperature in rectum 103.2° .

The right side of the chest moves more freely than the left. Heart's apex at fifth interspace in nipple line. The percussion-note is dull over the whole of the left side of the chest, but the dulness is greater below than above. Thus, in front complete dulness begins at third rib; behind at the spine of the scapula; and laterally dulness is only complete in the infra-axillary region. Resistance is only moderately increased. Vocal fremitus can be felt indis-

tinctly, and rhonchal fremitus is strong over the whole left back. With the stethoscope a subcrepitant rhonchus is heard all over the side with both inspiration and expiration. Above, it is coarse, but becomes finer towards the base, where it is mixed up with a sibilant rhonchus. The loudness of the rhonchus renders the breath-sound indistinct above, but from the angle of the scapula to the base the respiration is loud and intensely tubular, with bronchophonic resonance of the voice, and also at a spot a little to the inside of nipple it is tubular although less loud. The right side of the chest appears healthy except at the base behind, where the percussion-note has a wooden quality, and the respiration is blowing, with subcrepitant rhonchus. At the right supra-clavicular region also the percussion-note is a little high-pitched, but the breathing is natural. An occasional sibilant creak is heard over the right front.

On December 31 there were distinct evidences of consolidation of the right apex. Percussion was dull both above and below the clavicle, and also at the supra-spinous fossa. At the latter spot the breathing was weak, with sibilant rhonchus and coarse crepitation, while in front it was harsh, with a sibilant wheeze. At the right base behind there was also some dulness, and the respiration was faintly tubular, with coarse crepitation.

The left apex seemed, on the contrary, to be clearing, for the percussion-note at the supra-spinous fossa was only slightly high-pitched, and in front was resonant as far down as the upper border of the third rib. Above the left clavicle the respiration was diffused-blowing, with a fine crepitation; elsewhere at the upper part of that lung the breath-sound was quite concealed by loud subcrepitant rhonchus. In other respects the physical signs were the same as at the last report, with the exception that the rhonchus at the left back had a more distinctly bubbling character, and was more metallic in quality than before. Since December 27 the temperature had been very high, being seldom less than 105°

in the evening, with a fall to 102° or 103° in the morning ; pulse 140 to 145 ; respirations 60 to 84, slower in the morning than the evening.

On January 3 the signs of consolidation of the right apex were still more marked. The dulness reached in front as far downwards as the second rib, and the breathing there was bronchial, with coarse subcrepitant rhonchus. The same rhonchus was heard all over the right back, although the percussion-note was only dull at the apex and the extreme base. At the left back the dulness reached no higher than the spine of the scapula, and the supra-spinous fossa was quite resonant. In other respects the physical signs had undergone little alteration. Vocal fremitus was felt equally on both sides posteriorly. The coarse subcrepitant rhonchus was also heard over both sides of the back, and at the left base had developed into a large metallic bubbling. The temperature continued very high as a rule, but on the evening of January 1, after very profuse perspiration, it fell to 97.8° in the rectum ; by the next morning it had, however, returned to 104° . The cheeks continued flushed in the latter part of the day, and the nares acted slightly, but there were no signs of dyspnoea, although the number of respirations varied from 50 to 60 in the minute. The child's appetite was poor, and he was very thirsty, but he only coughed occasionally, and then sometimes expectorated offensive purulent matter. His breath had also somewhat of a putrid odour.

On January 7 there was great improvement. The temperature since last report had fallen, being seldom over 100° ; and his respirations were also much diminished in frequency—40 to 48. On the previous evening, at 7 p.m., temperature 99° ; pulse 96 ; respirations 42. The foetor of breath and sputa had been removed by the use of a creasote inhalation. The physical signs had also undergone a favourable change. At the right back the percussion-note was now healthy, but the respiration was faintly bronchial, with the

same rhonchus becoming rather metallic at the base. Vocal resonance was rather stronger than natural. At the left back the percussion-note was only completely dull from the angle of the scapula to the base. A loud metallic subcrepitant rhonchus accompanied the respiration. This, although coarse, was less large and less like gurgling than before, and at the base was mixed up with a dry, creaking rhonchus. The vocal resonance was still intensely bronchophonic over the lower half of the left back. The metallic subcrepitant rhonchus was heard all over the left front and side of chest, but only slightly at the right front, and the percussion at the right apex anteriorly was merely high-pitched, and no longer dull as before.

The child now began to improve. His temperature remained low, and the pulse and respiration diminished in frequency. His appetite also improved; he looked brighter in the face; was more cheerful and lively, and began to gain flesh. He had occasional attacks of diarrhoea, which rather depressed him at times, but these, after lasting a day or two, subsided readily under the influence of treatment.

The physical signs also improved very greatly. On January 17, at the left back the percussion-note, although generally high-pitched, was only dull at the base, where the respiration was rather blowing, and was accompanied by a faint, fine rhonchus. At the right side, posteriorly, resonance was natural and the respiration was healthy, except at the base, where it became faintly blowing in quality. Vocal fremitus was stronger on the left side than on the right. In front, the physical signs were natural, except that at the left side the breath-sounds were high-pitched and harsh, with here and there a little creaking rhonchus, and in the infra-axillary region some metallic crepitation. Last evening temperature 99° , pulse 84, respirations 29; this morning, temperature 98.4° . On this date (January 17) the bowels became again relaxed, and for several days after-

wards the motions, although not very frequent, were slimy and unhealthy-looking.

After this the improvement in the general health still continued, and the temperature remained little higher than natural— 99° to 100° ; the respirations were seldom under 50, and the pulse varied from 100 to 120. The left side continued to move less freely than the right, and the left intercostal spaces sank in more deeply in respiration. This state of things continued until February 5, the physical signs undergoing little change, except that the crepitating rhonchus reappeared on both sides, and became again very metallic at the left base behind.

On February 5 the temperature rose in the evening to 101° , and the pulse and respiration again increased in frequency. From this date the temperature remained high until the end. The severity of the general symptoms also returned: the appetite failed; the strength became reduced; and the boy began again to lose flesh. The physical signs also began slowly to increase. The dulness crept higher up in the left back, the breathing became more generally tubular, the bronchophonic resonance of the voice was no longer limited to the base, and the rhonchus was everywhere on both sides of the fine bubbling variety, with more or less of a metallic quality.

On February 14 the dulness reached at the left back as high as the spine of the scapula; and at the right back the percussion-note seemed less resonant than before, with greater resistance. All over both sides posteriorly bubbling rhonchus was heard, which at the left base became very large, metallic, and ringing in quality. The respiration there was almost cavernous, with intense bronchophony. Elsewhere at the back it was blowing or tubular. Temperature at 5 p.m. 102.6° .

On February 21 the boy was very much worse. His appetite was very bad, and he would take nothing but milk and broth. He often vomited, and was so weak that he

could hardly sit up in bed. The cough was frequent and loose, and the breath and expectoration had again become very offensive. There was no œdema of the legs.

At the left back percussion was dull from apex to base, with cavernous respiration even at the supra-spinous fossa. Rhonchus was heard only after a cough, and then was gurgling. Vocal resonance intensely bronchophonic. At the right back percussion was high-pitched as before, and dull at the base. Respiration bronchial or diffused-blowing from apex to base, accompanied, except at the apex, with small metallic subcrepitant rhonchus. Vocal resonance stronger than natural, but not bronchophonic. In front the percussion-note was dull all over the left side, with increased resistance; in the axillary and infra-axillary regions the note was rather tubular, with natural resistance. Respiration was everywhere on this side cavernous, with faint and rather distant small-sized bubbling rhonchus accompanying both expiration and inspiration. Vocal resonance natural. The right front was resonant, with loud, rather snoring, respiration, and no rhonchus. Vocal resonance natural. Temperature in the morning 100.2° , pulse 135, respiration 84; temperature in the evening 103.2° . The temperature continued high, with very rapid pulse and respiration. On February 23, at 9 a.m., temperature 103° , pulse 190, respirations 80; at 5 p.m., temperature 104° , pulse 188, respirations 86; and the child died on the following morning.

Autopsy.—Both lungs slightly adherent towards the base, but the adhesions feeble and easily broken down. Left lung small and shrunk, feels very solid and firm, and no part crepitates on pressure except a spot on the outer surface at the juncture of the upper and lower lobes, corresponding to the left axillary region. Bronchi, both large and medium-sized, very much dilated. Mucous membrane much injected. On section the surface is deep-red, not granular, and is marked with greyish intersecting lines or streaks, which are

the thickened interlobular septa, so that the lobules are unusually well defined. All over the surface are greyish points about the size of a small pin's head. These are evidently the sections of divided thickened bronchioles. From many of them minute yellowish semi-solid masses can be squeezed, and from others a thick yellowish fluid exudes on pressure. The substance of the lung is very firm, and is only broken down with difficulty by pressing the finger into it, but a section tears tolerably readily. Small portions sink in water. On section of the lower lobe the colour is a darker red, and the bronchi are excessively dilated. The dilatations are fusiform. In other respects the characters are the same as those already described. No grey granulations or cheesy masses are to be seen anywhere, nor are there any signs of ulceration of the bronchial tubes.

Right lung considerably larger than left. The two superior lobes crepitate pretty freely on pressure. The lower lobe and inferior half of the middle lobe alone are solid to the touch, and these on section show a dark-red non-granular surface, over which are scattered yellowish points of the size of a small pin's head. On pressure some of these escape as small masses of semi-solid yellowish matter similar to those squeezed out of the left lung; others cannot be so pressed out. The bronchial tubes are dilated, although to a less extent than in the opposite lung, and the fibrous tissue has not undergone the same development, for much fewer sections of bronchioles can be detected. The upper lobe is bright-red in parts, but still crepitates and is not easily broken down. At the apex a spot of collapse is seen forming a kind of cap to the summit of the lung. It is very superficial, and the part of the lung which it overlies is very red and crepitates less perfectly than the other parts of the same lobe. The bronchial tubes in this lobe are somewhat dilated, but to a less extent than elsewhere. There are no ulcerations. No grey granulations or caseous deposits can be seen in any of the organs. The other organs all seem

healthy, with the exception of the liver, which is very firm and rather fatty.

In the above case the fibroid thickening was only in an early stage. It was probably not entirely dependent upon the attack of pneumonia, from which the child suffered soon after his admission into the hospital. The history pointed to some chronic lung-mischief of considerable standing, for the boy had been subject for three years to a frequent cough, and this for two years had been very troublesome, and had been accompanied by the expectoration of purulent matter streaked occasionally with blood. A chronic bronchitis of such duration would probably lead to some increase in the fibrous tissue of the lungs, and to some dilatation of the bronchi. The insignificant character of the physical signs on the child's admission, and the absence of blowing or cavernous breathing—a change in the respiratory sounds which became afterwards very marked—seemed, however, to show that fibroid thickening, if present, was of small extent, and that the bronchiectasis was inconsiderable in amount; but it is possible that the degree to which the disease had advanced may have been concealed by the presence of compensating emphysema. The quality of the percussion-note was found, on the boy's admission, to be altered from a state of health; and the clubbing of the finger-ends indicated some long-standing impediment to the circulation, such as would be produced by a difficulty in the passage of the blood through the lungs. Besides, the rapid extension of the physical signs of consolidation to the front of the chest during the first attack of pneumonia—a phenomenon very unusual in an ordinary case of pulmonary inflammation—seemed to indicate a very abnormal condition of the lung.

At the time when I first saw the patient (December 27) the case presented many points of resemblance to one of pleurisy. The left side of the chest was completely dull over its lower half, both before and behind, and the breathing there was loud and intensely tubular, with bronchophonic

resonance of the voice—a condition often to be met with in the pleurisy of childhood. The heart, however, was not displaced; the dulness was not the dead, toneless dulness produced by a collection of fluid; vocal fremitus was not suppressed; and the subcrepitant rhonchus heard with the breath-sound differed widely from the superficial crackle heard in an inflamed pleura, and became finer and more like true crepitus at the base, where it was mixed up with sibilant rhonchus.

After admission into the hospital, the child suffered from two distinct attacks of catarrhal pneumonia. In the first attack the physical signs were those common to the subacute form of that complaint; but in the second a certain abnormal character was imparted to them on account of the extensive dilatation of the bronchi, which at this period had become decided. It was on this account that the crepitation, which was at first of the coarse variety common to that form of pneumonia, became much more bubbling and acquired an intense metallic ring, such as is often found in small cavities and softening caseous deposits. From the physical signs alone it would, no doubt, have been impossible to exclude breaking up of the lung, especially as the increasing weakness of the child, and his steady pyrexia, were quite compatible with such a condition. The frequency, however, with which severe catarrhal pneumonia is accompanied by dilatation of the bronchi, made it equally probable that bronchiectasis was the cause of this peculiarity in the auscultatory signs; and the post-mortem appearances confirmed this view of the case.

Besides the catarrhal pneumonia and the dilated bronchi, the autopsy showed a considerable excess of fibroid tissue in both lungs. The fibroid induration was as yet in an early stage. Had the boy recovered from the second attack of pulmonary inflammation, the bronchiectasis would no doubt have partially subsided, and a certain wooden quality of percussion-note, with harsh or bronchial breathing, might

have been the only physical signs remaining to indicate the abnormal condition of the lung. In fact, possibly little more would have been discoverable than was present when the child was first admitted into the hospital.

During an early stage of the disease the symptoms are very slight, and the physical signs are those which have just been enumerated. If the lesion extends over a considerable portion of the lung, the percussion-note is slightly high-pitched, with a certain wooden quality; resistance is rather increased; vocal fremitus presents nothing characteristic, for, if sometimes stronger, it is sometimes weaker than natural; the breathing is harsh or bronchial; and the vocal resonance, unless there be dilatation of the bronchi, is generally normal. If there be accompanying catarrh, some coarse bubbling or crepitating rhonchus may be heard at various parts of the chest. The patient is sensitive to cold, and prone to catarrhs and attacks of catarrhal pneumonia, but between such attacks the expansion of the side may be fair and the general health leave little to be desired.

When the induration is limited to the base of the lung, as it is sometimes found to be, as a consequence of an attack of pleurisy, there is dulness at the extreme base behind, with excessively weak bronchial breathing, and often coarse liquid râles. But fibroid induration from this cause is not common in the young subject.

As the induration of the lung increases and the fibroid tissue comes to be in great excess, the lung is reduced in bulk, the bronchi become dilated, and more or less contraction of the side is the result. The physical signs are then the following:—The wall of the chest immediately over the diseased part is flattened, and the respiratory movement is hampered, or, if the disease be extensive, even suppressed. The contraction of the lung draws in the soft parts in its neighbourhood, so that mediastinum is drawn towards

the diseased side. As a consequence of this, the opposite lung is found to project across the middle line of the sternum, and the heart is usually displaced. If the induration affect the upper part of the left lung, the heart is drawn upwards ; if the right lung be the one diseased, the heart is drawn towards the right side, and in extreme cases may be felt beating to the right of the sternum. In cases, however, where the pericardium has become adherent to the left pleura, this displacement is prevented. On account of the contraction of the chest-wall, the circumference of the thorax at the part affected is reduced. Vocal vibration is sometimes increased, but more frequently it is diminished, and is often hardly perceptible. The percussion-note is dull, with wooden or tubular quality, and resistance is usually, but not always, increased. The respiration differs in character according to the quantity of secretion in the tubes : if the tubes are nearly empty, it is of the blowing variety, either tubular or cavernous, and is generally very loud : vocal resonance is, as a rule, bronchophonic, and there is much clicking rhonchus mixed up with creaking sounds. If, however, there be great accumulation of secretion, the breath-sound is weak and bronchial, with but little rhonchus, and faint vocal resonance. The signs are in most cases limited to one side of the chest, as both lungs are rarely affected. Of the symptoms, the most characteristic is the violent paroxysmal cough, occurring at rare intervals and ending in retching efforts, and the expectoration of enormous quantities of offensive purulent matter. The foetor of the sputum is due to its retention in the tubes, on account of the loss of elasticity of the bronchi. The difficulty with which it is brought up is owing to the same cause, and in bad cases its expulsion is only effected by violent contractions of the diaphragm. The respiration is habitually rather hurried, and ranges from 30 to 40 in the minute. Hæmoptysis is rare, but is sometimes seen in older children. There is sometimes a tendency to diarrhoea ;

and in bad cases, towards the end of the disease, ulceration of the bowels may take place.

Unless the expectoration be very copious, or the diarrhœa troublesome, the general nutrition of the child is often good, for where the disease is not far advanced, emaciation is certainly not necessarily produced by such a condition of the lung. There is in many cases a considerable amount of plumpness, and the complexion is often healthy. When nutrition suffers, it is chiefly as a consequence of the catarrhal attacks to which these patients are so subject; but at the end of such attacks the temperature, which had been elevated, sinks to a natural level, the strength usually returns, and the child regains flesh and colour.

So long as it remains uncomplicated, fibroid induration of the lung does not give rise to pyrexia. If the temperature in any case is found to be high, the elevation is due, in the majority of instances, to the presence of catarrhal pneumonia, which is so frequent an accompaniment of the disease. Ulceration of the bronchial tubes will also cause the temperature to rise. In these cases the cause of the febrile symptoms may be discovered by a microscopical examination of the sputum. The expectoration must be boiled in a test-tube, with an equal quantity of caustic soda, and after a short time the mixture will be found to be converted into a perfectly transparent fluid. If any fragments of elastic tissue be present, they will sink to the bottom of the tube, whence they can be easily removed by a pipette, and placed under the microscope. Such fragments, if areolar, are conclusive evidence of ulcerative excavation.

The indurated lung offers a certain impediment to the pulmonary circulation. Consequently, if the disease be at all extensive, there is some enlargement of the right side of the heart, and the aortic venous system is fuller than natural; so that the veins of the neck and chest, and often of the limbs, are unusually prominent. The fingers soon

become clubbed. If any catarrh be superadded, the patients often present something of the congested, turgid appearance of the face which is seen in old emphysematous subjects.

In adults, a late stage of the disease is often accompanied by a certain amount of œdema of the extremities. This may, probably, be also true of children. In the only two cases, however, of fibroid induration complicated by œdema, which have come under my notice, the patients were the subjects of extensive amyloid degeneration of organs—a condition which would itself account for the serous effusion.

In the following case the induration of the lung is seen to have advanced to the stage of contraction. It presents a condition which is sometimes met with in children as a consequence of pneumonia :—

CASE XXVIII.—FIBROID INDURATION OF APEX OF LUNG WITH CONTRACTION OF CHEST.

Alice P., aged fourteen, was admitted into the East London Children's Hospital on March 24. Her illness had begun three years before with shivering, followed by fever and cough, and the girl kept her bed for a fortnight. Since that time the cough had continued, and she had slowly lost flesh. Last Christmas she became worse : her cough was more troublesome, she lost flesh faster, and often complained of feeling cold, and of pains in the left side of her chest.

Statement of Admission.—The girl is rather thin, but does not look ill. She has some colour in her cheeks ; appetite pretty good. She coughs a great deal, and expectorates thick greenish-yellow matter, but the sputa are not offensive, and there is no foetor of the breath. She states that some weeks ago the expectoration was much more profuse, and was rather offensive. It was brought up with retching efforts after prolonged fits of coughing. Temperature at

5 p.m., 99·8°. The left side of chest is flattened at the upper part, and is unnaturally hollow below the nipple. Respiratory movement very deficient on that side. Heart's apex half an inch below the nipple, and slightly to the inner side. The percussion-note is dull from above the clavicle to the third interspace. Just below the clavicle the note is rather tubular. Vocal fremitus is felt faintly above the clavicle, but is only just perceptible below over the dull area, and is very much weaker than on the opposite side. The respiration over the same area is cavernous, and is accompanied by a little metallic bubbling rhonchus and creaking sounds. At the base the breath-sound is excessively weak, and the same creaking and bubbling sounds are heard. The vocal resonance is bronchophonic. At the left back there is only dulness at the supra-spinous fossa, but the respiration from apex to base is bronchial and very feeble, with the same rhonchi as in front. Vocal vibration is faint, and vocal resonance is stronger than natural. In the axillary and infra-axillary regions percussion is natural. Respiration towards the armpit is blowing ; below it is merely weak and harsh. On the right side the chest is naturally resonant, both before and behind ; respiration is everywhere rather wheezing, with prolonged expiration ; and a little bubbling rhonchus, not metallic in character, is heard here and there. There is no creaking, such as is heard on the left side.

While in the hospital the girl improved considerably in general health, although during her stay she had an attack of pneumonia affecting the middle lobe of the right lung. This, however, passed off without producing any permanent injury, for the removal of the deposit appeared to be complete, although the percussion-note did not become perfectly resonant at the right scapular region until nearly three weeks after the beginning of the attack. The usual temperature, before the attack of pneumonia, was normal ; afterwards it remained a little higher than natural (99·6° to

100°). The pulse was, as a rule, just over 100 ; and the respirations varied from 30 to 38.

On April 26, shortly before the girl left the hospital, the physical signs were exactly the same as on her admission, and the bubbling and creaking rhonchus still persisted. Her general health had, however, greatly improved. She looked well, and had gained considerably in flesh.

Another illustration of the same condition is seen in the following case :—

CASE XXIX.—FIBROID INDURATION OF THE APEX OF THE LUNG WITH CONTRACTION OF THE CHEST.

Clara S., aged seven, had been healthy until the age of six years, when she had an attack of whooping-cough. Three months afterwards she was ill for six weeks with scarlatina, but had no dropsy. After this, however, her strength did not return, and three months before admission she caught cold and was very feverish. Since that time she had been losing flesh, and her cough had been very troublesome.

When admitted into the hospital, the child was pale, but tolerably stout. There was considerable flattening at the upper part of the left front, and the whole of that side moved less perfectly than the right during inspiration. The heart's apex was immediately to the inner side of the left nipple, and the impulse could be seen over the greater part of the præcordial region as the child lay on her back. On the left side there was dulness with increased resistance from above the clavicle as far downwards as the cardiac dulness. The percussion-note was also dull in the axillary region, and at the back from the supra-spinous fossa as far as mid-scapula. In front the dulness did not reach quite to the middle line, for the resonance of the right lung passed across the centre of the sternum and encroached somewhat upon the left pleural cavity. Over the whole area of dulness, vocal fremitus was excessively feeble, and this was

especially the case under the clavicle in front. At the base, however, and all over the right side, vibration was well marked. Respiration was loud and blowing, with prolonged hollow expiration above the clavicle. Below, it was weaker but still blowing, and near the acromial angle it was cavernous. The cavernous respiration was heard as far down as the nipple, becoming fainter as the stethoscope was moved downwards. At the base the breath-sound was merely harsh, and here and there a coarse superficial crackling sound was heard, which disappeared after a cough. In the axilla the breathing was loud and cavernous; behind it was merely harsh. Over both backs a little crackling rhonchus was heard, which was removed by coughing. Vocal resonance was everywhere bronchophonic over the dull area. On the right side the physical signs were perfectly healthy, except for the slight rhonchus before mentioned.

During the child's stay in the hospital the rhonchus disappeared, but otherwise the physical signs remained unaltered; still, her general health greatly improved, and she gained flesh. Her temperature was 99° , as a rule, both morning and evening, but occasionally it rose slightly, from some temporary cause unconnected with the pulmonary lesion.

In these two cases the fibroid induration occupied the upper part of the left lung, and the physical signs were sufficiently well marked to indicate with certainty the nature of the disease. In both cases the lung was evidently contracted, for the heart was displaced upwards, and the chest had fallen in. In the case of Clara S., it was noted, in addition, that the mediastinum was drawn to the left, so that the right lung encroached upon the left pleural cavity, and that the cardiac pulsations were plainly visible over the præcordial region—showing that the margin of the left lung had receded from between the heart and the wall of the chest. Such an amount of contraction of the lung should at once lead us to suspect something more than ordinary

caseous consolidation. Again, the vocal vibration of the thoracic walls was well marked on the healthy side of the chest ; but over the seat of disease it was almost completely suppressed. In the case of an unabsorbed pneumonic deposit, the vocal fremitus is increased over the seat of solidification. The auscultatory signs were also peculiar. Common bronchitic bubbling râles were mixed up with the creaking sounds which are considered characteristic of fibroid induration. Lastly, the stethoscope revealed signs of a cavity ; but a careful examination of the chest detected no indications of disease in the apex of the opposite lung. Now, it is well known that the softening process affecting a caseous pulmonary mass is very rapidly followed by signs of disease in the upper part of the lung on the other side of the chest ; and when the process has gone on to the formation of a vomica, the apex of the previously uninjured lung furnishes unmistakable evidence of disease. To this rule there are few, if any, exceptions. In the present cases, therefore, some other explanation of the cavernous signs must be sought for than that supplied by the supposition of a softening cheesy mass. Fibroid induration of the lung almost always leads after a time to dilatation of the bronchi ; but in the young subject signs of mischief in the opposite lung are rarely to be discovered—probably because absorption from the seat of disease is prevented by obliteration of the lymphatics of the part. In such a case, therefore, we should find evidences of consolidation of lung, with signs of a cavity, limited strictly to one side of the chest—the very conditions, in fact, which are met with in the two cases under discussion. In these cases the part of the lung concerned had been, no doubt, the seat of pneumonia, which had left behind it an unabsorbed cheesy mass. After a time the caseous matter had completely disappeared, and the lung was left with the connective tissue at the apex in great excess. The new tissue had contracted, indurating the lung ; consequently the bronchi had become

dilated, and the chest-wall had sunk in at the seat of disease.

In the following case the contraction of the lung, consequent upon fibroid induration, had reached an extreme degree. The case, indeed, presents a typical instance of the most advanced form of cirrhosis of the lung.

CASE XXX.—EXTREME FIBROID INDURATION AND CONTRACTION.

Elizabeth W., six years and a half old, had enjoyed good health until the age of four years, when she had an attack of whooping-cough. This was shortly afterwards followed by measles and varicella. The cough continued altogether for five or six months, and then disappeared completely, the child becoming again apparently perfectly well. Some time after the cessation of the cough, she was sent into Essex for several months, and while there had an attack of scarlatina, which was followed by suppuration of glands of the neck, but not by dropsy. She was ill altogether for three weeks.

In December, 1871, five months after the attack of scarlatina, the child began to cough again, and to expectorate large quantities of phlegm. Her body also began generally to swell. She became a patient at the Victoria Park Hospital, under the care of Dr. Baümle, who treated her for "chronic pneumonia secondary to scarlatina." At that time the physical signs, as reported by Dr. Baümle, were—"Tubular resonance over right front from apex to below nipple, at back from apex to spine of scapula. Within this area expiration prolonged, but no bronchial breathing. At right base, in front and at back, dulness, bronchial breathing, and râles. Some œdema of eyelids, especially of the right. Urine contains no albumen." Dr. Baümle did not remember that there was at that time any particular contraction of the right side of the chest.

On September 30, 1872, the child was admitted into the East London Children's Hospital. Her condition was then as follows :—Child very short for her age, and excessively anæmic. Skin of face thin and delicate ; eyelashes very long ; veins of forehead and of temples very full. Can stand, but limbs are very weak and thighs thin ; the legs are œdematous, and the right arm (the side on which she always lies) is slightly so. Ends of fingers clubbed. The skin of chest is dry and harsh—in fact, the skin generally acts imperfectly, for the child only perspires about the head and neck. Abdomen full, but does not fluctuate. The cervical glands are enlarged, freely movable, and painless, and one or two small glands are to be felt in the left axilla, but those of groins are hardly perceptible. Scars of old abscesses seen in neck. Liver and spleen large and resistant, and can be felt on each side about three inches below the false ribs. Urine scanty, seldom more than a pint in the four-and-twenty hours. It is straw-coloured, and contains about a third of albumen ; specific gravity 1024.

The cough is of a peculiar character ; it occurs in paroxysms at comparatively rare intervals. For hours together the child will not cough at all ; then a fit of coughing will begin, which lasts for five or more minutes, and ends in retching efforts and the expectoration of several ounces of frothy, purulent, offensive mucus.

The right side of the chest is visibly contracted, and is very much flattened at the upper part under the clavicle. The right shoulder and scapula are lower than the left, and the right nipple very much lower. Spine curved laterally, with convexity to the left. During inspiration there is little or no movement of the right side. Measurement, an inch above level of nipple on the right side, nine inches and fifteen-sixteenths ; on the left, eleven inches and a quarter. The heart's apex beats one inch below nipple in nipple-line ; sounds healthy. Vocal fremitus is stronger at the right supra-spinous fossa than at the left ; below, it

is weaker, although still perceptible as far as the angle of the scapula.

On percussion there is dulness of the whole right side of the chest from apex to base, both before and behind ; the note has a tubular quality in front from the clavicle as far as the level of the nipple, and behind at the supra-spinous fossa. In the axillary region there is tolerably good resonance as far as the level of the nipple. The resistance is nowhere much increased. Respiration is everywhere cavernous on the right side, with metallic bubbling rhonchus, and the vocal resonance is bronchophonic. The left side of chest is rather hyper-resonant, with loud, harsh breathing, and the resonance passes across the sternum to the right side, about the distance of one finger's breadth beyond the middle line. At the left back the respiration has rather a hollow quality, which appears to be conducted from the opposite side.

The child remained several months in the hospital, but without much alteration in her condition. The œdema varied in amount, and the physical signs underwent only slight changes, which appeared to be dependent upon the varying amount of purulent secretion in the tubes. Thus, if no expectoration had taken place for several hours, the respiration at the right base would be merely faint and bronchial, with vocal resonance weak or annulled. If now the child had a paroxysm of coughing and brought up a large quantity of sputa, the respiration at once became loud and cavernous, and the vocal resonance again became bronchophonic. The temperature was seldom materially elevated, and ranged usually from 99° in the morning to 100° in the evening ; the pulse varied from 115 to 125, and the respirations from 38 to 48.

In this case there were evident signs of contraction of the right lung. The diameter of the chest was very much diminished on that side ; the spine was curved, with the concavity to the right ; and the mediastinum was drawn in

the same direction, as was shown by the resonance of the left lung passing across the middle line of the sternum. The heart, however, was not displaced, owing, no doubt, to the existence of adhesions between the pericardium and left pleura. The falling of the shoulder, and lowering of the nipple on the right side, with the curvature of the spine, might have suggested pleurisy with retraction ; but there were many symptoms which pointed to a different conclusion. The contraction was not limited to the lower part of the chest, as in pleurisy, but was general over the whole side, and was indeed especially well marked above. On listening to the chest, loud cavernous breathing, with bronchophonic resonance of the voice, was everywhere audible, and the large metallic bubbling rhonchus, which accompanied the breath-sound, furnished by itself a strong presumption of the presence of bronchiectasis. Again, the offensive character of the sputa was very characteristic of bronchial dilatation, and the method in which this was expectorated—viz., at rare intervals, after a paroxysm of violent cough, ending in retching efforts—was a symptom very significant of fibroid induration of the lung, with great rigidity of the tubes.

The lung affection was not, however, the only disease from which the child was suffering. The size and firmness of the liver and spleen, the enlargement of the lymphatic glands in many situations, the albuminuria, and the marked anæmia, indicated the existence of amyloid degeneration. To this constitutional disease the emaciation, which was a prominent feature in the case, must no doubt be attributed, for in children fibroid induration of the lung is not necessarily accompanied by any loss of flesh. There was no history of syphilis, and the duration of the hepatic and splenic enlargement, as it had been unnoticed by the parents, could not be ascertained ; but no mention of any such disease occurs in the short note with which Dr. Bäumler kindly supplied me, and the absence of albumi-

nuria is especially remarked. The enlargement must, therefore, have occurred subsequently to the child's attendance at the Victoria Park Hospital, in December, 1871. It is probable that the profuse purulent expectoration from which the patient had suffered since that date may have been a determining cause of the disease in these organs. In early life there is a special tendency to amyloid degenerations, for the frequency with which such a result follows any suppurative drain upon the system, if the drain be at all prolonged, is a matter of common observation in a children's hospital.

The three preceding cases have been accompanied by falling in of the chest-wall; contraction of the lung is not, however, a necessary consequence of fibroid induration, even where the disease is of considerable standing, as will be seen in the following case:—

CASE XXXI.—EXTREME FIBROID INDURATION WITHOUT
CONTRACTION OF SIDE.

Jacob P., aged five years, was admitted into the East London Children's Hospital on February 20, 1874. The boy was said to have been healthy up to the age of sixteen months, when he had an attack of whooping-cough. Since that time he had been subject to a cough, and had frequently expectorated thick yellow phlegm. He had never had measles. A week before admission he began to complain of pain in the right side, and in the front of his chest. His cough became severe. Still his appetite remained fair, and he did not seem to lose flesh. Until the night before admission he slept well.

State on Admission.—Boy stoutly built, and rather short for age. In fair condition, rather fat than thin. Features coarse and thick. Complexion rather sallow; face a little congested, and resembles that of an old sufferer from winter cough. Chest well shaped. The right side moves little in

inspiration ; the intercostal spaces of both sides sink in. The nares act, but respiration is not at all laborious, and there is no recession of the lower part of the chest-walls during inspiration. The heart's impulse can be felt in the epigastrium, and, faintly, immediately below left nipple. A systolic murmur is heard all over præcordial region, and also in left axilla, and at the angle of left scapula. The percussion-note is dull all over the right side of the chest, both front and back, from apex to base. The note is not, however, completely toneless, and above the clavicle, although high pitched, is still somewhat resonant ; about the nipple the note becomes "cracked-pot" in quality. Resistance, although increased, is not markedly so, especially at back. The respiration is intensely cavernous everywhere on the right side, both before and behind. Vocal resonance is much stronger than natural, and a fine bubbling rhonchus, very loud, metallic, and ringing, is heard with both inspiration and expiration. About the middle of the right back the respiration is almost amphoric, with intense bronchophonic resonance of the voice. All over this side of the chest the vocal fremitus is very strong ; on the left side it is weak, the percussion-note is natural, and the respiration has a slight hollow quality, no doubt conducted from the opposite side. Measurement with the cyrtometer shows that the right side is rather larger than the left, measuring eleven inches and a half to eleven inches and three-eighths on the left side. Its outline also, as drawn by the cyrtometer, is less completely semicircular, being slightly flattened posteriorly and at the junction of the lateral and anterior thirds. Temperature at 6 p.m. $101\cdot6^{\circ}$.

On February 21, 8 a.m., temperature 101° ; pulse 120 ; respirations 48. At 7 p.m., temperature $101\cdot6^{\circ}$.

On February 24 the percussion-note had become less dull in front ; at the third interspace a "cracked-pot" sound could still be elicited. The respiration all over the right front of the chest was amphoric, and was accompanied by gurgling

rhonchus of an intensely metallic and ringing, almost musical quality. It was more intense and metallic at the base than above, and appeared to be seated immediately at the end of the stethoscope. Vocal resonance all over the right front pectoriloquous. At the right back the percussion-note was tubular, and was nowhere completely dull except at the extreme base, where the resistance was notably increased. Elsewhere resistance, although greater than natural, was not extreme. The vocal fremitus was stronger at right than at left back. Respiration at the supra-spinous fossa was cavernous; below, as far as the base, it was amphoric, and was accompanied by the same ringing, almost tinkling, rhonchus as in front. This was appreciably reduced in quantity by coughing, although its character remained the same. Vocal resonance at the back was intensely bronchophonic, the faintest whisper appearing to pass straight up the stethoscope. The cough was very violent, and during the paroxysms the jugular veins became much distended. Elsewhere the superficial veins were not unusually prominent. Since last report the temperature had varied from 102° to 103° in the evening, with a fall to 101° in the morning. The pulse had been slightly over 100, and the respirations 30 to 40. The boy's appetite was poor; his bowels were regular, and he slept well at night. No albumen was found in the urine.

After this date the case went on with little alteration in the symptoms for some weeks. The temperature continued high, being seldom less than 102° in the evening until March 11, when for the first time it was as low as 99° in the morning and 100.6° at 7 p.m. The boy was found to have gained two pounds since March 2, his weight being 2 st. 10 lbs. His appetite, too, had begun greatly to improve. The physical signs, however, had altered little. The dulness persisted over all the right side of the chest, and the note was tubular at the apex both before and behind. The vocal fremitus continued strong, and the resonance of

the voice intensely bronchophonic. The respiration was still amphoric, but less markedly so than before, except at the base in front, where it was very loud. The same metallic gurgling rhonchus accompanied both inspiration and expiration. The condition of the heart was unaltered. The murmur was still heard over the whole of the præcordial region but faintly at the base, and the apex was one inch and a half below the nipple and slightly to the outer side of the nipple-line.

On March 17 the dulness was found to be the same, but it was noted that the respiration had lost its amphoric quality and had become cavernous all over the right side of the chest. The bubbling rhonchus was still metallic, but without the musical tone before noticed, and the vocal resonance, although stronger than natural, was not bronchophonic. The vocal vibration still remained more marked than on the opposite side. The temperature was usually 99°, occasionally rising in the evening to 100°. Pulse generally 120 ; respirations 30 to 40.

On March 24 there were signs of clearing at the right apex. The percussion-note, although high-pitched, was tolerably resonant at the supra-spinous and supra-clavicular regions, and the respiration there was merely bronchial. The whole right back, indeed, was less dull than before, but the respiration was still cavernous, with the same metallic bubbling rhonchus. Over the front of the chest the percussion-note was dull below the clavicle, but resistance was almost natural. Respiration weak, and almost covered by the bubbling rhonchus. Vocal resonance was natural, except at base, where it was increased. In the axilla percussion was resonant, with slight tubular quality ; in the infra-axillary region it was still dull, with cavernous breathing.

After this date the physical signs gradually improved, and on April 10 the dulness was limited to the base, both before and behind, although over the whole side the percus-

sion-note remained more or less high-pitched. At the base the respiration was still cavernous, with crisp, large-sized bubbling rhonchus and increased vocal resonance. Elsewhere the breathing was bronchial, with the same rhonchus. Over the dull area the vocal fremitus was stronger than natural. The boy was now looking very well, and he coughed but little. His spine was quite straight, and the shoulders and angles of the scapulæ were on the same level. His chest measured on the right side ten inches and seven-eighths, and on the left eleven inches and three-eighths, one inch below the level of the nipple. His usual temperature was now 99° to 100°, pulse 108 to 120, respirations 34 to 40. He left the hospital on April 10.

On October 20 the boy was again admitted with a return of his symptoms. On this occasion there was noticed to be a little depression of the right shoulder, and the spine was very slightly curved, with concavity to the right. The circumference of the chest was now equal on the two sides. At the level of the ensiform cartilage, each side measured eleven inches and three-quarters. Enlargement of the liver was now noticed for the first time; the edge could be felt three fingers' breadth below the false ribs. The boy was fat and sturdy-looking. The ends of fingers markedly clubbed. His cough was violent, and often ended in retching, during which he expectorated large quantities of thin, offensive, purulent sputa. The breath was very offensive with the same putrescent odour as the expectoration. The physical signs were much the same as before. There was dulness at the lower half of the right side of the chest, both before and behind, but vocal fremitus was excessively faint over the dull area, although distinct elsewhere. Respiration was loud and cavernous, and was accompanied by large metallic bubbling rhonchus, and dry creaking sounds. Vocal resonance was intensely bronchophonic. Over the upper part of the chest, respiration was faintly bronchial, with non-metallic bubbling rhonchus. All over the left side of the

chest, respiration was loud and coarse, and was accompanied by faint and distinct metallic bubbling rhonchus—conducted, probably, from the diseased side. There was little movement of right side during inspiration. Heart's apex an inch and a quarter below left nipple, and half an inch to inner side. Soft systolic murmur heard as before at apex; heard also at angle of left scapula, and, faintly, at base of heart. Temperature, 7 p.m., 100·8°.

During the boy's stay in the hospital on this second occasion his temperature was habitually over 100° both morning and evening, and his respirations varied from 30 to 40, or even higher. His breath was always very offensive, and his sputa retained their putrescent odour, and were expectorated in large quantities at rare intervals. Occasional streaks of blood were noticed in the expectoration.

The boy left the hospital shortly before Christmas. At that time the dulness was only complete at the extreme right base, but the percussion-note was high-pitched, with increased resistance as high up as the nipple in front, and one inch above the angle of the scapula behind. Laterally, there was dulness in the infra-axillary region. Faint vocal fremitus was felt over the area of dulness. The respiration behind was cavernous, with gurgling rhonchus and bronchophony from the spine of scapula to the base; and in the supra-spinous fossa it was bronchial with the same rhonchus. In front the respiration was bronchial, with large bubbling râles at the upper part, but towards the base it became very weak and blowing, with crisp metallic bubbles. In the infra-axillary region the breath-sound was intensely cavernous, with gurgling rhonchus; and in the arm-pit it was blowing, with large metallic bubbles. There was some hypertrophy of the right ventricle, for the heart's impulse could be felt from below the left nipple to the right of the ensiform cartilage. The patient improved in flesh during his stay, and when he left weighed three stone within six ounces.

On February 5 he was brought again to the hospital. Immediately before the examination he had coughed violently, and had expectorated about an ounce of frothy, greenish, purulent mucus. His breath was still offensive. The fingers were much clubbed. There was only slight deformity of the chest. As the boy stood up, the angle of the right scapula was a very little (about half an inch) below the level of the left, and there was a scarcely noticeable curving of the spine, with concavity to the right side. The chest measured, one inch below nipple, on the right side eleven inches and three-eighths, on the left eleven inches and three-quarters. The base of the chest had now become nearly resonant; the percussion-note was dull on the right-side, above the clavicle in front, and from the supra-spinous fossa to midway between the spine and angle of scapula behind. Below these points to the base it was resonant, but rather wooden in quality. The same wooden or tubular note was found in the arm-pit and infra-axillary region. Everywhere on the right side the respiration was loud and cavernous, and was accompanied by large resonant clicking bubbles, which were more numerous and smaller at the base. Vocal resonance was everywhere intensely bronchophonic, and vocal fremitus was stronger than on the left side. All over the left lung the percussion-note was drum-like, and the resonance passed across the middle line, reaching to half an inch beyond the right edge of the sternum. Heart's apex as before.

On May 1, 1875, the boy was admitted for the third time. He was now excessively ill with very rapid breathing, high fever, and great prostration. The rare paroxysms of cough ended in retching and the ejection of large quantities of fetid purulent matter. Temperature on the first evening, 104.4° ; on May 2, 103.2° both morning and evening, pulse 132, respirations 52.

May 3.—Was found lying on right side, propped up in the bed. Expression anxious; nares dilating widely:

complexion dusky, purple on cheeks, lips, and end of nose, the rest of face and the neck of a peculiar earthy tint; eyes getting hollow. Had lost flesh considerably since his former visit. Seemed tender all over, and although generally very patient, cried if touched. Seemed to be drowsy, but was very restless, and could not sleep long at a time. At the back the right side was wanting in resonance all over, and was completely dull from the angle of scapula to the base. The respiration for the lower two-thirds was cavernous, with a peculiar high-pitched, ringing, stridulous quality, almost like the sound of a saw. At the angle of scapula a high-pitched ringing crepitation was heard towards the end of inspiration. The stridulous quality of breath-sound was heard over the right front of the chest, and also faintly on the left side, as if by conduction. Everywhere in front the breath-sound was accompanied by a large metallic bubbling. Pulse excessively small, feeble, and rapid; temperature at 8 a.m. 104.2 ; at 6 p.m. 104.6° . The boy gradually sank, and died on May 5.

On examination of the body the bronchial glands were found to be much enlarged; they were soft in consistence, and were not at all cheesy. The right lung was adherent over the upper half to the chest-wall, but the adhesions were not of long standing, and were broken down without difficulty. The lung itself was extensively indurated. The upper lobe still crepitated slightly on pressure, but the two lower lobes were very dense and solid. On section of the lung, white thick bands of fibroid tissue were seen radiating from the root of the lung towards the circumference. The bands were separated by purplish hollows or cavities, which on close examination was seen in each case to be continuous in the direct line with bronchial tubes. They were for the most part perfectly smooth in the interior, and were evidently dilatations of the bronchi. Here and there, however, some ulcerations in the walls were discovered, and one or two of the cavities appeared to be the result of ulcerative

destruction of tissue. The larger bronchi were very much thickened as well as dilated. In the case of the smaller air-tubes the walls were not especially thickened, but the tissue between them was very dense, and appeared to consist of fibroid tissue interspersed with emphysematous air-cells. This dense tissue was honey-combed in all directions by the dilated bronchi, which ended in spacious *culs-de-sac* almost immediately under the pleura. The left lung was extensively emphysematous, and presented large patches of collapsed tissue. It was the seat of bronchitis, otherwise its structure appeared to be unaltered. No cheesy deposits were found in either lung, nor were there any grey granulations either there or in any other organs. The liver was enlarged, reaching to level of umbilicus; its substance was dense, its edge firm and sharp, and its section was stained by a watery solution of iodine, showing the presence of amyloid degeneration. The spleen was slightly enlarged, and had a waxy look.

This boy had suffered for three years and a half from cough, with the expectoration of yellow phlegm. The attack, however, for which he was first admitted into the hospital had begun only a week before, with pain in the side. On examination there was found to be general dulness on percussion over the right half of the chest, both before and behind. As has been often before remarked in the course of these pages, dulness coming on suddenly, and occupying the front as well as the back of the chest, is due, in the large majority of cases, to pleurisy with effusion; but, in the present instance, the other physical signs failed to support this view. The loud cavernous breathing and bronchophony were indeed, in a child, quite compatible with the existence of even a large effusion, but the presence of vocal fremitus, which was throughout stronger on the diseased than on the opposite side; the character of the percussion-note, which was not completely toneless, and was not accompanied by much increase of resistance; the non-

displacement of the heart, and the shape of the chest, which, as drawn by the cyrtometer, had not the square outline which it invariably assumes when the fluid is at all considerable in amount,—seemed to imply that if pleurisy were present, it was accompanied by little effusion, and that the dulness must therefore be principally due to another cause. Besides, the abundant fine bubbling rhonchus heard all over the affected side of the chest was evidently intra-pulmonary, and was very different in character from the superficial pseudo-crepitation of an inflamed pleura.

The physical signs, therefore, taken altogether, pointed to consolidation of the lung, and tended directly to disprove the theory of liquid effusion. The next question, then, was to determine the nature of the consolidation. The history showed the existence of a lung affection dating from an attack of pertussis three years and a half before. Super-added to this was some recent acute attack, which was accompanied by a considerable elevation of temperature. The cavernous breathing, large metallic gurgling, and bronchophony indicated the presence of cavities of the lung ; but were these cavities the result of breaking down of lung-tissue, or were they merely dilatations of the bronchi ? The well-nourished state of the patient seemed to imply that excavation so extensive could not in this case be the consequence of ulcerative destruction of lung, for so serious a lesion must necessarily have been attended by a very considerable degree of emaciation. Besides, there was no evidence of deposits in the apex of the opposite lung. The signs in question must therefore have resulted from the presence of dilatations in the bronchi. Bronchiectasis may be either an acute or a chronic condition. It is acutely produced in many cases of catarrhal pneumonia, more especially in those where the disease assumes a subacute form. The dilatations are seldom noticed until the disease has lasted two or three weeks ; and after the subsidence of the inflammatory attack the air-tubes may gradually resume

their normal calibre. Sometimes, however, and usually in cases where the lung has been the seat of repeated attacks of catarrhal pneumonia, the bronchi become permanently dilated. There is then always considerable thickening of the air-tubes, and the fibrous tissue of the lungs is in great excess. Such cases are, in fact, cases of fibroid induration of the lung. In the present instance the disease had lasted too short a time to give rise to acute bronchiectasis; and therefore we must conclude that the right lung had suffered from several attacks of catarrhal pneumonia, which had led to fibroid induration of the lung-tissue and permanent dilatation of the bronchi. The present was evidently a similar attack, and during its progress further dilatation of the air-tubes appears to have taken place; for the respiration, which at first had been cavernous, became completely amphoric, and the bubbling rhonchus assumed an almost musical tone, approaching somewhat to the tinkling sounds generated in a case of pneumothorax. This extreme dilatation was, however, only temporary. As the inflammation subsided, the bronchi contracted again to a certain extent except at the base of the chest, where cavernous breathing appeared to be a constant phenomenon.

When the child left the hospital on April 10 there was no contraction of the side; but on his readmission, after an interval of six months, this was noticed to a very slight extent. The physical signs on his second visit were little altered, with the exception that the dulness was perhaps more extensive at the right base than was noted at the last report before his discharge, and that the vocal vibration of the chest-wall, which had before been strong, could now scarcely be detected. The disappearance of vocal fremitus over the seat of dulness is not an uncommon feature in fibroid induration, and is, indeed, the rule in cases where the disease is extensive. Sometimes, however, vibration of the chest-wall appears to vary according to the amount of fluid retained in the tubes. Where this is large, vibration

is suppressed, but when the air-tubes have been recently emptied, the fremitus becomes more perceptible. There was in this boy very little contraction of the chest-wall. This may be, no doubt, accounted for by the great dilatation of the bronchi; for when the account of bronchiectasis is sufficient to compensate for the loss of space by contraction of the lung, the chest-wall is prevented from falling in. The lung itself, however, was much diminished in size, as was shown by the resonance of the left lung passing across the middle line of the chest in front, and reaching, indeed, as far as half an inch to the right of the edge of the sternum. The pyrexia, which continued all through the child's second residence at the hospital, was symptomatic of a certain amount of subacute catarrhal pneumonia, kept up probably by the continued presence in the air-tubes of putrescent pus, which acted as a source of constant irritation.

Shortly before his final admission to the hospital the child caught a further cold and had a severe attack of bronchitis, under which he quickly died. Post-mortem examination showed, as was expected, a very marked degree of fibroid induration of the lung, with extreme and extensive bronchiectasis.

In the following case extensive fibroid induration was combined with a large excavation of the lung :—

CASE XXXII.—FIBROID INDURATION WITH EXTENSIVE EXCAVATION OF LUNG (FIBROID PHTHISIS).

William W., aged twelve, was admitted into the East London Children's Hospital on March 25. The boy had been fat and healthy up to the age of two years. He then caught a severe cold and began to lose flesh. The cough continued for some time afterwards. He had whooping-cough when three years old, and measles at the age of five.

Since that time he had been troubled with a frequent cough and often expectorated phlegm. His breathing also since the attack of measles had been habitually short. For five months before admission his urine had been noticed to be purulent, but for several months previously he had cried with pain whenever he passed water.

State on admission:—Patient a very thin boy; superficial veins very full and prominent along sides of neck and arms, and a network of veins seen on right side of chest close to sternum. The veins of neck swell up very much when boy coughs. Lymphatic glands, particularly those under the jaw, enlarged and hard, but not tender. Some flattening under right clavicle; right shoulder not lower than left. Heart's apex in normal situation, but the impulse is also plainly seen in the epigastrium and in the second and third interspaces immediately to right of sternum. Very little movement of either side of chest during inspiration. Abdomen is full, and liver reaches to level of umbilicus. Edge of spleen cannot be felt, but there is an abnormal sense of resistance below false ribs on left side. There is no œdema of the limbs. The percussion-note is wooden over the right front from above clavicle to nipple level, and at back from the apex to the angle of scapula. At the base both in front and behind the percussion-note is comparatively dull. Vocal fremitus is stronger at the right than the left side. Respiration is cavernous above, loud and blowing below, and over the dull area at the base very weak and blowing. Vocal resonance is everywhere bronchophonic, and at the base can be heard as a faint distant echo. With the exception of some dulness at the extreme base resonance is normal at the left back, but respiration all over the left side is loud and coarse, and is rather hollow at the left edge of the sternum.

The boy passes water tolerably freely: the urine is alkaline, sp. gr. 1010, and contains abundance of pus. There is some tenderness on pressure over the pubes, and

the boy complains of pain in the region of the bladder immediately before micturition.

For some days after admission the temperature was about 100° at night, but fell in the morning to the natural level. The pulse was slightly over 100, and the respiration 30 to 36. On April 5 the temperature rose in the evening to 101.4° , and was 100° on the following morning. During this time much coarse crackling rhonchus was noticed over the whole of the right side both back and front, except the posterior base, and a soft systolic murmur was noted to the right of the sternum over the seat of visible impulse.

From this time the boy became gradually weaker and thinner. His temperature varied, sometimes rising as high as 104° in the evening, and seldom falling lower than 101.8° after 6 p.m. In the morning it stood usually at 100° . On May 2 some ringing metallic bubbles were heard with respiration on the right side, mixed with cracking sounds. The urine remained loaded with pus. On May 20 the temperature fell to 98.6° in the evening, and remained low till May 27, when the boy died. At 8 a.m. on the morning of his death the temperature was 97° .

During his residence in the hospital the patient complained frequently of digestive disturbance. The bowels were often relaxed, and the tongue was always red and glazed-looking as if entirely denuded of epithelium. On admission his weight was 3 st. 4 oz. By April 12 he had gained 4 oz., but soon afterwards he began to lose flesh, and on May 18, when his weight was taken for the last time, it was only 2 st. 9 lbs. $3\frac{1}{2}$ oz.

The physical signs remained unaltered throughout, except for variation in the amount of rhonchus which was heard with the breath-sounds. This at times would be copious, very metallic and ringing, at other times scanty, and mixed up with creaking sounds. His breathing, although rather hurried, was not distressingly so, and he never complained of dyspnoea. He expectorated a good deal of purulent

matter at times, and the cough was sometimes very violent.

On examination of the body the liver was found to be very large, weighing $4\frac{3}{4}$ lbs. The spleen weighed $5\frac{1}{2}$ oz., and the kidneys $5\frac{1}{4}$ oz. each. All these organs were undergoing amyloid degeneration, and their sections were deeply stained with iodine. The right kidney was the seat of old pyelitis : the walls of the pelvis were very thick. In the substance of the organ was an abscess with thickened walls communicating with the upper part of the pelvis. Both ureters were dilated and hypertrophied. The walls of the right were $\frac{1}{8}$ in. in thickness. The bladder was much contracted, with thickened walls : the mucous membrane was granular, and felt very dense. No ridges of muscular fibre were seen as in cases of hypertrophy from stricture of the urethra.

The right lung was found to be forced upwards and backwards against the spine, so that the upper surface of the diaphragm reached as high as the third rib. The lung was everywhere adherent, and was much torn in removal. On section its substance was excessively hard, creaking under the knife ; the walls of the bronchi were thick and rigid, and throughout, the fibroid tissue of the lung was in such great excess that very little air-containing tissue appeared to be left. A large cavity was found occupying the antero-lateral part of the lung, and extending nearly from apex to base : many large bronchi opened into it. There was a small collection of purulent matter at the base of the lung between the opposed surfaces of the pleura. The left lung was emphysematous at the upper part with some collapse at the posterior base. On pressure some small nodular masses were felt in the interior, and on section the upper lobe and part of the lower were found to contain numerous small cheesy masses. In the upper part of the lower lobe at its posterior surface was a cavity the size of a filbert filled with thick purulent matter. The heart lay in its natural

position, being kept in place by adhesions between the pericardium and pleura. The right ventricle was somewhat thickened. The pulsation noticed during life to the right of the sternum was produced by the contraction of the right auricle.

There can be little doubt that in this case the origin of the disease was an attack of pleurisy on the right side of the chest. The lung had been apparently exposed for a considerable time to severe pressure from effused fluid, and had not afterwards expanded. The fluid had been gradually absorbed, except a small quantity which was still left separating the layers of the pleura at the base: elsewhere the opposed surfaces had formed firm adhesions. Except a certain amount of flattening under the clavicle there was little contraction of the right side of the chest, for the drawing upwards of the liver and diaphragm prevented the wall from falling in.

The fibroid induration of the lung took place, no doubt, as a consequence of the pleurisy, the fibroid overgrowth beginning at the circumference and spreading thence into the interior, until the greater part of the organ was converted into a fibroid hard mass. The large cavity discovered in the interior of the lung was in all probability a consequence of the fibroid induration. Ulceration of bronchi is a not uncommon result of such a lesion, and may lead, as in this case, to extensive excavation. The comparatively small amount of disease in the left lung seems to support this view, for had the ulcerative destruction of lung preceded the fibroid change, we should have expected much more extensive consolidation on the opposite side of the chest.

The chronic cystitis and pyelitis probably resulted from neglected catarrh, and the profuse purulent discharge to which they gave rise was quite sufficient to determine amyloid degeneration of organs, especially in a young subject. Children, as has been before remarked, have a special tendency to this form of degeneration, and in them a compa-

ratively small amount of suppurative drain is sufficient to excite it.

After the illustrations furnished by the preceding cases, little remains to be added upon the subject of diagnosis. The disease produces consolidation of the lung, combined with contraction of the diseased pulmonary tissue and dilatation of its bronchi. Its distinguishing marks will therefore be the physical signs indicative of those conditions : a falling-in of the chest-wall at the seat of disease, with drawing of the neighbouring organs towards the affected part ; dulness on percussion, with usually diminution in vocal fremitus ; loud blowing or cavernous breathing, with metallic crepitation and gurgling, and bronchophonic resonance of the voice ;—these signs confined to one side of the chest ; the other lung being healthy. Additional evidence is furnished by the character of the special symptoms. The cough is paroxysmal, and ends in the expectoration of large quantities of offensive purulent fluid, sometimes also in retching and sickness. The breath often has an offensive putrid odour. There are signs that the blood passes with difficulty through the lungs : the right ventricle is frequently hypertrophied ; the superficial veins of the chest, neck, and often of the arms, are unusually visible ; and the fingers soon become clubbed.

If the case be first seen during an attack of catarrhal pneumonia, it may be mistaken for one of pleurisy with effusion ; if there be much falling-in of the chest-wall, the disease may be confounded with pleurisy in the stage of retraction. The distinguishing marks, by means of which the two diseases may be recognised, have already been insisted upon (see page 215). Of these, the one which may be looked upon as of the chiefest importance in excluding pleurisy is the presence of metallic gurgling rhonchus ; for although, in the pleurisy of childhood, cavernous breathing and bronchophony are far from uncommon, the breath-sound is not

accompanied by rhonchus, and the crackling friction which is often heard can usually be readily recognised by its superficial character, and is never large and metallic like the sounds produced in a rigid, dilated air-tube. The cavernous character of the breath-sounds may cause the presence of ulcerative destruction of lung to be suspected, but the strict limitation of the disease to one side of the chest, which would not be the case in phthisis, may serve to establish the difference. Moreover, the general condition of the patient is here of importance. In fibroid induration of the lung nutrition is usually fair and the strength pretty good, while the third stage of phthisis is seldom unaccompanied by emaciation and great weakness. It must not, however, be forgotten that ulceration of the bronchial tubes may occur as a consequence of the continued irritation induced by the fetid contents of the tubes, and may lead to destruction of lung-tissue and the formation of cavities. In these cases (called "fibroid phthisis," by Sir Andrew Clark), cavities and dilated bronchi may be present in the same lung. Here, however, the opposite lung seldom escapes, and, if there be much destruction of tissue, general nutrition is seriously impaired. The child wastes and becomes more and more enfeebled; his temperature is high; and he dies with all the symptoms of consumptive disease. In most cases where ulceration of the bronchial tubes is taking place, a careful microscopic examination of the sputum will discover the presence of areolar fragments of elastic tissue.

The *prognosis* in fibroid induration of the lung in children is not necessarily unfavourable. So long as it remains simple it appears to exercise little influence upon the nutrition and general development of the child. It becomes dangerous only through the accidents which such a condition of lung tends especially to provoke. As such patients are very sensitive to changes of temperature, they are very prone to suffer from attacks of pulmonary catarrh :

and on account of the difficulty met with in expelling secretion from the rigid air-tubes, the sputum is apt to be retained and to putrefy. It then gives rise to great irritation, and after a time may excite an attack of catarrhal pneumonia. The seat of the induration is therefore of importance in prognosis, for if the apex of the lung be the part affected, the force of gravity will itself tend to prevent accumulation of secretion in the air-passages, and the chances of retention will be greatly diminished ; while in the case of disease at the base of the lung, all the conditions assist in opposing the escape of the contents of the tubes, and the danger of an attack of catarrhal pneumonia is materially increased. As the irritation of the bronchi is in direct proportion to the degree of putrescence of the expectoration, an offensive smell from the breath or the sputa is a certain sign that the lung is exposed to the risk of inflammation.

Catarrhal pneumonia, when it occurs, generally assumes the subacute, and therefore less severe type, and if the disease be not extensive and the nutrition of the child be good, often ends favourably after a time ; but its course is apt to be protracted, for the retained decomposing secretions act as a constant source of irritation. When the attack of pneumonia is acute and violent, the issue is usually as disastrous as in any other case of this serious disease.

Anything which reduces the strength of the child, such as attacks of diarrhoea, or of vomiting, or of severe hæmoptysis, must be looked upon as an unpromising symptom ; for healthy nutrition of the body generally is indispensable to favourable progress. Hæmoptysis is not necessarily a sign that ulceration of the bronchial tubes is taking place ; but it is often due to this cause, and when it occurs, a careful microscopic examination of the expectoration should be made. If amyloid disease of organs occur, especially if it be accompanied by œdema, the prognosis becomes very

serious, but even in these cases death may not take place until after considerable delay. Œdema, although unaccompanied by amyloid change, is itself a dangerous symptom, as it betokens a very impoverished condition of the blood.

The treatment of fibroid induration of the lung consists in doing our best to maintain healthy nutrition, and in preventing, if possible, the accidents to which this state of lung is necessarily prone. On account of the sensitiveness of such patients to changes of temperature, great care should be taken to avoid sudden chills. The child should be warmly dressed, and should never leave the house in damp or cold weather without some special protection for the neck and chest. All catarrhs should receive immediate treatment. The patient should be at once confined to bed, and saline diaphoretics should be administered so as to loosen the cough and relieve the irritation of the air-passages. Afterwards, stimulating expectorants can be employed to assist in the expulsion of the phlegm.

In cases where there is great rigidity of the bronchi, and little power to expectorate, and especially in cases where an offensive smell from the breath indicates that secretion has been retained and is putrefying in the tubes, disinfecting and astringent inhalations should be employed to remove the offensive character of the secretion and prevent its reproduction. Inhalations of steam impregnated with carbolic acid or creasote (twenty minims to the pint), may be made use of, and the addition of a teaspoonful of the compound tincture of benzoin, which is a powerful stimulating expectorant, will be found of service. When the amount of secretion is large, it can often be controlled by limiting the quantity of liquid taken by the patient, and by the use of tannin and other astringents in a spray. In these cases, the best medicines are tonic and astringent remedies, such as tinct. ferri perchlor. in large doses, with quinine, and a few drops of liq. morphiæ. Cod-liver oil is also useful. The expulsion of the contents of the tubes is greatly assisted

by an occasional emetic. The best emetic in these cases is the one which acts most quickly and produces the least nausea, such as a teaspoonful of mustard in water, or a couple of drachms of powdered alum in syrup. Sulphate of zinc is not to be recommended for children, as its action is very uncertain.

In cases where vomiting takes place after the fits of coughing, great interference with nutrition may be the consequence, for the paroxysms of cough are apt to occur immediately after a meal; and in such a case the greater part of the food may be returned. Now, the vomiting cannot be regarded as otherwise than a beneficial act, in that it assists in the expulsion of the injurious secretions. It is only hurtful to nutrition because it takes place at an inconvenient time. In these cases the terminal vomiting may often be controlled by small doses of liq. strychniæ, or by one drop of liq. arsenicalis given with iron and morphia; but as it is undesirable to put an end to the vomiting entirely, this may be encouraged at another time, as in the early morning, by draughts of warm water, preceded, if necessary, by a dose of alum or mustard and water.

Catarrhal pneumonia, when it occurs in these cases, must be treated in the usual way. When there is dropsy, tonic and astringent remedies are especially indicated.

CHAPTER VIII.

ACUTE GENERAL TUBERCULOSIS.

ACUTE tuberculosis a common and fatal disease in children—not to be confounded with pulmonary phthisis—resembles in many of its characters the acute specific diseases.

Pathology and causation.

Symptoms—General—Special—Tuberculosis of lung—Excites bronchitis and catarrhal pneumonia.

Case XXXIII.—Acute pulmonary tuberculosis with tubercular peritonitis—Comment.

Case XXXIV.—Acute general tuberculosis following old-standing disease of bone—Comment.

Diagnosis.

ACUTE tuberculosis is a very common disease in children of every age, and, as one of their most fatal disorders, requires especial study, more particularly as it is often very difficult of diagnosis.

By acute tuberculosis we mean an acute general affection giving rise at first to but few symptoms, and therefore difficult to identify, but of which the invariable anatomical expression and distinguishing mark is the grey granulation. The disease is not to be confounded with pulmonary phthisis (ulcerative destruction of lung), nor with bronchial and mesenteric glandular enlargement. It may be combined with these diseases—and, indeed, in the lung the grey granulation is often found secondary to pulmonary mischief,—but it may also arise quite independently of such affections, and, as a distinct disorder, with distinct pathological characters, cannot be said to hold any necessary relation to the so-called consumptive diseases. So far from being allied in its nature with the phthisical disorders,

acute general tuberculosis seems to partake more of the character of an acute specific disease, and in the insidious manner of its occurrence and the severity of its general symptoms may be compared to enteric fever, to which disorder, indeed, it often exhibits a very close analogy. It resembles the acute specific diseases also in the frequency with which it comes on unexpectedly in children of apparently the most robust constitution. It is not only weakly, pining infants, and thin, pallid, delicate-chested children who fall victims to this disease. These may, and frequently do, become the subjects of acute tuberculosis, but it is often in such cases a secondary complication—an intercurrent affection supervening upon already existing mischief in the lung. On the contrary, when it occurs as a primary disorder, it may seize upon children who are in apparently the soundest health, and in whom no signs have ever been noticed to indicate the likelihood of such an attack. This is especially the case when the tubercular outbreak falls principally upon the meninges.

Even in these cases, however, it seems probable that the tubercular affection is sometimes a secondary disease ; that it is a result of infection of the system from some previously unsuspected cheesy deposit. It is far from uncommon in the post-mortem examination of such cases to find old caseous masses in the body ; and many pathologists of eminence are now disposed to believe that such masses are capable of giving rise to a general infection of the system, even although they may themselves have induced no symptoms at all, or symptoms so slight as to have attracted no especial attention. According to this view, therefore, acute tuberculosis, although it is to be distinguished from pulmonary phthisis, may yet be directly excited by this condition of the lung. Caseous pulmonary deposits, as they soften and break down, may become direct sources of infection of the system, giving rise to the tubercular outbreak. So, also, cheesy enlargement of the lymphatic glands may be

followed by the same result. In this way may be explained the many cases of acute tuberculosis supervening upon measles or whooping-cough: the bronchial glands, which had become enlarged during the progress of these diseases, instead of returning to a healthy state, undergo a cheesy transformation, and after a time soften and lead to a general formation of tubercle* in the body, although the child may have appeared in the interval to recover completely from the original complaint. So, carious disease of bone of long standing may determine the occurrence of acute tuberculosis in children who have never manifested any signs of tubercular predisposition.

Acute tuberculosis may, therefore, be induced as a secondary affection in children who had previously presented but slight signs of ill-health, and in whom no family tendency to tubercular or other phthisical disease can be detected. It may, however, also occur as a primary disease in cases where the presence of a softening cheesy deposit cannot be suspected, and where, indeed, actual examination of the body has revealed no cause to which the outbreak of the tubercular disorder can be attributed. In such cases there is usually evidence of hereditary predisposition either on the side of the father or of the mother, or of both ; and we often find on inquiry that the case in question is not a solitary instance of the disease occurring in the family, but that some of the other children had been previously seized more or less suddenly with severe symptoms about the head, the chest, or the belly, and had rapidly died, with all the signs of acute tubercular disease. In many cases the predisposition extends to the exact form in which the

* The use of the word "tubercle" is here confined to the grey-granulation. The term "*grey granulation*" is used in deference to existing custom, although in children—and often indeed in adults—the nodules quickly lose their grey appearance and become yellow—so quickly in some cases that it is difficult to say that the colour was not yellow from the very first.

tubercular disorder is to manifest itself, and a family may lose several children in succession from tubercular meningitis.

When the hereditary tendency is thus strongly marked, there is often in the build of the child's body an approach to the so-called tubercular type of physical conformation. The chest, indeed, may be perfectly formed, and may show no signs of antero-posterior flattening, but in the clear complexion, the thin transparent skin, the large bright eyes and red lips, the small bones, the sensitive nervous system, and the general delicacy of organisation, we find evidences of a distinct constitutional condition in which the development of acute grey tubercle may be excited by comparatively trifling causes. This is not, however, always the case, for children may equally become the subjects of this complaint, who have never presented any such peculiarities, nor indeed ever exhibited in themselves any indications of a tendency to tubercular disease, although such indications may, perhaps, be found in the family history.

The cause of the tubercular outbreak must therefore in many cases remain unknown. Bad hygienic conditions, which are usually quoted as causes of this disease, may be capable of exciting a dormant tendency where this is strong, and may, by nourishing a propensity to caseous degenerations, tend indirectly to excite the formation of true grey tubercle ; but when the disease is found in children previously healthy, and reared in strict obedience to the laws of health, there must be other causes in operation of which we know nothing, and against which, therefore, it is impossible to guard the patient.

Acute tuberculosis is liable to occur after certain of the acute specific diseases, especially measles, typhoid fever, and variola. Whooping-cough is also sometimes followed by it. Indeed, where the predisposition is strong, a mild febrile attack or a slight shock to the system may produce sufficient impairment of nutrition to lead to the outbreak of the

disease. Thus, tuberculosis may come on after varicella, or may be set up by an accidental injury, such as a bad fall.

Acute tuberculosis is marked by the dissemination through the organs of the grey granulation. In a clinical work it is unnecessary to enter into strictly anatomical details as to the structure, etc., of a body with the appearance of which everyone must be familiar. It is, however, of practical importance to remember that the grey granulation is an outgrowth from the lymphatic system, and that one of its principal points of origin is the adenoid tissue which forms the lymphatic sheath round the minute arteries in many situations. This connexion with the blood-vessels may lead to great interference with the circulation, and produce extensive disorganisation. Thus in the pia mater the pressure of the perivascular growth causes great hyperæmia of vessels, followed by basilar meningitis, infiltration of the tissues with serous fluid, and softening of the substance of the brain. In the lungs its seat is not only in the lymphatic sheath, but also in the connective tissue between the lobules and the infundibula. Here the presence of the new formation causes a thickening of the alveolar walls, which may eventually lead to obliteration of the cavity of the cell. It also before long sets up catarrhal inflammation in its immediate neighbourhood, so that the larger nodules become surrounded by a circumscribed zone of catarrhal pneumonia. This, however, is a later change, and one which is not always to be seen, for, on account of the comparatively short course of acute tuberculosis, and the rapidity with which the disease hurries to a conclusion, death may take place before any inflammatory process has been excited in the tissue by the presence of the nodules.

The grey granulation may be found in all parts of the body—wherever, in fact, adenoid tissue exists. Its distribution, however, is not always general. It may be so, but often its presence is limited to special cavities or organs. The child, however, differs from the adult in one respect—

viz., that whereas in the latter, whenever tubercle exists in the body, the lungs rarely, if ever, escape, in a child dying of this disease the lungs are often found to contain no grey granulations at all, although the nodules may be plentifully distributed over other organs.

The symptoms of acute tuberculosis are difficult to describe, on account of the great variety of the shapes in which the disorder may manifest itself. As a constitutional disease affecting the body generally, it may give rise to general symptoms denoting the distress into which the whole system is thrown; and as a local disease affecting particularly special organs, it may excite special symptoms arising from the particular organ which is the principal seat of pathological change.

The general symptoms almost invariably appear before the local, and on this account they are often spoken of as *premonitory* symptoms, ushering in the more distinctive signs of disease of organs. When these latter occur, their intensity is usually so great as completely to overpower the symptoms by which they had been preceded. Thus, the symptoms of tubercular meningitis seldom come on suddenly in a healthy child, but are introduced by several weeks of obscure malaise which presents little that is characteristic, and merely hints vaguely at the presence of disease without offering any distinctive signs by which the nature of the complaint can be recognised. This malaise is the expression of a general as distinguished from a local disease, and denotes the influence of the constitutional disorder upon the entire system. The collection of symptoms more or less definite of which it is composed are the outward manifestation of the general disease—acute tuberculosis. The general symptoms consist of little more than pallor, languor, and fever, with gradual loss of flesh. The first change noticed is usually in the complexion of the child: his face loses its healthy tint, and becomes pecu-

liarly pallid and transparent. In the morning this is especially noticeable ; towards evening there is usually a bright flush on the cheek-bones, contrasting markedly with the intense pallor which still exists round the nose, mouth, and chin. Almost at the same time the child becomes languid and often irritable ; he looks wearied, complains of feeling tired, and lies about instead of amusing himself with the usual sports of his age. The wearied look of the child is an important symptom at this early stage. Although nothing very definite can be discovered in his state, and his organs give no sign of suffering, the haggard look in the patient's face and the pallor of his complexion show him to be really ill. That he should be so with so little to explain his indisposition is a noteworthy fact, and one tending in itself to turn our suspicions into a right direction.

The temperature is a very important symptom ; it rises in the evening to 100° or 101° , seldom higher,* and generally falls again to a certain extent towards the morning. This increase of temperature is not always readily estimated by the touch. The palms of the hands and the forehead often feel dry and hot ; but the general surface does not necessarily present to the hand of the observer any marked symptom of fever. The thirst, however, is usually great, the appetite poor, and the pulse quick and irritable. The fever sometimes takes on a remittent character, and children who are old enough to describe their sensations will complain of feeling cold at certain, often regular, intervals, and may even shiver ; but it is rare for actual rigors to take place. The feeling of chilliness may be merely a subjective symptom, or may be accompanied by an actual diminution of the temperature of the body. The regularity with which it occurs is often suggestive of ague, but the stages of cold, heat, and sweating do not follow one another with any

* An evening temperature higher than 101° generally indicates the presence of an inflammatory complication.

uniformity, and present many peculiarities which distinguish the case from one of intermittent fever. The chilliness begins generally in the evening, and occurs at about the same time every night. Shivering is very rare, and sweating is often absent. In the interval the state of the patient is very far from being satisfactory: it is not an intermission, but a remission of the fever, and the temperature is not normal between the attacks.

Loss of flesh soon follows the pyrexia, but the symptom is one which varies greatly in intensity. The emaciation may be rapid or slow, and sometimes, especially in children at the breast, proceeds so gradually as to cast doubts on the accuracy of the diagnosis. If there be any diarrhœa the wasting is more rapid. The bowels, indeed, are very often irritable early in the disease, but the looseness is apt to alternate with constipation.

In infants a common symptom is loss of elasticity of the skin. This change is best seen in the skin of the abdomen, which, when pinched up, does not recover its smoothness as quickly as a healthy skin would do. In extreme cases it remains wrinkled in small folds. This state of the skin is probably dependent upon imperfect action of the kidneys.

The urine often contains a little albumen, and occasionally a certain amount of œdema may be present in the hands and feet, and even in the face. Œdema is a symptom which may be produced by other diseases whose tendency is to cause much impoverishment of the blood, such as simple chronic non-tubercular diarrhœa; but when combined with a moderately elevated temperature it is very suspicious of tuberculosis.

In the above collection of symptoms there is nothing which marks out the disease with any distinctness so that it can be immediately recognised. The beginning is very gradual as a rule, and nutrition is interfered with but slowly. When fully developed the symptoms are merely those of fever, and indicate some general disturbance of the

system; hence, until local signs arise to draw attention to some particular organ, the nature of the disease must often remain—at any rate, for a considerable time—a matter of doubt. On account of the frequency with which a functional derangement of the bowels is present early in the disease, there is an especial danger that the case may be mistaken for typhoid fever, to which disorder the symptoms under such circumstances may present the very closest resemblance. In rare instances there is a mode of beginning which is almost identical with continued fever. The disease breaks out with more abruptness; there is headache, bleeding from the nose, and the bowels are relaxed or confined; the child is very feverish, restless, and stupid. Such cases are uncommon. In one which came under my own notice there was a peculiar dusky tint of the complexion, and after some days a strange, dull mottling of the skin was observed, which gave the case a close resemblance to true typhus.

The special symptoms arise from the head, the chest, and the abdomen; but these cavities are seldom equally affected, and usually one or another of them give rise to evidences of distress, showing that the contained organs have become the especial seat of the disease. In a description of acute tuberculosis it is convenient, for the sake of clearness, to take each cavity separately, but it must be remembered that the special symptoms are not necessarily limited to one group of organs—on the contrary, they are variously combined; and the same case may present at one time symptoms from each of the cavities of the body.

Most commonly it is the lungs which first give rise to special symptoms. In almost all cases the disease has not proceeded far before a short hacking cough begins to be noticed, indicating a certain irritation of the air-passages. The cough in the beginning is infrequent and dry, and an examination of the chest reveals no signs of pulmonary mischief. After a time, however, the cough is heard more

frequently, but still there is no dulness, and a stethoscopic examination rarely discovers anything but weak, harsh breathing, with a little scanty sibilant rhonchus scattered here and there about the lungs. This absence of physical signs is an important symptom in such cases; for if the cough be frequent and dry, the breathing hurried, and the temperature elevated, the absence of distinct physical signs should lead us to entertain strong suspicions of the tuberculous nature of the disease. The breathing is not, however, necessarily much quickened. There may be little increase in the number of respirations throughout the whole course of the disease—so long, at least, as the physical signs remain insignificant. If, however, as generally happens, a secondary bronchitis be set up, or pneumonic changes occur, the breathing may become hurried, and in such a case the physical signs would indicate the nature of the secondary lesion. It often becomes a nice question, when the patient is seen for the first time under such circumstances, to decide upon the tuberculous or non-tuberculous origin of these affections.

The most characteristic form, perhaps, is that in which a secondary bronchitis is induced by the presenee of the grey granulation in the air-passages. Here the breathing is rapid, and the nares act in respiration; the intercostal spaces sink in deeply; there is some lividity of the face, and the countenance indicates considerable distress. The pulse is small and rapid, and the temperature rises to between 102° and 103° . On auscultation of the chest, we find the breath-sounds covered by crisp medium-sized bubbles, which have a certain ringing quality, and occupy the whole extent of both inspiration and expiration. This rhonchus is heard at all parts of the chest, although it may be most abundant at the upper parts of the lungs. In other respects the physical signs are normal; the resonance on percussion is perfect, and the vocal resonance and the natural vibration of the chest-walls are unaltered. Some-

times in such cases, when the patient is very young, local collapses may occur, especially at the apices of the lungs. The percussion note then becomes slightly dull, and faint bronchial breathing is detected over the seat of the lesion. These signs occur with very great irregularity, being present one day, to disappear the next, and seldom persist longer than two or three days in succession. Often, however, a permanent dulness may be found at one or more spots; the respiration is then louder and bronchial or blowing, often completely tubular, and the rhonchus, where these signs occur, becomes drier in quality, and more distinctly crepitating. The temperature at the same time rises to 103° or higher, the lividity increases, the breathing becomes distinctly laborious, and there is much recession of the soft parts of the chest, and of the lower ribs, during inspiration. These symptoms are due to the occurrence of catarrhal pneumonia, and foretell a rapid ending to the disease. The lividity and distress increase, the respiration and pulse become more and more hurried, the emaciation proceeds with rapidity, and the child dies exhausted. Death is often preceded by convulsions, followed by unequal pupils, strabismus, paralytic lesions and other signs indicating implication of the cerebral meninges.

CASE XXXIII.—ACUTE PULMONARY TUBERCULOSIS WITH
ACUTE TUBERCULAR PERITONITIS.

The following case presents a good illustration of the pulmonary form of acute tuberculosis :—

Maria H., aged three years and six months, a child born of healthy parents in a family without consumptive history, was healthy up to the second week in October, when she had an attack of measles. After the disappearance of the eruption, the cough continued, her abdomen began to get large, and she was noticed to be wasting. For a month her bowels were relaxed, but they afterwards became more regular. Her appetite, however, was poor, and after eating

she often vomited the food she had just swallowed. All the time the child kept her bed; she was feverish, had a loose cough, and grew thinner and thinner.

On December 15, being still no better, she was brought to the East London Children's Hospital. Temperature on evening of admission 102° .

Child was small-boned and rather delicate-looking, with small features. Expression anxious. Lips dry and red. Tongue pink and moist. Abdomen rather full; liver projected for about two inches below edge of ribs, and the spleen was felt for about the same distance on the left side. There was some tension of the abdominal parietes, but tenderness was not marked, and nothing special was noted about the belly. Movement of chest in breathing normal, and intercostal spaces sank in. At the right base there was dulness, which reached upwards in front to the level of the nipple, and behind to within a finger's breadth of the angle of the scapula. Over the dull area the respiration was excessively weak and rather harsh. Elsewhere there was no dulness, but the respiration was very loud and coarse, and much clicking rhonchus was heard all over both sides of the chest. The dulness at the right base behind passed rather gradually into resonance above, without abrupt transition. For this reason, and also because the dulness reached upwards higher in front than behind, it was considered (and rightly, as the event proved) to be due to an enlarged liver.

After this report the child became gradually worse. The temperature remained high, varying from 101° to 103° in the evening. The pulse was quick, and the breathing hurried. The bowels continued rather loose, being open three, four, or more times in the day.

On December 19 the percussion-note was found to be a little higher-pitched at the right supra-spinous fossa than at the left, and the breathing there was harsher. These signs became more marked after a few days, when dulness

was noticed also above the right clavicle in front, and the respiration there and at the supra-spinous fossa became distinctly bronchial. On December 30 the physical signs about the chest remained the same, except that the clicking rhonchus previously noted had given place to a sibilant wheeze at the end of inspiration, and this was heard all over the chest, both at the back and at the front. The general symptoms all this time were little altered. The abdomen looked full to the eye, but there was no tenderness of the part on pressure, nor was the resistance of the abdominal wall increased. At no time did the patient show signs of pain in the belly. The child was weaker and thinner, and was very fretful, but she took her food well, and the looseness of the bowels had subsided; temperature at 8 a.m. 100° , 6 p.m. 103° .

After this date the emaciation proceeded more rapidly. The cough got more troublesome and hacking. The bowels were opened frequently, but the motions were very small, and were not loose.

On January 6 a crisp medium-sized bubbling rhonchus was heard all over the chest both at the back and at the front, but it was nowhere very copious. The percussion-note at each apex was high-pitched, with harsh or faintly bronchial breathing. The abdomen had now become much distended. The superficial veins were enlarged, but there was no pain, and pressure caused no sign of discomfort. The child was very weak, and slept with eyes half open. The appetite failed, and she was very peevish. There was no squint, inequality of pupils, or other cerebral symptom; nor was there any œdema of the legs or feet. Temperature at 8 a.m. 100.6° , 6 p.m. 103.4° ; pulse 148, respirations 48. Death took place on January 9.

On examination of the body on January 10, both lungs at their lower part were found to be attached by very slight adhesions to the diaphragm. The adhesions could be separated with perfect ease, and there was no fluid in the

pleura. Both lungs contained numerous grey granulations underneath the pleura, and also scattered through the substance of the organs. They crepitated less perfectly than natural, and large quantities of bloody serum escaped on pressure. Small patches of lobular collapse were seen here and there upon the surface. Both lungs contained also a few small caseous nodules of about the size of a pea; and in the apex of the right lung was a larger patch of catarrhal pneumonia, which was undergoing caseous degeneration.

On opening the abdomen the peritoneum was found to be covered by a thick layer of cheesy lymph, which penetrated between the coils of intestine, matting them together and gluing the upper surfaces of the liver and stomach to the diaphragm. A small quantity of thick purulent-looking fluid was found in the abdominal cavity. Some yellow tubercles were seen underneath the peritoneum. The intestine was so matted into a confused mass that it could not be laid open. The liver was large, weighing fifteen ounces, and felt firmer than natural. Its section was not stained with iodine.

There are two sequelæ of measles so commonly met with that they must be within the experience of all who have any familiarity with the diseases of children. One of these sequelæ is catarrhal pneumonia; the other acute tuberculosis. A persistent high temperature following an attack of measles, usually indicates the presence of one or the other of these disorders. On examination of the child whose case has just been related, it was seen that the disease could not be catarrhal pneumonia; for had it been so, a history which comprised two months of fever and cough following measles would have entitled us to expect some considerable evidence of lung consolidation. As it was, the only signs discoverable were those of bronchitis. This bronchitis was evidently of considerable duration, for the cough was stated by the mother to have been present ever since the attack of measles, and no history of any

more recent lung affection was hinted at. Bronchitis of such standing, combined, too, with a high temperature, could not be the result merely of severe pulmonary catarrh, but was evidently dependent upon some other cause ; and this other cause, there could be little doubt, was tuberculosis. All the other symptoms pointed to the same conclusion. Obstinate fever, wasting, cough, and digestive disturbance, lasting for two months, and without evidence of lung consolidation, could only be due to acute tuberculosis.

The existence of peritonitis was not recognised during life ; for, with the exception of some slight swelling of the abdomen, which was attributed to the flatulent distension so common in weakly children, there were no symptoms to draw attention to this part of the body. If on this account alone, the case is an interesting one, as showing with how few symptoms so serious a disorganisation may be associated. Had the peritonitis been recognised, it would have furnished additional evidence as to the nature of the disease.

In this case the tubercular outbreak must be attributed directly to the measles, by which it had been preceded, for the child had been healthy before the attack, and there was no history of consumptive tendencies in the family.

In the following illustration the tuberculosis supervened upon long-standing disease of bone :

CASE XXXIV.—ACUTE GENERAL TUBERCULOSIS FOLLOWING OLD-STANDING DISEASE OF BONE.

Robert T., aged $8\frac{1}{2}$ years, had been as an infant the subject of rickets, and could not walk before the age of three years. He was never a strong boy, and had always perspired profusely about the head at night. The parents were in bad circumstances, and for some months his diet had been poor and insufficient.

On October 22 he had a rigor, followed by thirst, hot skin, sore throat, and slight delirium. This attack lasted three days. On October 27 he complained for the first time of pain in the left hip, and was admitted into the East London Children's Hospital, under the care of Mr. Reeves. An abscess soon formed at the seat of pain. It was opened, and a large quantity of offensive pus escaped. After the operation the abscess did not heal up, but continued to discharge; the boy's health remained poor, and he lost flesh considerably. For some time the temperature was high, varying from 100° to $102^{\circ}6$ in the evening; but in the middle of November it became natural, and remained so until January 10, when it rose again to $103^{\circ}6$. From this time it was permanently elevated, rising in the evening to from 101° to 104 , and sinking to 99° in the morning. No disease could be detected in the bones of the hip, but the periosteum over the left trochanter sloughed off, leaving the bone exposed. There was considerable discharge from the wound, and the boy still wasted and looked weak and ill. He had no rigors, rapid fluctuations of temperature, or other signs of pyæmia. The spleen, however, was noticed to be enlarged. Examination of the chest was frequently made, but resonance was normal, and a faint sibilant wheeze caught occasionally towards the end of inspiration was the only abnormal sign to be detected.

The patient continued in much the same state, with a high temperature but a very trifling cough, and getting thinner and thinner, until August 16, when he complained of frontal headache, and vomited several times. The bowels were confined, and his pulse was noticed to be slow.

On the following day he had a convulsive fit. The temperature that evening was $101^{\circ}2$. During the three following days he had as many as twenty fits, which involved both sides of the body. The boy's intellect seemed also affected;

he talked incoherently, and his sight appeared to be impaired. Albumen was found in the urine.

After the fourth day the fits were not repeated, but the patient seemed much exhausted, and he passed gradually into a comatose state, in which he died on August 23.

A few days before death the temperature of the body fell to a point below the ordinary level of health, and remained low until the end.

In this boy the tubercular disease was found to be very widely distributed. Grey granulations were discovered in the brain, pia mater, lungs, peritoneum, liver, and spleen. The latter organ was enlarged to about three times its natural size, and seemed to consist of little more than masses of tubercle. The mesenteric glands were large, cheesy, and adherent to one another. The cervical and bronchial glands were also enlarged and cheesy. The ventricles of the brain contained much fluid, and there was a small tubercular tumour of the size of a filbert attached to the upper surface of the right lobe of the cerebellum. On examination of the left hip the great trochanter was found to be extensively necrosed, and to be entirely separated from the rest of the bone.

The tubercular attack in this case must be dated from January, when the temperature became permanently elevated. From that time the course of the disease was very suspicious of tuberculosis, although the trifling amount of cough, and the almost complete absence of physical signs about the chest, caused some hesitation in making the diagnosis.

A case such as the above, seen for the first time towards the end of the disease, might be easily mistaken. A young practitioner finding frequent convulsive attacks, combined with albumen and perhaps a few blood-casts in the urine, might be excused if from such symptoms he inferred the presence of Bright's disease with uræmic convulsions. Such a diagnosis might, however, be corrected by noting the extreme emaciation, the absence of œdema or of urinous

odour from the breath, and by inquiring into the general history of the case, especially remarking the absence of any history of scarlatina. Still, in some of these cases œdema is present, but it is never so general or in such quantity as to conceal the wasted condition of the body. Upon this, therefore, upon the absence of urinous odour, and upon the history of the case, the diagnosis would chiefly depend.

The diagnosis of acute tuberculosis presents many difficulties. So long as the disease is general, its symptoms are indefinite, and present no characteristic features by which the nature of the complaint can be immediately recognised. Even when other and local symptoms have arisen, showing direct implication of special organs, the new phenomena, being merely the result of familiar tissue-changes excited by the presence of the grey granulations, have no such unmistakeable character that their constitutional origin can be at once detected.

Acute tuberculosis begins insidiously, creeping upon the child by slow degrees. Nutrition is impaired almost imperceptibly, and there is no startling symptom of ill-health to arrest attention and excite the anxiety of the friends. When at last the state of the patient has attracted observation, the causes of wasting in young children—in infants especially—are so numerous, that out of the many morbid conditions which may give rise to a similar group of symptoms, the detection of the exact source of evil is often far from being an easy task. In all such cases the thermometer is of the utmost service, for a continued low temperature in the evening is incompatible with tuberculosis. Therefore, in all infantile diseases accompanied by wasting, general tuberculosis may be excluded if the evening temperature be normal for many days in succession. On the other hand, the temperature in a wasting infant may be high without the case being necessarily one of tuberculosis. Aphthæ of the mouth and inflammation of the gums are common complica-

tions in children suffering from abdominal derangements, and cause for the time a considerable amount of fever. So also in dentition, which is active at this period, the bodily heat is much increased. Dentition, indeed, is sometimes a cause of great and groundless anxiety. It is often accompanied by a certain amount of catarrh of the stomach, which interferes temporarily with nutrition, and which, if the process be a long one, may induce a very evident diminution in the weight of the body. If several teeth be cut in succession, the temperature may remain high for a considerable time, and continued wasting with fever may excite serious apprehensions as to the nature of the ailment. It is, then, of very great importance to exclude other causes of pyrexia in every case of suspected tuberculosis, and in infants the mouth and gums should always be examined. We may, however, get a hint as to the nature of the case by noticing the height to which the mercury rises. In uncomplicated tuberculosis the temperature is seldom at first over 101° in the evening, and it falls notably towards the morning ; while in dentition, when the gums are much inflamed and swollen, a temperature of 104° or 105° even in the morning is far from uncommon.

One of the peculiarities of acute general tuberculosis is the fewness of the symptoms by which it is accompanied ; and the diagnosis is often arrived at by exclusion, there being no other discoverable cause to which the pyrexia can be attributed. If we are called to an infant who is suffering from moderate fever and steady loss of flesh, and after a careful examination of the whole body can find no explanation of the elevation of temperature, we should strongly suspect tuberculosis. A pinched, wearied expression in the child's face, a loss of elasticity in the skin, and the existence of slight œdema of the extremities, furnish strong corroborative evidence ; and if we can discover any history of fatal illness of a similar kind in other children of the same family, little doubt can remain.

In older children, in cases where the symptoms remain for a considerable time indefinite, the most prominent among them being wasting with pyrexia, the diagnosis lies between tuberculosis and enteric fever.* The resemblance between these two diseases is often exceedingly close, more especially on account of the state of the bowels. A slight diarrhoea is common in acute tuberculosis; and looseness of the bowels is, as is well known, a frequent symptom in enteric fever. But in children suffering from the latter disease diarrhoea does not always assume the obstinate character which is so common a feature in the adult; indeed, in at least a third of the cases, the bowels, instead of being relaxed, are obstinately confined. The difficulty of diagnosis is increased by the slight severity of the other symptoms in many cases of typhoid fever in children. The patient is feverish, loses flesh, and his strength is considerably reduced. The headache, however, is insignificant; the tongue moist; diarrhoea absent or mild; the belly only moderately swollen, and not tender on pressure. The child sleeps well at night, without delirium, and shows no unwillingness to take food. Such a case, which would in former days have been classed under the heading of remittent fever, is often very perplexing. The temperature is, however, higher than that common to tuberculosis, reaching to 104° or 105° . On careful examination, too, we can perhaps discover a few of the characteristic rose-spots upon the belly, chest, or back. In children, however, the eruption is not unfrequently absent; but even in these cases the condition of the spleen seldom fails to furnish an important indication. In tuberculosis it is rarely enlarged until the disease is far advanced; while in typhoid fever, as early as the end of the first week, it may

* It is, of course, assumed that in every case all the organs are submitted to a careful examination, otherwise many important diseases might be passed over. To mention one amongst others: pleurisy may give rise to but few symptoms, and without an attentive examination of the chest may easily escape recognition.

be felt to project inwards towards the middle line from under the cover of the ribs. The position and aspect of the child also furnish important information. In tuberculosis the child's position in bed is often on the side, with his arms folded on his chest and his thighs flexed on his belly. In typhoid fever, as long as the temperature continues high, the decubitus is almost invariably dorsal. Again, in tuberculosis the features are pinched, and the face often wears an expression of distress which is out of all proportion to the actual severity of the symptoms. In typhoid fever, on the contrary, unless the disease be very severe, the facies is little altered. In grave cases the countenance may assume various expressions of terror, rage, etc., suggested by the delirium; but these are usually passing emotions, and although the child looks really ill, his face generally denotes rather stupid indifference than anxiety. The history of a strong family tendency to tubercular disease does not decide our doubts in favour of tuberculosis, for enteric fever is quite possible in such a constitution. In many cases it will be necessary to withhold a positive opinion, and to wait for time to decide the nature of the case. In typhoid fever the temperature seldom remains elevated longer than three weeks, and in mild cases often becomes normal at a still earlier period.

When, in a case of acute tuberculosis, severe chest symptoms arise, the physical signs are those naturally belonging to the pulmonary complication, and bear no special character which at once establishes the constitutional origin of the new lesion. When, therefore, a case is seen for the first time at this period, it is not always easy to decide at once upon its nature. We find a child suffering from all the symptoms of capillary bronchitis; he is agitated; his face is distressed and more or less livid; his pulse quick, his breathing laborious, his cough frequent and harassing; a fine bubbling rhonchus pervades the whole of both lungs. The thermometer placed under the arm or in the rectum marks

103° or 104°. This elevation of temperature indicates that the disease is not of a simple character, and the diagnosis lies between tuberculous bronchitis and broncho-pneumonia. Here a history of the disease is of the utmost importance. If the lung affection have been preceded by several weeks of wasting and pyrexia, it is probably of tuberculous origin, more especially if there be a history of consumptive tendencies in the family. If, in addition, we find the cough, although frequent, to be loose, and if a careful exploration of the chest discovers no spots of dulness or bronchial breathing; if, too, the rhonchus is everywhere of the crisp bubbling variety, without any approach to the dry crepitation peculiar to pneumonia, we may conclude that the case is one of tuberculous bronchitis. Broncho-pneumonia may, however, actually come on under such circumstances, and spots of local consolidation may be discovered. Here we are forced to rely upon the previous history of the child to determine whether the disease be a primary one or be secondary to acute tuberculosis of the lungs; and unfortunately in such a case little more than *probability* can be arrived at, so long as the special symptoms remain limited to the chest. If, however, the case be still further complicated by indications of intracranial disease, and convulsions occur with squinting, unequal pupils, ptosis, and rigidity about the joints, we may then safely infer that the case is one of acute tuberculosis involving especially the lungs and the meninges of the brain. The presence of these distinct signs of intracranial irritation is important, for the occurrence of convulsions alone will not give us the assurance we require. It is not uncommon for simple broncho-pneumonia in the child to be brought to a sudden termination by a convulsive seizure, and therefore we must look for the symptoms mentioned above, in order to be enabled to establish the tuberculous nature of the disease.

CHAPTER IX.

ACUTE TUBERCULOSIS OF THE PIA MATER—TUBERCULAR MENINGITIS—ACUTE HYDROCEPHALUS.

Two forms of the disease—primary and secondary.

Symptoms—Premonitory—First stage—Second Stage—Third and final stage—Variations in the symptoms.

Duration of the disease.

Case XXXV.—Ordinary attack of tubercular meningitis.

Cases XXXVI. and XXXVII.—Acute meningitis occurring in the course of acute general tuberculosis—Comment.

Case XXXVIII.—Tubercular meningitis occupying the convexity of the brain—Comment.

Diagnosis.

ACUTE tuberculosis of the pia mater is a very important and interesting form of the disease. The presence of the grey granulation in the pia mater is soon followed by inflammation of this membrane, and by effusion of fluid into the ventricles, and softening of the substance of the brain. This form of acute tuberculosis may occur at any period of life, but is most frequently met with between the ages of three and ten years. It is, however, far from uncommon even in infants of a few months old; but in such cases the distribution of the grey granulation is almost invariably general over the whole body, and the symptoms arising from the cranial cavity occur late in the disease, often in the course of an acute pulmonary tuberculosis, and confirm a diagnosis which the previously obscure symptoms may not have made sufficiently evident. In such cases the meningitis is spoken of as a *secondary* disease. This is the common form in children under twelve months or even two years old.

At a more advanced age children are more subject to the *primary* form, in which—whether the presence of the grey

granulation be or be not limited to the cranial cavity—the manifestations of cerebral distress are at any rate the first local symptoms to arise in the course of the general complaint. These local symptoms may indeed occur so early that the brain affection appears to come on suddenly in the midst of a state of health. This, however, is quite exceptional. As a general rule the meningeal disease is preceded by the signs which have been described as generally accompanying the constitutional malady, combined with others indicating some mental disturbance and irritability.

In the large majority of cases, then, a certain amount of languor, pallor, and loss of flesh precedes by a few weeks the more special symptoms. It would be a mistake, however, to suppose that there must be necessarily any marked emaciation. The eye can often recognise no difference in the bulk of the child, but the hand appreciates a certain softness and flabbiness of the muscles, and the scales show that the actual weight of the body is diminished. The child is drowsy by day and easily fatigued; he is fretful, listless, and unwilling to play, and at night sleeps uneasily, and by fits and starts. Perhaps one of the most characteristic symptoms at this time is increased emotional sensibility. The child shows exaggerated distress at the most trifling occurrences, and when spoken to may burst into tears. This usually alternates with sluggishness of mind and drowsiness. He may show unusual stupidity at his lessons, or may stop in his play and fall asleep on the floor of the room. The temperature at this time is slightly raised, although there need be no increase of bodily heat appreciable by the hand. Cephalalgia, slight in amount, is often present; it is increased by movement, or excitement, or by a bright light. Young children may be noticed to lift the hand frequently to the head, and older children will often complain spontaneously of headache. The pain is usually referred to the crown of the head. It is often accompanied by giddiness. At the same time there are

usually evidences of disturbance of the digestive functions ; the appetite fails, and the bowels become irregular. When such symptoms are present, the case should be always regarded with anxiety, and a careful watch should be kept over the patient so that the occurrence of any more characteristic symptoms may not escape notice.

The symptoms of acute hydrocephalus fall naturally into three groups. As a part of acute general tuberculosis, the disease is a constitutional one, but, like most other general maladies, its chief manifestations result from local changes of structure. The local lesion is at first limited, and consists of irritation or inflammation at the base of the brain, due to the presence of grey granulations in the pia mater. The irritation excited by this local condition is evidenced by symptoms of exalted activity in the parts supplied by nerves proceeding from the brain at the seat of disease. Afterwards, as effusion takes place into the ventricles and substance of the brain, other symptoms arise showing diminished nervous activity, and gradual destruction of the functions of life. We find, therefore, a stage of invasion breaking in upon the indefinite symptoms of the general disease, and first marking the presence of a local lesion; this is followed by a stage of irritation during which the nervous functions are exalted ; and lastly, the disease passes into the third and final stage of paralysis and death.

The first of the special symptoms pointing more directly to the nature of the disease is vomiting. The vomiting occurs not only after food has been taken, but at other times, and is often excited by raising the child up in his bed. The first occurrence of vomiting marks the first day of the disease ; it is generally accompanied by constipation, which may be obstinate. Vomiting coming on in a child when the stomach is nearly empty, especially if accompanied by constipation, should always be regarded with very grave suspicion. The abdomen is normal to the eye, but an early and characteristic symptom is a peculiar flabbiness and

compressibility of the parietes. The muscles seem to have lost all elasticity, and yield before the pressure of the fingers with a readiness and absence of recoil which reminds one of unbaked dough. The child looks ill; his features are more or less pinched, and the expression is anxious or spiteful. The appetite is lost, and the tongue may be furred thickly, although it is sometimes clean and moist. A convulsive fit occasionally occurs at this time, but as a rule it is a later symptom. The pulse is rapid; the breathing unequal and interspersed with sighs; and the temperature rises to between 100° and 101° . The headache increases, and is often accompanied by vertigo, so that the child will stagger in his walk, or if lying down will say that his bed is falling. He is sullen and drowsy, or excessively peevish and fretful.

The above symptoms constitute the invasion period of the disease. The beginning of the next stage—the stage of irritation—is marked by a sudden fall in the rapidity of the pulse. It becomes slow and irregular, sinking usually to between 60 and 70, but is quickened by movement. The irregularity affects rhythm as well as force. The irregularity of rhythm is particularly well marked, because, as a rule, the intermission is complete. This is the more evident on account of the slowness of the pulse. Several beats strike the finger in regular succession, then suddenly one beat is entirely omitted, and there is a pause until the rhythm is again taken up by the succeeding beat. The frequency of the intermission varies in different cases, but the intervals are seldom regular.

The headache often now becomes so violent that the child cries out with the pain. This is especially noticeable at night when gas-jets are burning, for a bright light seems to have the effect of greatly increasing the pain; and the child will be seen to squeeze his eyelids tightly together, and to cower down behind the bedclothes. At night, indeed, he is very restless, crying out frequently, and show-

ing great agitation. The cries are probably often due to delirium, for they cannot be always attributed to the headache. A sudden sharp squeal, common to this period, has been considered so characteristic of the disease that it has been called the hydrocephalic cry ; but this symptom is not present in every case.

The pupils, which during the invasion period are usually small, now dilate, and are often unequal in size, although they still act regularly with light. There is very often transient strabismus of one or both eyes. If an ophthalmoscopic examination be made, the optic disc and retinal vessels will be found to be congested. The expression of the child at this time is peculiar: the brows are contracted, and there is a peevish, often spiteful, look on the face which at once attracts attention.

The respiration increases in rapidity, and continues to be very irregular. The child sighs deeply, and makes long pauses in breathing, the chest often remaining motionless for several seconds. His face flushes, and any—even the slightest—irritation of the skin, such as pressure with the fingers upon the forehead, is followed by a vivid blush—the *tache cérébrale* of Trousseau. After three or four days the vomiting usually ceases, but the bowels remain confined, and only act under the influence of a stiff aperient. The belly is retracted, often deeply hollowed ; and the parietes retain the flabbiness and want of elasticity which has been already remarked upon.

After a few days the case enters into the final stage. The symptoms of irritation or exalted sensibility give place to those of depression, and there is a gradual extinction of the functions of life. In this period the headache ceases, but the delirium continues. There is stupor, and the sensitiveness to light is lost. Delirium and stupor often alternate with one another, the child dozing heavily for a few minutes, and then waking up with a passionate cry. At this stage the appearance of the child is often characteristic.

He lies with cheeks brightly flushed, and with an expression of complete placidity, apparently asleep ; but his eyelids close imperfectly, allowing some of the white to be seen, and he occasionally partially raises and then lowers his eyelids, as if an inclination to rouse himself were overcome by returning drowsiness. If the eyelids are lifted up by the finger, the pupils will be seen to be widely dilated and often unequal in size, and there may be an evident squint. The somnolence gradually deepens into coma, and the insensibility of the child is complete. At first, however, he may look up when his name is called loudly, but he shows no other sign of consciousness, and closes his eyes again almost immediately. Sometimes in these cases the child suddenly appears to wake up, and may be noticed for a few seconds to move his eyes from side to side as if looking about him, but it is difficult to say whether at this time there is any actual return of consciousness. It may, however, happen that the stupor really clears off for a while, and for several hours the intellect may appear almost natural, the child's interest in his toys returning, as if he were about to recover. But false hopes should not be excited by this apparent improvement. It will be noticed that the squint or inequality of pupils still persists ; and although in rare cases consciousness is preserved until the close, the stupor usually returns after a longer or shorter interval.

During the coma the flush on the face often fades, and is replaced by pallor. The sighing and irregularity of respiration continue. The pulse may remain slow, but generally in this stage it quickens again, and becomes rapid and more regular, although it may still occasionally intermit. A difficulty in swallowing is sometimes noticed at this time, and there may be retention of urine. If the urine be examined, it is often found to contain albumen.

Soon paralytic lesions begin to be observed. One eyelid droops ; the squint may become permanent ; there may be

paralysis of the arm or leg—one or both—on one side, with often rigidity about the joints; the jaws may get stiff, and there may be a stiffness of the muscles at the nucha, with dragging backwards of the head. Although the child is perfectly insensible, automatic movements continue. Thus, he may push away a spoon held to his lips, or with his hand he may grasp objects within his reach, as the bars of his cot. In such a state a child may be seen lying, with half-closed eyes, perfectly motionless on one side of his body, while the other side is in constant movement, the knee being alternately raised and lowered, and the hand wandering to his head or lips, or being thrown about on the bedclothes.

Tremors and convulsive movements generally occur at this time. The eyeballs can be seen through the half-shut lids to have a rapid quivering movement; convulsive twitchings take place on one side of the face or in one or more of the limbs, and subsultus tendinum greatly interferes with observations upon the rapidity of the pulse. The temperature rises from 101° to 103° or higher, and sometimes before death may mount to 106° or 108° . Occasionally, however, instead of rising, the temperature falls, and during the last few hours of life the heat of the body, as registered by a thermometer placed within the rectum, may be below—sometimes very considerably below—the level of health.

In cases where ossification of the cranial bones is still incomplete, the fontanelle is generally found to be elevated and tense. Sometimes, however, it is depressed, and even in cases where bending back of the head is a prominent symptom, and where a post-mortem examination shows large effusion, with much flattening of the convolutions, the fontanelle, so far from being elevated, has been noted to be depressed below the level of the surrounding bones from the beginning to the end of the disease. The state of the fontanelle must not, therefore, be relied upon as a test of

the amount of effusion. Towards the end of this stage the constipation often yields to a slight relaxation of the bowels. The motions, which are very offensive, are, with the urine, passed in the bed.

Ophthalmia often occurs before death, and the cornea may become ulcerated. Death may take place in a convulsive fit, or the child may pass away quietly. The end is usually preceded by aphthæ of the mouth, swelling of the previously sunken abdomen by gaseous distension, frequent pulse, profuse sweating, and copious secretion into the air-passages of the lungs, for on auscultation much fine bubbling rhonchus may be heard all over the chest.

To complete the picture of this form of tuberculosis, some variations in the symptoms must be noticed. Thus, the bowels are not always confined in the beginning; sometimes they are relaxed, with liquid offensive stools, but the looseness soon gives place to constipation, which then persists to nearly the end of the disease.

Again, the pulse may be infrequent from the very first, or, if rapid in the beginning, may soon fall and remain infrequent until the close. In other cases its rapidity may be maintained until a few days before death. Different cases will be found to vary greatly in this respect, but in all the pulse, if carefully watched, may be noted to be infrequent at some period of the disease, if only for a few hours. The pulse of tubercular meningitis is excited by very gentle movements, and therefore it is important that observations upon its rapidity should be made while the child is quiet.

The temperature also is subject to no fixed rule. In the majority of cases it is only moderately raised, and for days together may maintain a uniform level of less than 100° . Thus, in a boy aged six years, the evening temperature on the eleventh day was 99.2° , and on the twelfth, thirteenth, and fourteenth days 99° , after which it became more elevated. This is exceptionally low, but until towards the

end of the disease the mercury seldom reaches higher than 102° in the evening.

The cerebral flush is not a constant symptom, and the amount of redness produced by irritation of the skin will be found to vary very much in different cases. It is often not greater than that produced by the same means in a healthy child. The headache also is very variable—sometimes it is exceedingly violent and paroxysmal, and appears to be accompanied by tenderness of the scalp; at other times it is so slight as to be scarcely complained of; but in almost all cases the discomfort produced by a bright light may be observed. In the same way the intensity of most of the symptoms will be found to differ very much in different cases—sometimes one, sometimes another, assuming the chief prominence,—but in their main features all varieties of the disease show a very close resemblance.

The average duration of the illness, counting from the first day of vomiting, is twelve days. It seldom ends sooner than seven days or lasts longer than twenty-one.

The following case furnishes a very good example of an ordinary attack of acute hydrocephalus, and it is especially interesting from the fact that the patient was under observation for some time previous to the beginning of the first definite symptoms of the disease.

CASE XXXV.—ORDINARY ATTACK OF PRIMARY TUBERCULAR MENINGITIS.

William G., aged four years, was admitted into the East London Children's Hospital on December 4. He had been a healthy boy until a month before admission. At that time he had an attack of measles, and on the disappearance of the eruption did not thoroughly recover. The child seemed ailing; he coughed occasionally, and was a little feverish. He slept well, however, and had a good appetite; his bowels were regular, and he did not notably lose flesh.

On admission he was found to be suffering from pulmonary catarrh, and the percussion-note was thought to be a little high-pitched at the right apex, with some hollow quality of breathing there and at the left side, but nothing very definite was discovered. For a few days the evening temperature was 100° , but on December 10 it became natural, and remained so until the 17th, when the child was sent as convalescent into the country. On December 30, however, he was sent back to the hospital as not improving. For the five following days he was noticed to be quiet and silent. He seemed unusually drowsy, and his appetite began to fail. The bowels were not confined, but acted two or three times in the day. During these five days his evening temperature was successively 100.4° , 98.4° , 98° , 99.6° , and 97.6° ; and on this last evening (January 4) the pulse, which had been always over 100, was found to have fallen to 74 and to be irregular. On the next morning (January 5) he vomited, and his bowels became confined. The temperature in the morning was 99.4° ; pulse 72, intermittent; respirations 36, irregular; temperature in evening, 101.6° . These invasion symptoms marked the first day of the established disease. The vomiting on this and the following day was very troublesome; he retched frequently, but with little effect. He was very drowsy, dozing and waking up from time to time with a scream. There was no squinting, and the pupils were equal.

January 6 (second day).—9 a.m.: Temperature 100.2° ; pulse 62, very intermittent; respirations 18. 6 p.m.: Temperature 100° ; pulse 78; respirations 26.

7th (third day).—Very stupid, but answers intelligently if roused. Occasionally gives a short angry cry, and then lies quiet. Conjunctivæ injected, and transient internal squint of both eyes. Pupils equal, and act naturally. Belly not much retracted. 9 a.m.: Temperature 100.6° ; pulse 80; respirations 22. 6 p.m.: Temperature 101.2° .

8th (fourth day).—Dulness increases, but he still can be

made to answer intelligently. He passes his water, however, in the bed. Bowels acted to-day (first time for forty-eight hours) after a purgative followed by an injection, and a large *ascaris lumbricoides* came away with the stool. Face rather flushed, but *tache cérébrale* not well marked. 9 a.m.: Temperature 101° ; pulse 84, still intermittent; respirations 28, still irregular and sighing. 6 p.m.: Temperature 102° ; pulse 96; respirations 40.

9th (fifth day).—Lies with eyelids half closed, apparently asleep; contracts brows, and every few minutes gives a peculiar squeal; expression of face spiteful; cheeks flushed, but not much redness produced by irritation of the skin; nares acting, but not with each breath; squints slightly with left eye, and left pupil is larger than right. When called by name opens eyes, but does not appear to understand question. Respirations 16, with frequent pauses and occasional deep sighs; pulse 160, regular in force and rhythm. Decided rigidity of muscles at back of neck, but none of muscles of limbs. Abdomen not particularly retracted, and walls flaccid. Percussion-note at right side of sternum over the second and third ribs rather toneless, and respiration high-pitched, but vocal resonance not increased. Elsewhere chest seems healthy. At 9 a.m., temperature 102.2° ; pulse 116; respirations 28. At 7 p.m., temperature 102.4° ; pulse 152; respirations 24.

10th (sixth day).—More comatose, and quieter; screams less frequently; lies with eyes half open and expression rather stupid than spiteful, but still is not completely insensible, for complains if disturbed. Face paler. Pupils nearly equal, and of moderate size; some internal squint of left eye. Bowels still confined; passes water in the bed. The nurse states that he has been rather convulsed, his limbs and eyes "working." Muscles generally flaccid, but still stiffness at back of neck, and some rigidity about right knee-joint. 9 a.m.: temperature 102° ; pulse 130, regular, but weaker; respirations 26, irregular both in depth and

frequency. 7 p.m. : Temperature 102° ; pulse 116 ; respirations 30.

11th (seventh day).—Quiet ; has ceased screaming ; seems less stupefied ; has spoken distinctly, and takes drink without difficulty. Pupils act well with light ; generally unequal. Belly rather distended ; bowels very confined. 9 a.m. : Temperature 103.2° ; pulse 116, regular ; respirations 30, sighing. 7 p.m. : Temperature 102.6° .

12th (eighth day).—Coma deeper ; face flushed and perspiring. Ophthalmia of both eyes ; pupils unequal. Irregular spasmodic movements of limbs, especially right arm and left leg. 9 a.m. : Temperature 101.8° ; pulse 112, regular ; respirations 40, very irregular. 7 p.m. : Temperature 103° .

13th (ninth day).—Lying as if asleep ; right cheek flushed. Pupils equal, large, contract with light. When eyelids are touched he opens both eyes and makes a moaning sound, but appears quite unconscious, although he is stated to have said "no" in answer to a question from his mother. Sighs frequently. Bowels relieved last night by injection. Same stiffness about knee-joints. Large bubbling rhonchus heard about chest. 9 a.m. : Temperature 101.6° ; pulse 116, irregular in force, regular in rhythm ; respirations 40, excessively irregular. 7 p.m. : Temperature 102.4° . In afternoon had a fit in which the right side of face alone was convulsed.

14th (tenth day).—State much the same. Coma seems deeper. Rigidity of extensor muscles of leg. Moans occasionally. Ulceration of left cornea. 9 a.m. : Temperature 103.2° ; pulse 128 ; respirations 44. 7 p.m. : Temperature 103.6° .

15th (eleventh day).—Sinking. Face covered with large drops of sweat. Abdomen more retracted than before. Respirations very irregular and laborious, 68 ; pulse excessively rapid and weak ; temperature 9 a.m. 104.4° , 3 p.m. 105.8° , 6 p.m. 106° . Patient died at 11 p.m.

On examination of the body the pia mater was found much injected at the base of the brain, and contained numerous small grey granulations; corpus callosum much softened. Lateral ventricles contained a large quantity of fluid, and the convolutions were flattened. The lungs contained patches of collapse and a few caseous nodules, but no grey granulations were found in them or in any of the other organs. The bronchial glands were enlarged and cheesy.

The preceding description represents a typical case of acute hydrocephalus, in which the brain symptoms are the first local phenomena to occur in the general disease. It is then called primary tubercular meningitis to distinguish it from cases where the grey granulation is more generally diffused over the body, and where the symptoms arising from the cranial cavity occur subsequently to others proceeding from the chest or the abdomen. In this the secondary form of the disease, the symptoms of the earlier stages are usually masked by the more prominent symptoms emanating from the organs previously attacked. In such cases the first manifestation drawing attention to the brain may be a violent convulsive fit, a paralytic lesion, or other sign indicative of the third stage of the disease; and it is not uncommon for acute tuberculosis with severe chest symptoms to be brought to a sudden close by this complication.

CASE XXXVI.—SECONDARY TUBERCULAR MENINGITIS.

John W., aged sixteen months, the illegitimate child of a consumptive mother, was said to have been healthy up to the age of six months, when he was attacked with sickness and diarrhoea, which lasted eight weeks. After this, however, he seemed to recover, and remained pretty well until the beginning of May, when he began to cough, and his breathing seemed oppressed. After some time these symp-

toms improved, and the child appeared, according to the mother's account, to be going on well, when, on June 9, he had a fit which lasted a quarter of an hour. During the following forty-eight hours the fits were frequently repeated, and three days afterwards (June 12) the child was admitted into the East London Children's Hospital. The temperature on the evening of admission was 102.8° . On examination of the chest on the following day, signs of consolidation were found at the upper part of the right lung, and the respiration in the right axilla was tubular.

The fits continued, and were very violent. During the attacks both sides of the face twitched, the tongue was bitten, and all the limbs became rigid and were convulsed. Between the fits, consciousness was not restored; the eyeballs oscillated, and the eyes were often turned upwards. There was no squint. The convulsions continued with short intervals until the evening of June 16, when they ceased. The temperature all this time was over 102° , both morning and evening. The pulse was quickened, but remained regular, 140 to 150; and the breathing was hurried and irregular, with deep sighs.

June 17.—No fit since yesterday evening. Child quite unconscious; face pale; nares act; eyeballs oscillate, and are turned rather upwards; pupils contracted, but act slowly with light; tongue ulcerated from being bitten during the fits. Breathing very rapid (75), but not laborious; intercostal spaces sink in rather deeply during inspiration. Pulse 157, regular. Skin wet with perspiration; very little flushing on irritation of the skin.

On examination of the chest, dulness was found at the upper half of the right side. The respiration in the axilla was intensely tubular; elsewhere on that side it was covered by a fine crisp bubbling rhonchus, which was heard all over both sides of the chest. There was also dulness at the lower half of the left back, and the respiration was blowing, becoming tubular at the base. 9 a.m.: Temperature 104.8° ;

pulse 174; respirations 75. 7 p.m.: Temperature 105°. The child died quietly the next day. Temperature a few hours before death, 105·8°.

On examination of the body, numerous small grey granulations were found in the pia mater, the lungs, and the spleen. The pia mater was much injected at the base of the brain, and a patch of effused lymph was seen on the left side. The whole brain seemed softened, and the lateral ventricles contained a large amount of fluid. Both lungs were the seat of pneumonic consolidation, and in the right were several small cavities.

In this case the occurrence of a convulsive attack was the first alarming symptom noticed by the mother. Before this event took place, although the child was no doubt feverish, and was, as we know, troubled with cough and some oppression of breathing, there was yet nothing in its condition which appeared sufficiently urgent to require medical advice.

On admission, the obstinate recurrence of the fits, the continuance of insensibility between the attacks, the oscillation of the eyeballs, and the other symptoms, made it evident that the convulsions were not merely sympathetic with the condition of the lung. In certain forms of pulmonary disease in the child, sympathetic convulsions are not uncommon. Thus, in the course of acute capillary bronchitis they are often observed, especially if that disease be complicated by collapse of the lung. They are also not unfrequent at the close of acute catarrhal pneumonia; and at the beginning of many acute diseases in the child convulsions may occur, taking the place of the rigor which often ushers in an acute disease in the adult; but it is very rare for sympathetic convulsions to appear first in the course of an acute pulmonary consolidation. Where this happens we should always, as in the present case, suspect the existence of some complication. All the symptoms here pointed to meningeal inflammation, and the

history of the attack, combined with the pulmonary symptoms, made it evident that the case was one of acute general tuberculosis. Where, as in the present instance, the first symptoms pointing to the brain are those of the third stage, it is rare to find a slow pulse. In the third stage of acute hydrocephalus, the pulse, as a rule, has become rapid and often regular, the period during which it is slow and intermittent having gone by.

CASE XXXVII.—SECONDARY TUBERCULAR MENINGITIS.

Louisa P., aged six months, born of a healthy family, had been herself a healthy child up to the age of five months, when, after a severe attack of acute diarrhœa, she began to waste, and for the following month seemed low and ailing. She was then (December 1) admitted into the East London Children's Hospital. The child was thin and weakly-looking, weighing only twelve pounds three ounces. The fontanelle was rather depressed, and she was very peevish and fretful. The cervical glands were enlarged and also those of the axilla and groins. The abdomen was very firm and tense, and the liver and spleen seemed to be slightly enlarged. Her bowels were opened twice a day. There were signs of slight consolidation of the right apex, and all over the right back the respiration was weaker than on the opposite side. Large bubbling rhonchus was heard all over the chest. The evening temperature at first varied between 99° and 100°, the pulse was small and rapid (120), and the breathing hurried (40).

The child remained very fretful until December 14, when she became more composed, her appetite improved, and she seemed generally better. She coughed less, and the bronchitis from which she had been suffering appeared to be subsiding. Her weight on this date (December 14) was

exactly the same as on the day of her admission into the hospital.

During the next fortnight the temperature varied between 99° and 101° , and the child seemed growing daily weaker. Her cough, too, returned and was very troublesome. On December 29 much consolidation was found on the right side. In addition, however, she had begun to squint with the right eye, and had occasionally slight twitchings of the muscles of the face. The temperature rose, and both the pulse and the respiration became quickened.

On December 31, 8 a.m., temperature 103° , pulse 150, respirations 60; 6 p.m., temperature 103.2° , pulse 156.

On the following day she seemed about the same; the squint, however, had ceased, and the pupils were equal. The temperature remained high (103° at 6 p.m.), and the fontanelle was much elevated.

On January 3 she had a violent convulsive fit early in the morning, after which the left pupil was noticed to be larger than the right. In the evening the fits returned, and the child died in the course of the night. The evening temperature during the last three days was 103° , 103° , and 102.8° .

A post-mortem examination showed the body to be the seat of acute general tuberculosis. The lungs were studded with grey granulations, and there was pneumonic consolidation of the right lung, which had softened into a small cavity towards the base. Numerous grey granulations were also found in the heart, kidneys, intestines, and in the meninges of the brain. The lateral ventricles were full of fluid, and there was besides a yellowish nodule the size of a small pea in the white matter of the brain near the lateral ventricle of one side.

The two preceding cases furnish very good examples of the way in which the earlier symptoms of acute hydrocephalus are obscured when the disease occurs in the course of an attack of acute tubercular phthisis. In such cases the

occurrence of marked brain symptoms is a sign that the disease is about to run a rapid downward course, and that the end is not far distant. The case of Louisa P. is interesting as showing that the emaciation does not always proceed with regularity, for during the first fortnight of the child's residence in the hospital her weight remained stationary. The patient's appetite was good, and the quiet and the better food she was taking at the time, no doubt contributed to this result.

The seat of tubercular meningitis is almost always at the base of the skull as in the preceding instances. In rare cases, however, the lesion is seated at the convexity of the brain. The symptoms are then very indefinite, neither partaking of the severity which distinguishes simple meningitis when seated at the convexity, nor exhibiting any of those marked characteristics which accompany tubercular meningitis when it occupies its more usual situation.

CASE XXXVIII.—TUBERCULAR MENINGITIS OCCUPYING THE CONVEXITY OF THE BRAIN.

Annie H., aged two years and a half, one of five children, two of whom had died of pertussis, had been delicate since infancy. She had never had measles, whooping cough or scarlatina, but from the age of three months had been very liable to cough, from which she was seldom free. At the beginning of July the child had a fit, and the glands of the neck began to swell, first on one side then on the other. They suppurated; and the discharge continuing, the child was brought to the East London Children's Hospital, and was admitted on August 16.

Her temperature that evening was 103.2° , but it fell the next morning to 101° , and remained between 101° and 102° until the morning of August 20, when it rose again to 103° . During this time the child had two fits.

On August 20 her condition was as follows:—Child has

all teeth, fontanelle about the size of a shilling. Sloughy-looking sores are seen on each side of neck below the angle of the jaw, and the cervical glands are enlarged. There is slight discharge from the right ear, but no offensive smell. Child lies on back with open eyes and perfectly placid expression; she moves her eyes from side to side, as if watching the persons around her cot. Pupils natural, and act well with light. Face pale, and is said not to flush; nares act in breathing; respiration very irregular, occasionally is suspended altogether for many seconds, while at other times several breaths are taken in quick succession. Pulse 144, regular. Twitching of tendons of wrist. Abdomen retracted, and very flaccid. Edge of liver can be felt one inch below false ribs: spleen not enlarged. No paralysis of facial muscles, or loss of power or rigidity anywhere.

On examination of the chest some slight dulness was found at the right supra-spinous fossa, and the respiration there was blowing. Much large clicking rhonchus was heard at all parts of the chest. Heart-sounds natural.

The bowels now became relaxed, and by August 24th the child was found to have lost five pounds in weight. The temperature still remained very high, and at this date was, at 9 a.m. $104^{\circ}8'$, at 6 p.m. $105^{\circ}4'$. Pulse 180, very weak, but regular. Respiration 56, and now quite regular. The child is still perfectly conscious; face pale, and nares act sometimes with respiration. Expression of face rather peevish, but features not pinched. She moves her eyes from side to side as if watching objects in the ward. The pupils act well with light, but the left is half as large again as the right. Abdomen now not retracted; wall quite flaccid, and no tenderness. Sudamina on chest.

Very little movement in respiration at the upper part of left front, and the interspaces there are motionless, although elsewhere they sink in. Breathing slightly laboured. All over the chest a fine bubbling rhonchus accompanies both

breath-sounds, especially on the left side. A little dulness under the left clavicle towards the sternum.

On August 25 the temperature fell to between 102° and 103° both morning and evening, but rose again on the following morning to 103.6° .

On August 27 the child had a fit, which lasted two minutes. With this exception consciousness appeared to be preserved. The eyes remained bright and intelligent-looking, but she was excessively apathetic, and made no complaint at being taken from her cot for examination. The physical signs remained the same. After this the pyrexia persisted, and the child died on August 29. The temperature about an hour before death (at 6 p.m.) was 104.8° .

On examination of the body, the pia mater covering the convexity of the brain was found to be inflamed and thickened. The two hemispheres were adherent, and numerous grey granulations were found on each hemisphere and in the large median fissure. The base of the brain, the medulla and upper part of spinal cord, the ventricles and levum interpositum, were free from granulations, and there was little excess of fluid. The substance of the brain appeared to be healthy.

In the lungs numerous granulations were found which had become yellow; these were all of the size of a pin's head, and were scattered over the upper part of each lung. No caseous nodules were found. The pleuræ were adherent, and the bronchial glands were enlarged and cheesy. No grey granulations were seen on the surface of the liver, spleen, or intestines. The mesenteric glands were enlarged.

This case furnishes a good example of the difficulty of diagnosis of tubercular meningitis when the inflammation is limited to the pia mater covering the convexity of the brain. There was no great excess of fluid, and no increase of pressure upon the brain, and as there was little irritation at the base of the skull, the interference with nerve function was correspondingly slight. Consciousness was retained,

and there was no rigidity or paralysis. The only signs, indeed, from which cerebral disease could have been suspected were the extreme irregularity of breathing and the retracted abdomen noted on August 20, the difference in the size of the pupils seen at a later period, and the occasional convulsive attacks from which the child suffered after her admission into the hospital.

In this case the diagnosis was not made during life. The physical signs were those of acute capillary bronchitis, passing into catarrhal pneumonia, and there was nothing in the history of the case to suggest acute tuberculosis. Still the progress of the disease might have given rise to suspicions, for convulsive attacks (themselves very suspicious when occurring in the course of an illness such as the present), combined with breathing so irregular and a retracted abdomen, and followed after a short interval by unequal pupils, formed a very suspicious combination of symptoms. Markedly irregular breathing is one of the most characteristic symptoms of cerebral disease, and the occurrence of inequality of the pupils is a sign of the very gravest import.

It is important to note the absence of the severe headache, delirium, and violent convulsive attacks which characterise simple acute meningitis of the convexity; for although the child had several attacks of convulsions, yet the fits were mild and short, and were followed by no paralysis, rigidity, or other sign of organic cerebral lesion.

The *diagnosis* of tubercular meningitis in the child is often very difficult in the early stage. The danger lies not so much in overlooking the disease when it is present, as in suspecting its existence in cases where many of the symptoms are simulated by other affections. On account of the activity of the nervous system in young children, and the commotion into which it is thrown by the presence of acute illness, symptoms indicating functional disturbance of the brain may accompany many acute disorders. We should

not, therefore, because such symptoms are present, rashly assume that the disease must be a cerebral one, still less that it must be tubercular meningitis. M. Rilliet has given a wise caution against hurry in the diagnosis of this complaint. An opinion should be formed with deliberation, and be rested not upon one or two symptoms, but upon a broad view of the whole aspect of the case, otherwise we shall be very likely to fall into error.

There are numerous symptoms indicative of cerebral disturbance, which, although frequently present in cases of organic disease of the brain, are yet perfectly compatible with mere functional derangement: vertigo, headache, delirium, twitching, convulsive movements, and irregularity of pulse, all belong to this class. One or all of these may be present in a case without necessarily denoting more than nervous irritability and distress, which may be of purely functional origin. Tubercular meningitis is essentially an insidious disease, and there is a certain regular progression in the symptoms which should be always looked for. The condition of the child before the actual outbreak, then, becomes of especial value, and a history of several weeks or months of languor and steady wasting greatly heightens the importance of the more definite symptoms of invasion. Vomiting, combined with constipation, and an elevated temperature, coming on under such circumstances, is always very suspicious. If afterwards the pulse become slow and irregular, the breathing also irregular and sighing, and there be drowsiness, with occasional squinting or inequality of pupils, we can have little doubt as to the nature of the complaint. In all such cases it is important to attend to the expression of the face. During the invasion stage the child looks really ill, and the features are pinched and drawn. In a case where tubercular meningitis is suspected, if this haggard expression be noted, however slight the other indications may be, we are justified in entertaining the most serious apprehensions.

A slow, irregular pulse, especially if the intermission be complete, very greatly increases the value of the symptoms with which it may be associated. It must, however, be remembered that during convalescence from acute disease, the pulse in a child often becomes slower and irregular, and may even completely intermit, especially during sleep. But in such cases the condition of the child is improving, while in the case of tubercular meningitis the slackening of the pulse is accompanied by increased mental heaviness, without any amelioration in the other symptoms.

In a typical case, where all the symptoms of invasion are present, the diagnosis is not a difficult one; but very often some are absent. Thus, the disease may begin without vomiting, or this symptom may occur once and not be repeated. In other cases the usual constipation is absent, and the bowels may even be relaxed; but if there be headache and drowsiness, with unequal breathing and deep sighs, and if the pulse become slow and irregular, we have grounds for the very gravest suspicion. Irregular breathing is in such a case of the utmost importance, and, if markedly unequal, is perhaps more to be relied upon than any other *single* symptom in the diagnosis of tubercular meningitis during the early period. When in doubt at this stage, we should look for the slighter forms of nervous derangement, as these are more to be trusted than the severer, which occur, as a rule, later on in the disease, and are only occasionally present in the beginning. Thus, it is the exception for the disease to begin with a fit of convulsions, or violent delirium, or other form of intense nervous excitement. In ordinary cases the symptoms are much less startling in their character, and must be looked for in order to be observed. Thus, the child may be noticed to contract his brows, wink his eyes, and avoid the light; his face wears a discontented or spiteful expression; the cheeks flush up and then grow pale, and the features are suddenly contracted at times with a strange grimace; the child puts his

hand to the crown of his head, complains of headache, and sighs frequently; he is sullen and morose, has a peculiar fixedness of look, and answers questions dryly or not at all. Symptoms such as these are very valuable, because they, or at any rate some of them, are almost invariably present during the stage of invasion.

Typhoid fever sometimes, in the beginning, presents a close resemblance to tubercular meningitis, for the fever may be marked at an early stage by drowsiness, severe headache, and a slow pulse. But a diagnosis can usually be arrived at by attention to the following points:—In fever the pulse is regular and not intermittent; the respiration, too, is regular, without the pauses and sighs so often noticed in inflammation of the cerebral meninges; and the belly, instead of being retracted, with soft doughy walls, is full if not swollen. Moreover, in typhoid fever there is usually cough, and the spleen is enlarged. These symptoms are never met with in primary tubercular meningitis.

The diagnosis of tubercular meningitis has often to be made by exclusion. In all cases of acute illness beginning with nervous symptoms and with vomiting, the meningeal disease may be suspected, and we can only wait for further information before venturing on an opinion. In the case of the exanthemata the appearance of the specific eruption enlightens us upon the nature of the case. Croupous pneumonia is sometimes accompanied by nervous symptoms which bear a close resemblance to those of tubercular meningitis. The diagnosis between the two conditions is discussed elsewhere (see page 55). Cases, however, often occur, where the symptoms are anomalous, and where, however carefully the patient be examined, the most experienced physician must be at fault as to the nature of the disease. It is, therefore, very necessary to proceed with caution, visiting the child frequently, and watching for fresh signs which may lighten up what was dark and put an end to the anxiety of doubt.

There are two forms of disease in which mistakes are particularly likely to happen. Thus, exhaustion in children often manifests itself by vomiting and constipation, with drowsiness, and great restlessness and distress. But here on inquiry we generally find a history of some recent acute attack, and the temperature is not elevated. Still, it is often far from easy to give a decided opinion at once upon the nature of such an illness. Again, in that form of chronic abdominal derangement, which I have elsewhere described under the name of "mucous disease,"* a sudden attack of gastric catarrh, with vomiting, constipation, and headache, may occasion very great anxiety, as the languor and wasting by which the disease had been accompanied may be easily mistaken for the premonitory stage of acute hydrocephalus. But in these cases, also, the temperature is generally low, and after a day or two, the gastric disorder having subsided, the child begins to improve. In gastric catarrh the heat of the body is not as a rule much elevated, often not at all, and the face does not show the pinched, haggard expression which is so common in intracranial disease.

During its later periods, tubercular meningitis may be confounded with any other cerebral lesion. To make a distinction, it is important to obtain a clear account of the history and progress of the illness, for the insidious beginning of tubercular meningitis is an important element in the diagnosis. The disease ought not to be confounded with simple meningitis, for the course and character of the two maladies are essentially different.

Simple meningitis begins suddenly in a state of good health or after an accident, with either violent convulsions, high fever, and rapid breathing, or with severe headache, fever, and sickness; the patient then gets drowsy, and is excessively restless; the restlessness passes into wild deli-

* See "Wasting Diseases of Children," fourth edition, p. 208.

rium ; and after a very few days collapse comes on, and the child dies exhausted. Constipation, is slight, and the pulse, although intermittent, is not slow as a rule. The disease, then, begins violently, and increases in violence as it goes on. Its duration is short, and it rarely lasts longer than a week.

Cerebral tumour also manifests itself suddenly, generally by a convulsive fit, or some other severe nervous symptom, such as a paralytic lesion, with violent paroxysmal headache, and perhaps vomiting. It is the exception for it to begin with the invasion symptoms common to tubercular meningitis—viz., vomiting, constipation, and headache alone. In any case the course of the two diseases is very different. In tumour the convulsions are violent, frequently repeated, and are followed by paralysis or rigidity in the convulsed parts. The paralytic lesions are various, and usually include some one of the organs of special sense. As a general rule, the intellect is preserved till late in the disease, and there is none of that drowsiness and dulness of mind which is one of the most prominent symptoms in tubercular meningitis. If, however, the tumour be tubercular, there are often granulations in the meninges causing meningitis. In such a case the symptoms of cerebral tumour would be confused, and combined with those dependent upon the complication.

In cases of disease of the temporal bone with the formation of a cerebral abscess, the symptoms are often obscured by those of purulent meningitis, which is often present at the same time. As a rule, they begin suddenly. The child screams with pain in his head, is drowsy and stupid, and refuses his food. The temperature is moderately elevated, 102° , the pulse slow, 75 ; the pupils are contracted. The drowsiness deepens into stupor and convulsions occur, or rigidities are noticed in the joints. Paralysis or paresis is usually to be discovered. It is variously distributed according to the seat of the abscess, but there is usually

hemiplegia. Eventually the child dies exhausted. Pyæmia may be present, still further to complicate the case.

The occurrence of hemiplegia in a case where drowsiness, convulsions and coma have supervened upon a severe attack of otitis, should point to the nature of the lesion. The existence of long-continued otorrhœa, with a history of previous deafness, is a valuable assistance in the diagnosis of such cases.*

Tubercular meningitis is almost inevitably fatal, and we should not therefore allow ourselves to be deceived, by any sudden amelioration in the symptoms, into a belief that the child is about to recover. Such sudden changes are not uncommon, but they are most deceptive, and when they occur, it is our duty to warn the friends of the patient of the little confidence that can be placed in the apparent improvement.

* For a full description of the preceding diseases and the peculiarities they assume in early life, the reader is referred to the author's larger work, "A Practical Treatise on Disease in Children."

CHAPTER X.

TUBERCULOSIS OF THE PERITONEUM—TUBERCULAR PERITONITIS.

COURSE less acute than in other forms of tuberculosis—Obscurity of early symptoms.

Symptoms—General and local—The latter occasionally absent.

Case XXXIX.—Tubercular peritonitis—Apparent amendment—Relapse and death—Comment.

Case XL.—Tubercular peritonitis complicated with ulceration of bowels.

Diagnosis—Importance of tenderness as a symptom.

Treatment of tuberculosis—Of general tuberculosis—Preventive treatment. Treatment of pulmonary tuberculosis—Of tubercular meningitis—Of tubercular peritonitis.

TUBERCULOSIS, when it affects the abdominal cavity, is usually a chronic complaint. As a rule, the disease runs a very sluggish course, and the only acute or pressing symptoms are those which mark the closing period of the illness. The disease, indeed, differs so widely from all other forms of tubercular inflammation, that one is sometimes tempted to doubt whether tuberculosis as it attacks the abdomen, can be always the same disease as tuberculosis as it affects the head or the chest. The frequent absence of general symptoms, the slower course of the illness and its occasional favourable termination are all features which seem to distinguish this form of tuberculosis from the tubercular disease as it affects other parts of the body.

Although chronic as a rule, the course of tubercular peritonitis is not always so prolonged. In rare cases the inflammation is acute. In this exceptional shape the abdominal mischief forms part of an attack of general

tuberculosis. In many of the cases of acute tubercular peritonitis which have come under my notice, the peritoneal inflammation has been curiously latent. A good illustration of this form is seen in case (XXXIII.) narrated on a previous page (see page 247). Sometimes, as in that case, no symptoms are met with to direct attention to the belly. In other instances pain is complained of in the abdomen, but an examination of the part discovers no tenderness or tension. Often a certain fulness to the eye (a symptom of little value in the case of a child) is the only peculiarity to be detected. The general symptoms may be limited to feverishness and the vague indications of ill-health which are common to tuberculosis before local symptoms have had time to appear. Usually, however, secondary symptoms are not slow to arise, springing from the chest or the head, and showing clearly the tubercular nature of the case; but the abdominal lesion may remain latent throughout, and be only discovered when the body is examined after death.

When the abdomen is involved as part of an outbreak of general tuberculosis, the course of the peritoneal inflammation is always more rapid than it is in cases where the tubercular growth is limited—at any rate, primarily—to the peritoneum. In this latter form—chronic tubercular peritonitis—the course of the disease is usually very protracted, and if—as may happen—there are periods of improvement and relapse, many months may pass before the illness is at an end.

Chronic tubercular peritonitis is not a disease of early childhood. It is rarely seen in children under three years old, and not often at an earlier age than six or seven. The disease is of course a constitutional one, but the general symptoms which precede any manifestation of local distress are often curiously trifling and indefinite; the fever is moderate, the pallor slight, and the loss of flesh very gradual. Indeed, until peritoneal inflammation comes on to compli-

cate a case of abdominal tuberculosis, there is often little in the condition of the child to indicate the presence of serious disease.

The first symptom noticed is fulness of the abdomen, and there may be a little pain accompanied, perhaps, by slight tenderness. The pain occurs at first only on movement, so that the child neglects his usual games, and prefers sitting quietly to indulging in any bodily exertion. The tenderness is usually discovered by the child leaning his body against a table or chair, when he will say that the pressure hurts him. Often, however, tenderness is a later symptom, and towards the beginning of the illness handling of the abdomen elicits no expression of discomfort. At this time the appetite of the child is often good; there is no vomiting or other sign of digestive disturbance; and, except for some disinclination to exercise, the patient appears to be in his usual spirits. His general health is, however, evidently impaired. He wastes slowly but perceptibly, and the bodily heat is increased, the temperature rising every evening to 101° or 102° .

The pain after a time increases and becomes markedly paroxysmal, the tenderness is more pronounced, and the bowels are often capricious, so that attacks of diarrhœa alternate with constipation. The belly then begins to swell slightly, and palpation can sometimes detect a decided fluctuation. Before, however, this stage of the disease is reached, a considerable time may elapse. Slight pain, with a little tenderness, may be in existence for weeks, or even months, before further change occurs; and may even disappear altogether for a time, although the symptoms usually return after a longer or shorter interval.

The swelling of the abdomen is caused by an accumulation of gas. It is seldom the result of ascites, for the amount of fluid is rarely sufficient to cause distension of the belly. So far, indeed, from this being the case, fluctuation is not

unfrequently absent altogether, and we may have to make our diagnosis without it. Instead of fluctuation, there is in many cases a sense of vague impulse conveyed between the fingers placed on each side of the abdomen. The sensation is one which differs very much from the definite tap of a wave of fluid. It is an impulse transmitted by the coils of intestine more or less agglutinated by lymph, and holding a certain quantity of fluid between their folds. The presence of fluid can be often demonstrated by percussion. Thus, if fluid be present, there will be dulness in each flank as the child lies on his back; but when he is made to turn upon his side, the note produced by percussion upon the side turned uppermost after a few seconds becomes clear. Sometimes, when the quantity of fluid is small, fluctuation can be detected by placing the child on his elbows and knees and palpating the belly in this position. The fluid gravitates to the most depending part, and is thus brought within reach of the finger. It is only in exceptional cases that fluid is present in any quantity. It may, however, happen that there is copious ascites, and if at the same time tenderness of the abdomen is absent, the origin of the dropsy may be doubtful. A boy, aged seven years, was lately admitted into the East London Hospital for Children, suffering from ascites of four months' standing. The belly was large and globular, and measured thirty-two inches in girth and fourteen inches from the ensiform cartilage to the pubes. There was very little tenderness. Six and a half pints of greenish-yellow liquid were removed by paracentesis; and after the boy's death, which occurred a few days afterwards, several pints were found retained in the abdominal cavity. The cause of the effusion was tubercular peritonitis.

Another symptom generally present is a resistance of the abdominal wall to pressure from increased tension of the muscles. The tension is not a constant symptom, and is found to vary from time to time in degree. It is always

present when the belly is tender, but may sometimes be noticed in cases where no discomfort is complained of when pressure is made upon the abdomen. In rare cases this symptom, like the tenderness, is altogether absent, and the abdominal wall remains soft and yielding throughout.

Increased tension of the parietes is not the only cause of the resistance to pressure. As the deposit of lymph accumulates in the peritoneal cavity, and the coils of intestine become massed together and adherent, a sense of solidity is conveyed to the hand pressing upon the abdomen, which is very characteristic; and the more so that the resistance varies in different spots, being well marked in some places, while in others the parietes yield naturally to the pressure of the hand. This localised increase of resistance is one of the diagnostic tests of the disease. In certain cases hard masses of cheesy lymph can be felt through the parietes.

When the disease is confirmed, the shape of the belly undergoes a characteristic alteration; it is oval, with a plain uniform surface, which presents none of the slight prominences and markings natural to the abdominal wall in a state of health. If there be much distension, the superficial veins may become visible, especially if, as is often the case, the disease is accompanied by enlargement of the mesenteric glands. The pain when paroxysmal is usually severe, but it is not always a prominent symptom; and the amount of tenderness may be comparatively insignificant, and much less strongly marked than the nature of the complaint would lead us to expect. Sometimes neither pain nor tenderness are complained of throughout the course of the disease. In such cases, unless the abdomen be much distended, there may be nothing in the symptoms to draw attention to the belly, and, indeed, until signs arise indicating tubercular disease of some other part of the body, the condition of the child may be such that medical advice is considered unnecessary. Such an instance has already been related (see case of Maria H., page 247). Few cases,

however, occur where movements which give a sudden shock or jerk to the body are not a source of pain; and the child's own consciousness of this fact will induce a caution in his movements which is very peculiar. Thus, he will be noticed to go downstairs backwards so as to diminish the jerk; or to steady the abdomen with his hand as he walks, so as to control the vibrations produced in his body by the movement.

When the local symptoms become pronounced, the temperature rises, and the thermometer often marks 104° or even higher. At the same time the plumpness of the child rapidly diminishes, and the face assumes a pinched, worn expression, indicating great suffering, and highly characteristic of abdominal disease. He lies in bed with his knees drawn up, and movement of any kind seems to increase his distress. The pulse is rapid and weak; the lips dry, and often covered with sordes; diarrhoea is frequent; and death is usually preceded by extreme emaciation. The symptoms may remain throughout limited to the abdomen, or indications may be noticed showing that the disease is not confined to one cavity of the body. Cough and other symptoms arising from the chest may be observed; or death may be preceded by squinting or convulsions.

A very obstinate diarrhoea, with the passage of dark, watery, offensive stools, containing shreds of mucus and flaky matter, with small black clots of blood, is very suspicious of accompanying ulceration of the bowels.

The grey granulation is sometimes limited to the peritoneum lining the abdominal cavity or enveloping the contained organs; but often also it is found in the substance of the liver, kidneys, and spleen; and the mesenteric glands are usually enlarged and cheesy. In many of these cases the liver has been found to be in a state of acute cirrhosis.

CASE XXXIX.—TUBERCULAR PERITONITIS—APPARENT AMENDMENT—RELAPSE AND DEATH.

Selina L., aged seven years, was admitted into the East London Children's Hospital on January 14. The child had been well during the previous four years, but nothing could be discovered as to her health before that time, as her mother had long been dead.

Her present illness began about a fortnight before Christmas with pain in the region of the umbilicus, which was increased by coughing. There must also have been some tenderness of the abdomen, for she complained of being hurt by her sister who slept with her. About this time the belly looked large, and it has since increased in size. She has been losing flesh; has often complained of thirst; and her skin has been noticed to be hot. She has been rather languid and listless, and has neglected her usual amusements; but this was partly on account of the pain in the belly produced by exertion.

State on Admission.—Child in pretty good condition, not thin. Skin of body generally rather harsh, and of a peculiar sallow tint. Conjunctivæ slightly yellow. Belly very full, with free fluctuation all over; it does not seem to be tender, and she says she has now no pain. Liver does not project below the edges of the ribs. Behind, the liver dullness reaches to within a finger's breadth of the angle of the scapula. Spleen cannot be felt. Urine acid; specific gravity 1012; contains a slight trace of bile pigment; no albumen.

After she had been some days in the hospital her bowels became relaxed. The fluid then disappeared from the abdominal cavity and did not return. She remained an in-patient until February 22, for although she had ceased to have any abdominal symptoms, the temperature remained persistently high, varying from 100° to 101°. It was still

high when she left the hospital. In other respects she seemed well.

The patient was readmitted on April 16. She now appeared to be very ill. Her face looked pinched, and the skin was very hot. Temperature at 6 p.m., 104° . There was indistinct fluctuation of the abdomen, which was very tender, but was not especially full or otherwise altered in appearance. At the umbilicus a hard substance was felt, which appeared to be seated immediately beneath the abdominal wall. The induration was of about the size and shape of a crown-piece; the centre was at the navel, and the circumference was distinctly marked, especially at the upper part. It was extremely tender, and pressure upon it caused great pain. The child now began to waste very rapidly; her face became haggard, and her cheeks and temples hollow; the bowels also became relaxed, and her appetite failed. The temperature was always over 100° , varying from 100° to 104° . Her favourite position in the bed was on her back, inclined to the left side, with her knees drawn up. The abdomen remained very tender, and looked full, although the walls were flaccid. The indurated substance previously felt lost its circular shape, and became less distinctly circumscribed.

On May 6 the upper margin was distinctly to be felt immediately above the umbilicus, but laterally it faded gradually away, and was lost in general hardness along the sides of the belly, so that the impression produced to the finger was that of a straight line of induration passing outwards on each side from the umbilicus. Below this level the resistance of the abdominal wall was very much increased; the whole of the lower part of the belly gave more or less of a firm feeling to the touch, and was more tender than the upper part. No friction was heard when the ear was applied to the abdominal wall. On examination of the chest some want of tone was found at each base behind,

and some large-sized bubbling rhonchus was heard about both backs.

The child got rapidly weaker and more emaciated, the mouth and fauces became aphthous, and she died on May 18, the temperature shortly before death being 102.4° .

On post-mortem examination of the body the intestines were found matted together by recent lymph. The lymph deposit was especially abundant towards the central and lower parts of the belly, and was the cause of the hardness which had been noticed during life. The abdominal cavity contained about two ounces of dark yellow turbid fluid. On opening the bowel, several ulcers, circular or oval, and very cleanly cut, were found in the large intestine and ileum. At the upper part of the ileum was a spot where there was a collection of small masses varying in size from a peppercorn to a large pea. These masses were situated under the mucous membrane; they were yellowish-white and cheesy, and were evidently tubercles. The mucous membrane at this spot was not inflamed, but there were two or three small, cleanly cut ulcers with thickened edges. These had perforated quite through the coats of the bowel.

In other parts of the small intestine, so far at least as it could be followed out—for in places, on account of gluing of different portions of the bowel together by lymph, the canal could not be laid open—a few scattered tubercles, like those already described, were seen here and there.

Many cheesy glands of different sizes were found adherent to the intestine and scattered through the mesentery.

The liver was congested, and was firmly adherent to the diaphragm above, and to the kidney and stomach below. At its anterior border it presented several yellow projecting masses, varying in size from a pea to a marble. These showed a bright yellow granular surface on section, and were apparently softening. These masses were similar in appearance to those in the intestine.

The lungs were collapsed at their posterior surfaces, but contained no tubercles ; neither did the spleen or kidneys.

This case well illustrates the insidious beginning of peritoneal tuberculosis. There was fever, wasting, and languor, but the pain and tenderness were slight, and there was no vomiting or other sign of digestive disturbance. Effusion soon took place into the peritoneal cavity, but this disappeared after an attack of diarrhœa. The pain and tenderness also subsided, and, except for the continued pyrexia, the child seemed to be well. It was not until three months after the appearance of the first symptoms that the child appeared to be in any danger. At that time the inflammation returned with greater violence, the temperature became still more elevated, the pain and tenderness were severe, and an examination of the belly showed signs of extensive disorganisation.

The absence of vomiting is especially to be noticed. In simple peritonitis, the beginning of the inflammation is generally accompanied by much gastric disturbance ; but in cases where the inflammation occurs secondarily to tuberculosis, sickness is rarely met with, and, if present, is not troublesome, and occasions little inconvenience to the patient.

The frequent absence of signs of local suffering in the earlier stage of tubercular peritonitis is a point which it is necessary to insist upon.

Even when pain in the belly is complained of, it is often at first only an occasional symptom ; and tenderness of the same part, although distressing at one time, may be quite absent at another, and for days together may be unnoticed. Fulness of the belly is often the only local sign to be observed. With this, we may sometimes detect a slight increase of the abdominal tension even although the pressure of the hand is said to give no pain ; but increased tension of the abdominal wall, like pain and tenderness, is not a necessary consequence of peritoneal inflammation.

CASE XL.—TUBERCULAR PERITONITIS, ETC.

A girl, aged thirteen years, was admitted into the East London Children's Hospital on January 22. The child was one of twelve. One of her brothers was said to be the subject of phthisis, but there was no other history of this disease in the family, and the parents were reported to be healthy.

The girl had had measles and scarlatina in her infancy, but had never been considered delicate until the age of eleven years, when she had had an attack of what was called "acute rheumatism." Her illness was said to have lasted for three or four months, and she had since never completely recovered her strength. A little exertion tired her, and she had become habitually languid and dull. She complained occasionally of nausea after food and of vague pains about the belly.

The illness for which the girl was admitted was of six weeks standing. After a dinner of bullock's liver she was seized with severe pain in the abdomen, which appeared to be of a colicky character, for she was said to have been bent double by it. The pain lasted for four or five days. Ever since, the belly had been rather tender, and was thought to have increased in size. Now and then a little abdominal pain was complained of, but it was not severe. Since the beginning of her illness the girl had become very pale and weak, and had lost much flesh. She had passed the greater part of her time in bed, and had seemed incapable of any exertion. For three or four days she had coughed a little, and expectorated some frothy mucus.

On admission the girl was thin and pale, with hollow cheeks and an anxious look. The belly was a little full, but not at all tender. The edge of the liver could be felt just below the costal margin: the spleen could not be felt. No abnormal physical signs were found in the lungs, but a double superficial rub was detected at the base of the præcordial

region. The temperature was normal. Cod-liver oil and iron were prescribed, and the girl was ordered to take two ounces of port wine daily.

A few days later the temperature had risen to 101° . The patient expressed herself as quite free from pain, but the abdomen was full and distended, and seemed a little tender on pressure. The bowels acted two or three times a day, and the stools were loose and offensive.

On January 30, the abdomen looked large, but there was no fluctuation, no pain, nor any increase in the tension of the abdominal wall. The natural markings of the belly, however, were lost; there was slight tenderness, and, on pressure, a certain sense of solidity of the abdominal contents could be detected. The bowels still remained loose, and the girl admitted that she felt some occasional discomfort in the abdomen, although there was no actual pain. Her temperature was still 101° .

She was ordered to take a scruple of carbonate of bismuth, with three grains of pulvis cretæ comp. cum opio, every four hours.

On February 3 some dulness was noted in the right flank. There was a faint sense of fluctuation at that part, but none from side to side of the abdomen. The abdominal wall was slightly œdematous at the lower part. Temperature, 101° .

After this report the looseness of the bowels persisted. The temperature remained high and the child wasted rapidly. The abdomen became very tender, and the child began to complain of pains in the belly. The œdema of the abdominal wall increased and extended from the hypogastric region to the epigastrium. The distension of the abdomen was more marked at the lower than the upper half, and the skin below the navel became red. Free fluctuation was also noticed.

About the middle of February signs of fluid were discovered in the right chest. After this time the weakness

of the patient increased rapidly, although she continued to take food well. On February 21 she passed into a drowsy, half-delirious state, and died on the following day.

On examination of the body :—Grey granulations were found in the right pleura, and the lung on this side was compressed by a quantity of serous fluid. There were no tubercles in either lung. The abdomen was the seat of extensive peritonitis, which matted together the viscera and lined the whole cavity with lymph. Grey granulations were seen dotted over the intestines, the spleen, the gall bladder, and the liver. The latter lay in an encysted collection of clear fluid, which amounted to several pints. The omentum, mesentery and parietal peritoneum, were coated with a dense layer of thick lymph. The intestines contained a few tubercular ulcers.

The insidious way in which tubercular peritonitis may creep on is well illustrated by the above case. The child came of a family of evident consumptive tendencies, and entered the hospital with a history strongly suggestive of abdominal mischief. The suspicion thus aroused was supported by the aspect of the girl. She was pale, with a spare frame ; her expression was anxious, and her face looked haggard. She had, indeed, a distinctly tubercular appearance. Still, the closest examination failed to discover sufficient grounds to warrant the diagnosis of tuberculosis. No abnormal physical signs were noted about the lungs. The abdomen, although a little full, was not tender, and there was no enlargement of the liver or spleen. It was only gradually that signs of peritoneal inflammation became developed ; but eventually there was great distension and tenderness of the belly ; free fluctuation was noticed ; signs of effusion into the chest were observed ; the child wasted rapidly, and died, apparently from exhaustion. This mode of death is the less common termination of general tuberculosis in the child. It is not often in young subjects that the cerebral meninges escape ; indeed, in the majority of cases

the illness closes with all the symptoms of the third stage of tubercular meningitis.

The frequent absence of pain and tenderness in the earlier period of tubercular peritonitis is a fact of great clinical importance. In cases where the peritoneal inflammation, although tubercular, is acute, the absence of pain and discomfort is, according to my experience, the rule. Even the normal tension of the parietes may not be increased. It is important to remember this fact, so that when the general symptoms point to tubercular mischief, we may not exclude peritonitis because we can press upon the belly without provoking a cry. It is a good rule, indeed, to take the contrary course, and in every case where tuberculosis is probable, to suspect the abdominal complication whenever the abdomen is continuously full, even although there be no tenderness, no pain, no abnormal tension of the parietes, no fluctuation, nor any increase in the resistance of the abdominal contents.

On account of this obscurity of the earlier symptoms, tubercular peritonitis is frequently overlooked at its beginning. Abdominal pain is common enough in children, and is usually at once attributed to digestive derangement. It is well, however, in all such cases, to make special examination of the belly; for although the symptom is probably due to indigestion of food, it may arise from a more serious cause; and, should the child afterwards succumb to an attack of tubercular peritonitis, the failure to recognize the early signs of so grave a disease is a mistake which the friends of the patient are not likely readily to forgive.

When a careful examination of the abdomen detects the presence of evident tenderness, the symptom is a very important one. Flatulent pains are not increased by firm, steady pressure: and if, while diverting the attention of the child, we find that such pressure causes a cry of distress, we should look for some other cause of the discomfort than mere flatulent spasm. Again, a persistent temperature of over 100° is not produced by a chronic indigestion;

and whenever associated with abdominal tenderness and pain, forms a very suspicious combination.

We must not, however, in cases where this group of symptoms is complained of, allow our attention to be directed too exclusively to the abdomen, or we may overlook the existence of a pleurisy. The pain of pleuritic inflammation is often felt in the belly, where there may even appear to be some tenderness, and the temperature is generally elevated. An examination of the chest must, therefore, on no account be omitted.

Persistent abdominal pain is no evidence of tubercular disease, so long as there is no tenderness and the temperature is natural. Such pains are of common occurrence. The following case is the type of a class very frequently to be met with. The symptoms are due merely to a functional derangement, and have no gravity whatever, but they often persist for months together if not treated judiciously.

A little girl, aged eight years, who had cut twelve of her second crop of teeth, was brought to me with the following symptoms:—For some weeks, although apparently in her usual spirits, she had been losing flesh and had complained of pain in the belly, coming on at different times in the day and sometimes in the night. It had been noticed that during the day she often complained shortly after taking food. The pains were seated towards the lower part of the abdomen, and appeared to be griping in character; but the child did not attempt to relieve them by bending the body as in ordinary flatulent pains. They were not accompanied by any feeling of sickness, but when severe, took all colour from her cheeks. The girl's appetite was good, but not excessive; her bowels regular, but inclined to be costive, and she sometimes strained at stool. At night she was restless and her sleep was disturbed; but she had no worms, nor had mucus been noticed in the evacuations. Her tongue was clean; her complexion rather pale, but not at all sallow; and there was no yellowness

of the conjunctivæ. There was no sign of distress in the face.

On examination, the abdomen was found to be natural, without tenderness or increase of tension, and the child's temperature was that of health. She had been treated with aperients, but with little benefit. This patient was very quickly cured by an alkaline mixture, with a bi-weekly dose of syrup of senna, and the use of a flannel bandage to the belly. The parents, too, were directed to be sparing for a time in the amount of farinaceous matter allowed in her diet. Potatos, in particular, were strictly forbidden.

Tenderness of the abdomen is, then, an important element in the diagnosis of tubercular peritonitis. If in a child who complains frequently of pains in the abdomen, there is general tenderness and fulness of the belly, and especially if the temperature of the body be higher than that of health, we should suspect the beginning of tubercular peritonitis. The tenderness, when it is slight, is often elicited by slight jerks or shocks, and these cause distress to the patient out of all proportion to their apparent violence. Therefore, caution in the movements of a child—so contrary to childish habits—should never be disregarded ; and if the patient be seen to steady his abdomen with his hand as he walks, the sign is a very suspicious one.

Although pain and tenderness, when present, are important symptoms, it must not be forgotten that the disease may exist without either. On the other hand, slight pain and tenderness, even of long duration, are not to be taken as an indication that the case must necessarily be one of tubercular peritonitis. A child who appears to be weakly and has a distressed face is said to be subject to frequent attacks of diarrhœa. In the intervals between the attacks his bowels tend to be costive. He often complains of colicky pains in the belly, and these are apt to come on at about the same hour every day. There is slight tenderness on pressure

of the abdomen, with some increase in the tension of the parietes. The stools are either offensive and watery, with a shreddy deposit, which may contain little black clots; or semi-solid, scanty, with stringy mucus and perhaps streaks of blood. These symptoms, which are due to abrasions or ulceration of the intestinal mucous membrane, are distinguished from those set up by peritonitis by noticing that the belly is not unduly distended, and has not the globular shape common to serous inflammation; that the natural markings of the surface are not obliterated; that the abdomen moves freely in respiration; and that the temperature of the body is not raised. Moreover, in cases of uncomplicated ulceration of the bowels there is no fluctuation. It must be remembered that ulceration of the bowels is not seldom a complication of peritonitis, and that in such cases the intestinal lesion may be suspected from the persistence of the purging and the appearance of the stools, which have the character above referred to. In the case (XL.) last related, the mere obstinacy of the purging was in itself a strong argument in favour of ulceration of the intestinal mucous membrane.

It is only during the earlier period of the disease that there is any cause for uncertainty as to the nature of the illness. At a later stage, when the symptoms have become pronounced, we can scarcely overlook the presence of peritonitis; and we infer that the inflammation is of a tubercular character from the family history of the patient, and, especially, the chronic course of the illness. For all practical purposes a chronic peritonitis in the child may be taken to be of tubercular origin.

In the treatment of acute general tuberculosis, little, unfortunately, can be done after the disease is fully established. It is only while the child is as yet free from the complaint, but when from family history or other cause we have reason to apprehend its occurrence, that we are able to do much in preventing the development of this fatal illness.

If other children of the same parents have died of acute

tuberculosis, very careful attention should be paid to the diet and general management of those who are left. A judiciously arranged dietary—avoiding excess of farinaceous food, on account of its tendency to promote acid dyspepsia,—regular meals, early hours, airy bed-rooms, plenty of fresh air and exercise, and plenty of sleep, are all matters to which our care should be directed.

Such children are often very active, mentally as well as physically, and it is important that the brain should not be unduly exercised or excited. A mid-day sleep should be enforced as long as possible, as, by this means, the brain is insured a necessary amount of repose. It is well, also, in such cases not to be too precipitate in forwarding the mental culture of the child. The lessons at first should be easy and short, and every care should be taken to avoid undue strain upon the faculties, so that the nervous system may not be overtasked.

In the selection of a residence, attention should be directed to the character of the soil: preference should be given to sand or gravel; and low-lying clay lands ought, if possible, to be avoided.

An infant may be nursed by the mother if she herself be healthy and her milk suitable to the child; if not, a good wet-nurse should if possible be provided.

All digestive derangements should receive early attention, and in particular looseness of the bowels, however apparently trifling it may be, must never be neglected, as no functional disorders to which children are liable have a more immediate and depressing influence upon the processes of nutrition. Inflammatory chest affections also should be treated with the greatest care, and in the case of pleurisy early paracentesis is advisable, if at the end of three weeks from the beginning of the disease no signs of removal of the effused fluid have been noticed. The outbreak of any of the acute specific diseases should be regarded with very great anxiety, and for some time afterwards careful watch

should be kept over the patient, so that the occurrence of any subsequent symptoms may be met by suitable treatment. In cases where the means of the parents permit the expense, change of air for a time to a dry and bracing but sheltered spot is of great value, and will do more than any amount of expenditure in drugs to remove the risk of after ill-consequences. This is especially true in the case of measles and whooping-cough.

When the disease has actually declared itself, no drugs appear to have any influence upon its progress, or any power in preventing a fatal termination. We are obliged to content ourselves with the treatment of symptoms and of the secondary complications to which the disease almost invariably gives rise. It is highly desirable to reduce the pyrexia, as it is to this that the wasting is principally due. But, unfortunately, remedies which are found to lower the temperature in inflammatory chest affections have but slight value in cases where the fever is symptomatic of the formation of the grey granulation. Quinine and digitalis—drugs which are so useful in pneumonic phthisis—are here of little avail; and the hypophosphites, whose influence in the same cases appears sometimes to be almost magical, are of no service in acute tuberculosis. Iron and tonics generally appear to have no other effect than that of increasing the patient's discomfort; while cod-liver oil is very apt to derange the stomach, and if it agrees is seldom of any benefit. I have never seen a case of acute tuberculosis arrested or even retarded by cod-liver oil. We can do little more than place the child in an airy room, supply him with a diet regulated with due regard to his powers of digestion and the irritable condition of his bowels, order a mild saline, such as citrate of potash, and wait for special symptoms to declare themselves.

The diarrhoea, which is so common in this disease, is not always readily arrested. During the earlier period of the illness, when the looseness probably depends upon mere

functional derangement, it is amenable to the milder astringents, and may be stopped by a few doses of aromatic sulphuric acid with a small quantity of laudanum; but afterwards, when inflammation or ulceration of the intestinal mucous membrane is present, stronger measures are required, and we must have recourse to lead and opium, gallic acid, and other remedies, such as are found useful in an adult suffering from the same complication.

Inflammatory chest affections, when they occur secondarily to tubercle, must be treated in exactly the same way as if the complication were the primary disease; but if we are satisfied of the correctness of our diagnosis, we can have little hope that treatment will have any influence in retarding the fatal issue.

When the disease assumes the form of tubercular meningitis, care should be taken to keep the head cool and the feet warm, and to act upon the bowels by suitable aperients. The head must be shaved, and a bladder of ice kept constantly applied to it. The ice should be broken up into small pieces, and all air should be squeezed out of the bladder before closing it up, so that the latter may fit like a cap upon the shaven scalp. Other applications to the head, such as leeches or counter-irritation by blisters or irritating ointments, appear to be actually injurious, and to have no other effect than that of hastening the fatal termination. If the feet are cold they should be warmed by hot bottles. The diet of the child should consist of beef-tea, milk, yolk of egg, and, as he becomes weaker, the brandy and egg mixture.

The best aperient is a combination of calomel and jalap, and the action of the bowels may be assisted by enemata. The only other drugs from which any good ever seems to result are the iodide of potassium and perchloride of mercury. Recovery has in some cases been known to follow the employment of the iodide in large doses; but such cases in all probability were not cases of true tuber-

cular meningitis. A simple meningitis at the base of the brain might present symptoms differing little from the tubercular form of the disease, and there is reason to think that a syphilitic inflammation of the cerebral membranes is occasionally mistaken for tubercular meningitis.

In tuberculosis of the peritoneum, when pain and tenderness are complained of, the child should be at once confined to his bed, a succession of hot linseed-meal poultices should be applied to the abdomen, and perfect quiet should be enforced. If the pain be very severe at any particular point of the belly, a leech or two may be applied; but care must be taken to arrest the bleeding early, so that the strength of the child may not be reduced. Diarrhoea must be treated by opium and astringents, and the diet should consist principally of milk and broths. If the case be seen early, this treatment may have the effect of quieting the symptoms for a time, so that the child may appear to recover. When the case is more advanced and the symptoms are severe, with rapid wasting and high fever, the condition of the child is hopeless. Little more can be done than to endeavour to relieve the pain by hot applications and the administration of opium by the mouth or by hypodermic injection. When the strength begins to fail, brandy and egg mixture should be prescribed.

INDEX.

A.

	PAGE
ABDOMINAL wall, tension of, in peritonitis	105
Acute disease in wasted infants	5
„ general tuberculosiſ. See <i>Tuberculosis</i> .	
„ tuberculosiſ, ſecondary, in chronic catarrhal pneumonia .	178
„ „ „ pleuriſy	98, 103
Ægophony in pleuritic effuſion	110
Albuminuria in acute tuberculoſiſ	244
Alkalies, value of, for infants	17
„ „ in chronic catarrhal pneumonia	189
Amphoric breathing in eaſe of dilated bronchus	218
Amyloid degeneration, a reſult of empyema	105
„ „ „ cirrhoſiſ	224
Anæmia induced by mercury	19
Antipyretics in erupous pneumonia	69
Aperients in acute cheſt diſeaſes, care in ordering	118
Aromatics, value of, in treatment	18
Arsenic, in phthiſiſ	189
„ toleranee of children for	19
„ to check vomiting in cirrhoſiſ of lung	236
Aſciteſ in tubercular peritonitiſ.	289
Aſtringents in cirrhoſiſ	235
„ tuberculous diarrhœa	305
Atelectaſiſ, poſtnatal. See <i>Lung, Collapse of</i> .	

B.

Bandage, flannel, in diaphragmatic pleuriſy	117
Baſic lung diſeaſe from pleuriſy	104
Belladonna for night ſweats	191
„ toleranee of children for	18
Blood, ſpitting of, from epiſtaxiſ	168
„ „ in fibroid induration of lung	205
Bone diſeaſe a cauſe of acute tuberculoſiſ	251
Brain, tumour of	284
„ abſceſſ of	284

	PAGE
Breathing, laborious, in acute catarrhal pneumonia	130
" rapidity of, in acute tuberculosis	246
" " croupous pneumonia	45
" " catarrhal pneumonia	130
" " collapse of lung	26
" " infants	8
" " pleurisy	74
" shallow, in pleurisy	74
" " collapse of lung	26
Breath sounds, peculiarities of, in children	161
Bright's disease, diagnosis from acute tuberculosis	253
Bronchi, dilatation of. See <i>Dilated Bronchi</i> .	
" ulceration of.	194
Bronchial catarrh, dangerous in wasted infants	40
" " treatment of	149
" glands, conducting power of	161
Bronchiectasis. See <i>Dilated Bronchi</i> .	
Bronchitis, tubercular	246
Broncho-pneumonia. See <i>Catarrhal Pneumonia, Acute</i> .	
Bulging of chest in pleurisy	77

C.

Caseous deposits	156
" " absorption of	157, 164
" " a cause of tuberculosis	159
" " causes of	154
" " seat of, in lung	165
" " softening of	157
" " treatment of	188
Catarrh, bronchial, a great danger in wasted infants	40
" " treatment of	149
Catarrhal pneumonia, acute	126
" " causes of	128
" " diagnosis of	146
" " " from lobular collapse	147
" " dilatation of bronchi in	143
" " distinction between, and croupous pneumonia	127
" " favourable ending of	139
" " impassiveness in	135
" " pathology of	127
" " physical signs of	131
" " subacute type of	135
" " symptoms of	130
" " treatment of	149
Catarrhal pneumonia, chronic	154
" " cavernous breathing in	166, 168
" " cavities of lung in	158
" " diagnosis of	183
" " dilatation of bronchi in	166, 173
" " hæmoptysis in	168
" " pathology of	155

	PAGE
Catarrhal pneumonia, chronic, physical signs of	162
" " prognosis in	186
" " softening in	157
" " symptoms of	160
" " treatment of	187
Cavernous breathing in chronic catarrhal pneumonia	166, 168
" " pleurisy	80, 110, 168
" " not always the sign of a cavity	168
Cavities in the lung	158
" " diagnosis of	166
" " physical signs of	166, 167
Cerebral abscess	284
" pneumonia	52
" tumour	284
Chest, distortion of, in cirrhosis	212, 216
" " pleurisy	77, 97
" " examination of	9
" " shape of, in pleurisy	77, 97
Children, clinical examination of	6
" constitutional peculiarities of	2
" respiration in	8
" temperature in	7
Chloralum for washing out chest in empyema	124
Circumscribed pleurisy	83, 112
Clinical examination of infants	6
Collapse of lung. See <i>Lung, Collapse of.</i>	
Complexion in acute tuberculosis	242
" " pleurisy	76, 111
Convulsions in cerebral tumour	284
" collapse of lung	26
" pneumonia	50
" simple meningitis	283
" tubercular meningitis	265
" tuberculosis	258
Cough, paroxysmal, in cirrhosis of lung	205
Counter-irritation in pleurisy	118
" " pulmonary catarrh	149
Crisis in croupous pneumonia	49
Croupous pneumonia	43
" " convulsions in	50
" " delirium in	44, 51, 65
" " diagnosis of	64
" " " from pleurisy	109
" " epistaxis in	51
" " gastro-intestinal catarrh in	57, 60
" " physical signs of	46
" " prognosis in	63
" " resolution in	49
" " symptoms of	44
" " temperature in	44
" " " reduction of	69
Cyrtometer, use of, in pleurisy	77

D.

	PAGE
Death, sudden, in pulmonary collapse	26
Deformity of chest in pleurisy	77, 97
Delirium in croupous pneumonia	44, 51, 65
Dentition, temperature in	7, 184
" wasting from	243
Diagnosis, general considerations	11
" of abscess of brain	284
" acute catarrhal pneumonia	146
" acute tuberculosis	254
" cerebral tumour	284
" chronic catarrhal pneumonia	184
" collapse of lung	37
" dilated bronchi	177
" fibroid induration of lung	109, 232
" hydrothorax	10, 115
" pleuritic effusion	108
" pleurisy with retraction	215
" pneumonia, croupous	64
" pulmonary tuberculosis	185, 244
" simple meningitis	283
" tubercular meningitis	282
" " peritonitis	299
" typhoid fever	59, 244
Diarrhœa in acute tuberculosis	244
" chronic catarrhal pneumonia	172
" tubercular peritonitis	291
Diet, dry, in pleurisy	118
" in acute catarrhal pneumonia	152
" acute general tuberculosis	303
" chronic catarrhal pneumonia	188
" collapse of lung	42
" croupous pneumonia	68
" empyema	125
" pleurisy	117
Dilated bronchi, cough in	205
" " diagnosis of	177
" " in acute catarrhal pneumonia	128, 143
" " chronic catarrhal pneumonia	166, 173, 177
" " fibroid induration of lung	193, 201, 224
" " retention of sputa in	193, 205
Displacement of heart in cirrhosis	205
" " pleurisy	77
Drainage-tubes in empyema	123
Dulness, quality of, in collapse of lung	38
" " fibroid induration of lung	205
" " pleurisy	78

E.

	PAGE
Empyema, absorption in	93
„ a cause of amyloid disease	105
„ „ tuberculosis	94
„ diagnosis of	113, 185
„ diet in	125
„ often mistaken for phthisis	108, 113
„ paracentesis in	119
„ prognosis in	115
„ results of	94
„ spontaneous evacuation of fluid in	94
„ symptoms of	92
„ temperature in	93
„ treatment of	119
Epistaxis in acute tuberculosis	245
„ children	168
„ pneumonia	51
Examination, clinical, of infants	6
„ of the chest in children	10, 160
„ of sputum	206
Excavation of lung	158, 167
„ „ in chronic catarrhal pneumonia	166
„ „ in fibroid induration	225
Expectorants, relative value of	150, 190

F.

Fever, remittent in acute tuberculosis	243
„ typhoid, diagnosis from acute tuberculosis	256
„ „ „ croupous pneumonia	59
„ „ facies in	257
„ „ spleen in	256
„ „ temperature in	256
Fibrinous pleurisy	104, 182
Fibroid induration of lung	192
„ „ „ causes	193
„ „ „ diagnosis	232
„ „ „ pathology	192
„ „ „ physical signs	205
„ „ „ prognosis	233
„ „ „ symptoms	205
„ „ „ temperature in	206
„ „ „ treatment of	235
„ phthisis	194
„ „ case of	227
Finger-ends, clubbing of, in cirrhosis	205
Fœtor of pus in empyema, to correct	124
„ sputum in cirrhosis of lung	205
Fontanelle in tubercular meningitis	265
Friction, pleural, peculiarities of, in children	79
Functional derangements, frequency of, in infants	4

G.

	PAGE
Gastric catarrh during dentition	255
Glands, bronchial, enlarged, conducting power of	161
Granulations, grey	239
" " quickly become yellow	239

H.

Hæmoptysis in children	168
Headache in tubercular meningitis	262, 267
Heat, reduction of, in croupous pneumonia	69
Heart, displacement of, in fibroid induration of the lung	205
" pleurisy	77
"Hydrocephalic" cry	263
Hydrocephalus, acute. See <i>Meningitis, Tubercular</i> .	
Hydrothorax, diagnosis of	115
Hypophosphites, value of the	190, 304

I.

Inequality of pupils in tubercular meningitis	263, 264
Infant, case of acute catarrhal pneumonia in an	131
Infants, constitutional peculiarities of	2
" examination of chest in	10, 161
" pulse of	7
" respiration of	8
" to take temperature of	7
Inhalations in cirrhosis of lung	235
Invasion stage of tubercular meningitis	261
Iodine liniment for phthisis	189
" pleurisy	118
Iron, large doses of, in cirrhosis of lung	235

K.

Kidney, amyloid degeneration of, in cirrhosis of lung	230
---	-----

L.

Laryngismus stridulus	4
" " complicating pulmonary collapse	32
Liver, enlarged, diagnosis of, from pleurisy	114
Lobar pneumonia. See <i>Croupous Pneumonia</i> .	
Lobular pneumonia. See <i>Catarrhal Pneumonia, Acute</i> .	
Localised pleurisy, diagnosis of	112
" " symptoms of	85

	PAGE
Lung, collapse of	21
" " anatomical characters of	25
" " causes of	22
" " diagnosis of	37
" " danger of, in young babies	40
" " in pertussis	31
" " in phthisis	163
" " mechanism of	21
" " physical signs of	27
" " sudden death in	26, 28
" " symptoms of	26
" " treatment of	41
Lungs, often small in phthisical children	160

M.

Measles, a cause of catarrhal pneumonia	172, 250
" " tuberculosis	172, 250
" sequelæ of	250
Meningitis, simple	283
" " duration of	284
" tubercular	259
" " constipation in	261, 280
" " diagnosis of	282
" " duration of	267
" " fontanelle in	268
" " headache in	262, 267
" " invasion of	261
" " of convexity	276
" " ordinary attack of	267
" " paralysis in	264
" " premonitory symptoms of	260
" " pulse in	262, 266
" " second stage of	262
" " secondary	271
" " " cases of	271, 274
" " squint in	263, 264
" " stages of	261
" " symptoms of	260
" " temperature in	262, 265
" " third stage of	263
" " treatment of	305
" " vomiting in	261
Mercury, effects of, on children	19
Mouth, examination of	11
" open, during examination of chest	11, 151

N.

Necrosis of bone a cause of tuberculosis	251
Nervous symptoms, in tubercular meningitis, value of	281

	PAGE
Nervous system, excitability of	2
" " " often lost in chronic disease	5
Nose, bleeding of, in croupous pneumonia	51
Nutrition in chronic catarrhal pneumonia	162
" fibroid induration of the lung	206

O.

Œdema in acute tuberculosis	244
Otorrhœa, a cause of cerebral abscess	285

P.

Paracentesis, mode of performing	120
Paralysis in cerebral tumour	284
" tubercular meningitis	264
Pars petrosa, disease of	285
Percussion note, quality of, in collapse of lung	38
" " pleurisy	70, 78
" with two fingers, importance of	10
Peritonitis by extension of inflammation from the pleura	105
" latent	108
" tubercular	286
" " cases of	292, 296
" " diagnosis of	299
" " insidious beginning of	287
" " symptoms of	288
" " temperature in	288, 291
" " treatment of	306
Pertussis a cause of catarrhal pneumonia	129
" " pulmonary collapse	31
" " tuberculosis	240
Phthisis, acute pulmonary	157
" chronic pulmonary	154, 160
" common form of	160
" definition of	160
" diagnosis of, from pleurisy	113, 185
Physical signs in acute catarrhal pneumonia	131
" " pulmonary tuberculosis	246
" " chronic catarrhal pneumonia	163
" " collapse of lung	27
" " croupous pneumonia	46
" " empyema	76
" " fibroid induration of lung	205
" " pleurisy	76
Pleural friction in children	79
Pleurisy	72
" localised	85
" deformity of chest from	97
" diagnosis of	108

	PAGE
Pleurisy, diet in	117, 125
" effect of, in causing absorption of pneumonic deposits.	89
" fibrinous	104, 182
" paracentesis in	119
" physical signs of	76
" prognosis in	115
" retraction of side in	97
" secondary tuberculosis from	98
" symptoms of	73
" termination of	89
" treatment of	116
Pneumonia, catarrhal. See <i>Catarrhal Pneumonia</i> .	
" cerebral	52
" croupous. See <i>Croupous Pneumonia</i> .	
" of apex of lung	56
" " rapid resolution of	56
" secondary	60
" " treatment of	71
Pneumothorax, traumatic	121
Prognosis, general considerations on	14
Pulmonary collapse. See <i>Collapse of Lung</i> .	
" disease, peculiarities of, in children	160
" tuberculosis. See <i>Tuberculosis</i> .	
Pulse in collapse of the lung	26
" infants	7
" " precautions in counting	7
" tubercular meningitis	251
Pulse-respiration ratio in catarrhal pneumonia	130
" " collapse of lung	26
" " croupous pneumonia	45
" " pleurisy	74
Pupils in tubercular meningitis	263, 264
" inequality of, as a sign of cerebral disease	279
Pyrexia from teething	7
" remittent, in tuberculosis	243

Q.

Quality of voice in inherited syphilis	8
Quinine in croupous pneumonia	70
" pleurisy "	119

R.

Resolution of croupous pneumonia	48
Respiration, rapidity of, in acute catarrhal pneumonia	130
" " " tuberculosis	246
" " children	8
" " collapse of lung	26

	PAGE
Respiration, rapidity of, in croupous pneumonia	45
" " fibroid induration of lung	205
" " pleurisy	74
" " tubercular meningitis	263
" " unequal in tubercular meningitis	252, 263, 279
Rickets a cause of collapse of lung	23, 33
Rigidity of joints in tubercular meningitis	264

S.

Scapula, tilting backwards of, in pleurisy	98
Secondary tuberculosis from chronic catarrhal pneumonia	178
" " from pleurisy	98
Sighing in tubercular meningitis	263
" value of, in diagnosis of tubercular meningitis	281
Sleep, caution concerning, in collapse of lung	42
" irregularity of pulse in	8
Softening in phthisis, symptoms	165
Soil, dryness of, important for phthisical patients	188
Spleen, enlargement of, in typhoid fever	256
" " tuberculosis	253, 256
Sputa, elastic tissue in	167, 206
" factor of, in cirrhosis of lung	205
" microscopical examination of	206
Stethoscope in the examination of infants	10
Stimulants in the treatment of disease	16
" the best for children	191
Subacute type of catarrhal pneumonia	135
Swallowing, difficulty of, in collapse of lung	27
Symptoms, local, value of	3
" " sometimes absent	3
" " " deceptive	4
" of abscess of brain	284
" acute catarrhal pneumonia	130
" " general tuberculosis	245
" " meningitis	283
" cerebral tumour	272, 284
" chronic catarrhal pneumonia	160
" collapse of lung	26
" croupous pneumonia	44
" emphyema	92
" fibroid induration of lung	205
" pleurisy	73
" pulmonary tuberculosis	245
" softening of caseous mass	165
" tubercular meningitis	260
" " peritonitis	288

T.

	PAGE
Tâche cerebrale	252, 254
Temperature in acute catarrhal pneumonia	130
" acute general tuberculosis	243
" children	7
" chronic catarrhal pneumonia	163
" collapse of lung	26
" croupous pneumonia	44
" empyema	93
" fibroid induration of lung	206
" pleurisy	75
" tubercular meningitis	262, 265
" " peritonitis	288, 291
Tint of face in pleurisy	76
Treatment of acute catarrhal pneumonia	152
" " general tuberculosis	302
" chronic catarrhal pneumonia	187
" collapse of lung	41
" croupous pneumonia	67
" empyema	120
" fibroid induration of lung	235
" general considerations on	14
" pleurisy	116
" pulmonary tuberculosis	289, 305
" tubercular meningitis	305
" " peritonitis	306
Traumatic pneumo-thorax, ease of	121
Tubercular infiltration of Laennec	156
" meningitis. See <i>Meningitis</i> .	
" peritonitis. See <i>Peritonitis</i> .	
Tuberculosis, acute general	237
" albuminuria in	244
" diagnosis of	254
" emaciation in	243
" etiology of	240
" expression of face in	243, 257
" œdema of limbs in	244
" pathology of	238
" symptoms of	242
" temperature in	243
" treatment of	302
" urine in	244
Tuberculosis, pulmonary	245
" breathing in	246
" ease of	247
" cough in	245
" diagnosis of	185, 257
" secondary complications	247, 258
" symptoms of	245
" treatment of	302
" secondary, to catarrhal pneumonia	178
" " pleurisy	98

U.

	PAGE
Ulceration of bowels in pulmonary phthisis	159
" " tubercular peritonitis	291, 302
" " symptoms of	302
" " bronehi	194
" " temperature in	206
" " cornea in tubercular meningitis	266
Urine in acute tuberculosis	244
" purulent, in a case of eirrrosis of the lung	228

V.

Varicella, an exciting cause of acute tuberculosis	241
Veins, distension of superficial, in acute catarrhal pneumonia	130
" " " fibroid induration of the lung	206
Vocal fremitus in eroupous pneumonia	66
" " collapse of the lung	28
" " fibroid induration of the lung	204
" " pleurisy	78, 100, 110
" " uncertainty of, in children	66
Vocal resonance in children	161
" " chronic catarrhal pneumonia	155
" " collapse of the lung	28
" " croupous pneumonia	47
" " fibroid induration of lung	205
" " pleurisy	80
Voice in inherited syphilis	8
Vomiting in fibroid induration of lung	205
" " " to check	236
" " tubercular meningitis	261
" " rarity of, in tubercular peritonitis	288

W.

Wasting in acute tuberculosis	242
" " chronic catarrhal pneumonia	162
" " fibroid induration of lung	206
Weak breathing, an early sign of pleurisy	79
" " not always a sign of disease	162
Weakly infants, danger of pulmonary collapse in	5
" " case of pulmonary collapse in	28
Whooping-cough, a cause of catarrhal pneumonia	129
" " " pulmonary collapse	31

BY THE SAME AUTHOR.

A PRACTICAL TREATISE ON DISEASE IN
CHILDREN. 8vo., 22s.

ON THE WASTING DISEASES OF INFANTS AND
CHILDREN. Fourth Edition, post 8vo., 8s. 6*d*.

LONDON: J. & A. CHURCHILL.

J. & A. CHURCHILL'S
MEDICAL CLASS BOOKS.

ANATOMY.

BRAUNE.—An Atlas of Topographical Anatomy, after Plane Sections of Frozen Bodies. By WILHELM BRAUNE, Professor of Anatomy in the University of Leipzig. Translated by EDWARD BELLAMY, F.R.C.S., and Member of the Board of Examiners; Surgeon to Charing Cross Hospital, and Lecturer on Anatomy in its School. With 34 Photo-lithographic Plates and 46 Woodcuts. Large Imp. 8vo, 40s.

FLOWER.—Diagrams of the Nerves of the Human Body, exhibiting their Origin, Divisions, and Connexions, with their Distribution to the various Regions of the Cutaneous Surface, and to all the Muscles. By WILLIAM H. FLOWER, F.R.C.S., F.R.S. Third Edition, containing 6 Plates. Royal 4to, 12s.

GODLEE.—An Atlas of Human Anatomy: illustrating most of the ordinary Dissections and many not usually practised by the Student. By RICKMAN J. GODLEE, M.S., F.R.C.S., Assistant-Surgeon to University College Hospital, and Senior Demonstrator of Anatomy in University College. With 48 Imp. 4to Coloured Plates, containing 112 Figures, and a Volume of Explanatory Text, with many Engravings. 8vo, £4 14s. 6d.

HEATH.—Practical Anatomy: a Manual of Dissections. By CHRISTOPHER HEATH, F.R.C.S., Holme Professor of Clinical Surgery in University College and Surgeon to the Hospital. Sixth Edition, revised by RICKMAN J. GODLEE, M.S. Lond., F.R.C.S., Demonstrator of Anatomy in University College, and Assistant Surgeon to the Hospital. With 24 Coloured Plates and 274 Engravings. Crown 8vo, 15s.

11, NEW BURLINGTON STREET.

CHEMISTRY—*continued.*

CLOWES.—**Practical Chemistry and Qualitative Inorganic Analysis.** Adapted for use in the Laboratories of Schools and Colleges. By FRANK CLOWES, D.Sc. Lond., Professor of Chemistry in University College, Nottingham. Fourth Edition. With Engravings. Post 8vo, 7s. 6d.

FOWNEs.—**Manual of Chemistry.**—*See WATTS.*

FRANKLAND AND JAPP.—**Inorganic Chemistry.** By EDWARD FRANKLAND, Ph.D., D.C.L., F.R.S., and F. R. JAPP, M.A., Ph.D., F.I.C. With 2 Lithographic Plates and numerous Wood Engravings. 8vo, 24s.

JAMES.—**Notes on the Detection of the Acids** (Inorganic and Organic) usually met with in Analysis. For the use of Laboratory Students. By J. WILLIAM JAMES, Ph.D., F.C.S., Demonstrator and Assistant Lecturer in University College, Cardiff. 8vo, 1s.

MORLEY.—**Outlines of Organic Chemistry.** By H. FORSTER MORLEY, M.A., D.Sc., Assistant Professor of Chemistry at University College, London. Crown 8vo, 7s. 6d.

TIDY.—**A Handbook of Modern Chemistry,** Inorganic and Organic. By C. MEYMOTT TIDY, M.B., Professor of Chemistry and Medical Jurisprudence at the London Hospital. 8vo, 16s.

VACHER.—**A Primer of Chemistry, including** Analysis. By ARTHUR VACHER. 18mo, 1s.

VALENTIN.—**Chemical Tables for the Lecture-room and Laboratory.** By WILLIAM G. VALENTIN, F.C.S. In Five large Sheets, 5s. 6d.

VALENTIN AND HODGKINSON.—**A Course of Qualitative Chemical Analysis.** By W. G. VALENTIN, F.C.S. Sixth Edition by W. R. HODGKINSON, Ph.D. (Wurzburg), Senior Demonstrator of Practical Chemistry in the Science Schools, South Kensington, and H. M. CHAPMAN, Assistant Demonstrator. With Engravings and Map of Spectra. 8vo, 8s. 6d.

The Tables for the Qualitative Analysis of Simple and Compound Substances, with Map of Spectra, printed separately on indestructible paper. 8vo, 2s. 6d.

11, *NEW BURLINGTON STREET.*

CHEMISTRY—*continued.*

WATTS.—Physical and Inorganic Chemistry.

By HENRY WATTS, B.A., F.R.S. (being Vol. I. of the Thirteenth Edition of Fownes' Manual of Chemistry). With 150 Wood Engravings, and coloured Plate of Spectra. Crown 8vo, 9s.

By the same Author.

Chemistry of Carbon - Compounds, or

Organic Chemistry (being Vol. II. of the Thirteenth Edition of Fownes' Manual of Chemistry). Edited by WM. A. TILDEN, D.Sc., F.R.S. With Engravings. Crown 8vo, 10s.

CHILDREN, DISEASES OF.

DAY.—A Manual of the Diseases of Children.

By WILLIAM H. DAY, M.D., Physician to the Samaritan Hospital for Women and Children. Second Edition. Crown 8vo, 12s. 6d.

ELLIS.—A Practical Manual of the Diseases

of Children. By EDWARD ELLIS, M.D., late Senior Physician to the Victoria Hospital for Sick Children. With a Formulary. Fifth Edition. Crown 8vo, 10s.

GOODHART.—The Student's Guide to Diseases

of Children. By JAMES F. GOODHART, M.D., F.R.C.P., Physician to Guy's Hospital; Physician to the Evelina Hospital for Sick Children. Second Edition. Fcap. 8vo. 10s. 6d.

SMITH.—On the Wasting Diseases of Infants

and Children. By EUSTACE SMITH, M.D., F.R.C.P., Physician to H.M. the King of the Belgians, and to the East London Hospital for Children. Fourth Edition. Post 8vo, 8s. 6d.

By the same Author.

A Practical Treatise on Disease in Chil-

dren. 8vo, 22s.

STEINER.—Compendium of Children's Dis-

eases; a Handbook for Practitioners and Students. By JOHANN STEINER, M.D. Translated by LAWSON TAIT, F.R.C.S., Surgeon to the Birmingham Hospital for Women, &c. 8vo, 12s. 6d.

11, NEW BURLINGTON STREET.

DENTISTRY.

GORGAS.—**Dental Medicine :** a Manual of Dental Materia Medica and Therapeutics, for Practitioners and Students. By FERDINAND J. S. GORGAS, A.M., M.D., D.D.S., Professor of Dentistry in the University of Maryland; Editor of "Harris's Principles and Practice of Dentistry," &c. Royal 8vo, 14s.

HARRIS.—**The Principles and Practice of** Dentistry; including Anatomy, Physiology, Pathology, Therapeutics, Dental Surgery, and Mechanism. By CHAPIN A. HARRIS, M.D., D.D.S. Eleventh Edition, revised and edited by FERDINAND J. S. GORGAS, A.M., M.D., D.D.S. With 750 Illustrations. 8vo, 31s. 6d.

SEWILL.—**The Student's Guide to Dental** Anatomy and Surgery. By HENRY E. SEWILL, M.R.C.S., L.D.S., late Dental Surgeon to the West London Hospital. Second Edition. With 78 Engravings. Fcap. 8vo, 5s. 6d.

STOCKEN.—**Elements of Dental Materia Medica** and Therapeutics, with Pharmacopœia. By JAMES STOCKEN, L.D.S.R.C.S., late Lecturer on Dental Materia Medica and Therapeutics and Dental Surgeon to the National Dental Hospital; assisted by THOMAS GADDES, L.D.S. Eng. and Edin. Third Edition. Fcap. 8vo, 7s. 6d.

TOMES (C. S.).—**Manual of Dental Anatomy,** Human and Comparative. By CHARLES S. TOMES, M.A., F.R.S. Second Edition. With 191 Engravings. Crown 8vo, 12s. 6d.

TOMES (J. and C. S.).—**A Manual of Dental** Surgery. By Sir JOHN TOMES, M.R.C.S., F.R.S., and CHARLES S. TOMES, M.A., M.R.C.S., F.R.S.; Lecturer on Anatomy and Physiology at the Dental Hospital of London. Third Edition. With many Engravings, Crown 8vo. [Preparing.]

EAR, DISEASES OF.

BURNETT.—**The Ear: its Anatomy, Physio-**logy, and Diseases. A Practical Treatise for the Use of Medical Students and Practitioners. By CHARLES H. BURNETT, M.D., Aural Surgeon to the Presbyterian Hospital, Philadelphia. Second Edition. With 107 Engravings. 8vo, 18s.

DALBY.—**On Diseases and Injuries of the Ear.** By SIR WILLIAM B. DALBY, F.R.C.S., Aural Surgeon to, and Lecturer on Aural Surgery at, St. George's Hospital. Third Edition. With Engravings. Crown 8vo. 7s. 6d.

11, NEW BURLINGTON STREET.

EAR, DISEASES OF—continued.

JONES.—**A Practical Treatise on Aural Surgery.** By H. MACNAUGHTON JONES, M.D., Professor of the Queen's University in Ireland, late Surgeon to the Cork Ophthalmic and Aural Hospital. Second Edition. With 63 Engravings. Crown 8vo, 8s. 6d.

By the same Author.

Atlas of the Diseases of the Membrana Tympani. In Coloured Plates, containing 59 Figures. With Explanatory Text. Crown 4to, 21s.

FORENSIC MEDICINE.

ABERCROMBIE.—**The Student's Guide to Medical Jurisprudence.** By JOHN ABERCROMBIE, M.D., F.R.C.P., Senior Assistant to, and Lecturer on Forensic Medicine at, Charing Cross Hospital. Fcap 8vo, 7s. 6d.

OGSTON.—**Lectures on Medical Jurisprudence.**

By FRANCIS OGSTON, M.D., late Professor of Medical Jurisprudence and Medical Logic in the University of Aberdeen. Edited by FRANCIS OGSTON, Jun., M.D., late Lecturer on Practical Toxicology in the University of Aberdeen. With 12 Plates. 8vo, 18s.

TAYLOR.—**The Principles and Practice of Medical Jurisprudence.** By ALFRED S. TAYLOR, M.D., F.R.S. Third Edition, revised by THOMAS STEVENSON, M.D., F.R.C.P., Lecturer on Chemistry and Medical Jurisprudence at Guy's Hospital; Examiner in Chemistry at the Royal College of Physicians; Official Analyst to the Home Office. With 188 Engravings. 2 Vols. 8vo, 31s. 6d.

By the same Author.

A Manual of Medical Jurisprudence.

Eleventh Edition, revised by THOMAS STEVENSON, M.D., F.R.C.P. With 56 Engravings. Crown 8vo, 14s.

ALSO,

On Poisons, in relation to Medical Jurisprudence and Medicine. Third Edition. With 104 Engravings. Crown 8vo, 16s.

TIDY AND WOODMAN.—**A Handy-Book of Forensic Medicine and Toxicology.** By C. MEYMOTT TIDY, M.B.; and W. BATHURST WOODMAN, M.D., F.R.C.P. With 8 Lithographic Plates and 116 Wood Engravings. 8vo, 31s. 6d.

HYGIENE.

PARKES.—A Manual of Practical Hygiene.

By EDMUND A. PARKES, M.D., F.R.S. Sixth Edition by F. DECHAUMONT, M.D., F.R.S., Professor of Military Hygiene in the Army Medical School. With 9 Plates and 103 Engravings. 8vo, 18s.

WILSON.—A Handbook of Hygiene and Sanitary Science.

By GEORGE WILSON, M.A., M.D., F.R.S.E., Medical Officer of Health for Mid Warwickshire. Sixth Edition. With Engravings. Crown 8vo, 10s. 6d.

MATERIA MEDICA AND THERAPEUTICS.

BINZ AND SPARKS.—The Elements of Therapeutics; a Clinical Guide to the Action of Medicines.

By C. BINZ, M.D., Professor of Pharmacology in the University of Bonn. Translated and Edited with Additions, in conformity with the British and American Pharmacopœias, by EDWARD I. SPARKS, M.A., M.B., F.R.C.P. Lond. Crown 8vo, 8s. 6d.

LESCHER.—Recent Materia Medica. Notes

on their Origin and Therapeutics. By F. HARWOOD LESCHER, F.C.S., Pereira Medallist. Second Edition. 8vo, 2s. 6d.

OWEN.—A Manual of Materia Medica; in-

corporating the Author's "Tables of Materia Medica." By ISAMBARD OWEN, M.D., F.R.C.P., Lecturer on Materia Medica and Therapeutics to St. George's Hospital. Second Edition. Crown 8vo, 6s. 6d.

ROYLE AND HARLEY.—A Manual of Materia

Medica and Therapeutics. By J. FORBES ROYLE, M.D., F.R.S., and JOHN HARLEY, M.D., F.R.C.P., Physician to, and Joint Lecturer on Clinical Medicine at, St. Thomas's Hospital. Sixth Edition, including addition and alterations in the B.P. 1885. With 139 Engravings. Crown 8vo, 15s.

THOROWGOOD.—The Student's Guide to

Materia Medica and Therapeutics. By JOHN C. THOROWGOOD, M.D., F.R.C.P., Lecturer on Materia Medica at the Middlesex Hospital. Second Edition. With Engravings. Fcap. 8vo, 7s.

WARING.—A Manual of Practical Therapeu-

tics. By EDWARD J. WARING, C.I.E., M.D., F.R.C.P. Fourth Edition, revised by the Author and DUDLEY W. BUXTON, M.D., M.R.C.P. Crown 8vo, 14s.

MEDICINE.

BARCLAY.—A Manual of Medical Diagnosis.

By A. WHYTE BARCLAY, M.D., F.R.C.P., late Physician to, and Lecturer on Medicine at, St. George's Hospital. Third Edition. Fcap. 8vo, 10s. 6d.

CHARTERIS.—The Student's Guide to the

Practice of Medicine. By MATTHEW CHARTERIS, M.D., Professor of Therapeutics and Materia Medica, University of Glasgow; Physician to the Royal Infirmary. With Engravings on Copper and Wood. Fourth Edition. Fcap. 8vo, 9s.

FAGGE.—The Principles and Practice of Medi-

cine. By the late C. HILTON FAGGE, M.D., F.R.C.P., Edited by P. H. PYE-SMITH, M.D., F.R.C.P., Physician to, and Lecturer on Medicine at, Guy's Hospital. 2 Vols. 8vo. Cloth, 36s.; half Persian, 42s.

FENWICK.—The Student's Guide to Medical

Diagnosis. By SAMUEL FENWICK, M.D., F.R.C.P., Physician to the London Hospital. Sixth Edition. With 114 Engravings. Fcap. 8vo, 7s.

By the same Author.

The Student's Outlines of Medical Treat-

ment. Second Edition. Fcap. 8vo, 7s.

FLINT.—Clinical Medicine : a Systematic Trea-

tise on the Diagnosis and Treatment of Disease. By AUSTIN FLINT, M.D., Professor of the Principles and Practice of Medicine, &c., in Bellevue Hospital Medical College. 8vo, 20s.

WARNER.—The Student's Guide to Clinical

Medicine and Case-Taking. By FRANCIS WARNER, M.D., F.R.C.P., Physician to the London Hospital. Second Edition. Fcap. 8vo, 5s.

WEST.—How to Examine the Chest: being a

Practical Guide for the Use of Students. By SAMUEL WEST, M.D., F.R.C.P., Physician to the City of London Hospital for Diseases of the Chest, &c. With 42 Engravings. Fcap. 8vo, 5s.

WHITTAKER.—Student's Primer on the Urine.

By J. TRAVIS WHITTAKER, M.D., Clinical Demonstrator at the Royal Infirmary, Glasgow. With Illustrations, and 16 Plates etched on Copper. Post 8vo, 4s. 6d.

MIDWIFERY.

BARNES.—*Lectures on Obstetric Operations*, including the Treatment of Hæmorrhage, and forming a Guide to the Management of Difficult Labour. By ROBERT BARNES, M.D., F.R.C.P., Consulting Obstetric Physician to St. George's Hospital. Fourth (and cheaper) Edition. With 121 Engravings. 8vo, 12s. 6d.

BURTON.—*Handbook of Midwifery for Midwives*. By JOHN E. BURTON, M.R.C.S., L.R.C.P., Surgeon to the Liverpool Hospital for Women. Second Edition. With Engravings. Fcap 8vo, 6s.

GALABIN.—*A Manual of Midwifery*. By Alfred LEWIS GALABIN, M.A., M.D., F.R.C.P., Obstetric Physician and Lecturer on Midwifery, &c., to Guy's Hospital, Examiner in Midwifery to the Conjoint Examining Board for England. With 227 Engravings. Crown 8vo, 15s.

RAMSBOTHAM.—*The Principles and Practice of Obstetric Medicine and Surgery*. By FRANCIS H. RAMSBOTHAM, M.D., formerly Obstetric Physician to the London Hospital. Fifth Edition. With 120 Plates, forming one thick handsome volume. 8vo, 22s.

REYNOLDS.—*Notes on Midwifery: specially designed to assist the Student in preparing for Examination*. By J. J. REYNOLDS, L.R.C.P., M.R.C.S. Second Edition. With 15 Engravings. Fcap. 8vo, 4s.

ROBERTS.—*The Student's Guide to the Practice of Midwifery*. By D. LLOYD ROBERTS, M.D., F.R.C.P., Lecturer on Clinical Midwifery and Diseases of Women at Owen's College, Physician to St. Mary's Hospital, Manchester. Third Edition. With 2 Coloured Plates and 127 Engravings. Fcap. 8vo, 7s. 6d.

SCHROEDER.—*A Manual of Midwifery; including the Pathology of Pregnancy and the Puerperal State*. By KARL SCHROEDER, M.D., Professor of Midwifery in the University of Erlangen. Translated by C. H. CARTER, M.D. With Engravings. 8vo, 12s. 6d.

SWAYNE.—*Obstetric Aphorisms for the Use of Students commencing Midwifery Practice*. By JOSEPH G. SWAYNE, M.D., Lecturer on Midwifery at the Bristol School of Medicine. Eighth Edition. With Engravings. Fcap. 8vo, 3s. 6d.

MICROSCOPY.

CARPENTER.—**The Microscope and its Revelations.** By WILLIAM B. CARPENTER, C.B., M.D., F.R.S. Sixth Edition. With about 600 Engravings. Crown 8vo, 16s.

LEE.—**The Microtometist's Vade-Mecum; a Handbook of the Methods of Microscopic Anatomy.** By ARTHUR BOLLES LEE. Crown 8vo, 8s. 6d.

MARSH.—**Microscopical Section-Cutting: a Practical Guide to the Preparation and Mounting of Sections for the Microscope.** By Dr. SYLVESTER MARSH. Second Edition. With 17 Engravings. Fcap. 8vo, 3s. 6d.

MARTIN.—**A Manual of Microscopic Mounting.** By J. H. MARTIN. Second Edition. With Plates and Wood Engravings. 8vo, 7s. 6d.

OPHTHALMOLOGY.

HARTRIDGE.—**The Refraction of the Eye.** By GUSTAVUS HARTRIDGE, F.R.C.S., Assistant Surgeon to the Royal Westminster Ophthalmic Hospital. Second Edition. With 94 Illustrations, Test Types, &c. Crown 8vo, 5s. 6d.

HIGGENS.—**Hints on Ophthalmic Out-Patient Practice.** By CHARLES HIGGENS, F.R.C.S., Ophthalmic Surgeon to, and Lecture on Ophthalmology at, Guy's Hospital. Third Edition. Fcap. 8vo, 3s.

MACNAMARA.—**A Manual of the Diseases of the Eye.** By CHARLES MACNAMARA, F.R.C.S., Surgeon to, and Lecturer on Surgery at, the Westminster Hospital. Fourth Edition. With 4 Coloured Plates and 66 Engravings. Crown 8vo, 10s. 6d.

NETTLESHIP.—**The Student's Guide to Diseases of the Eye.** By EDWARD NETTLESHIP, F.R.C.S., Ophthalmic Surgeon to, and Lecturer on Ophthalmic Surgery at, St. Thomas's Hospital. Third Edition. With 157 Engravings, and a Set of Coloured Papers illustrating Colour-blindness. Fcap. 8vo, 7s. 6d.

POLLOCK.—**The Normal and Pathological Histology of the Human Eye and Eyelids.** By C. FRED. POLLOCK, M.D., F.R.C.S.E., and F.R.S.E., Surgeon for Diseases of the Eye, Anderson's College Dispensary, Glasgow. With 100 Plates, containing 230 Original Drawings by the Author, Lithographed in black and colours. Crown 8vo, 15s.

OPHTHALMOLOGY—*continued.*

TOSSWILL.—Diseases and Injuries of the Eye and Eyelids. By LOUIS H. TOSSWILL, B.A., M.B. Cantab., M.R.C.S., Surgeon to the West of England Eye Infirmary, Exeter. Fcap. 8vo, 2s. 6d.

WOLFE.—On Diseases and Injuries of the Eye : a Course of Systematic and Clinical Lectures to Students and Medical Practitioners. By J. R. WOLFE, M.D., F.R.C.S.E., Senior Surgeon to the Glasgow Ophthalmic Institution, Lecturer on Ophthalmic Medicine and Surgery in Anderson's College. With 10 Coloured Plates, and 120 Wood Engravings, 8vo, 21s.

PATHOLOGY.

JONES AND SIEVEKING.—A Manual of Pathological Anatomy. By C. HANDFIELD JONES, M.B., F.R.S., and EDWARD H. SIEVEKING M.D., F.R.C.P. Second Edition. Edited, with considerable enlargement, by J. F. PAYNE, M.B., Assistant-Physician and Lecturer on General Pathology at St. Thomas's Hospital. With 195 Engravings. Crown 8vo, 16s.

LANCEREAUX.—Atlas of Pathological Anatomy. By Dr. LANCEREAUX. Translated by W. S. GREENFIELD, M.D., Professor of Pathology in the University of Edinburgh. With 70 Coloured Plates. Imperial 8vo, £5 5s.

SUTTON. — An Introduction to General Pathology. By JOHN BLAND SUTTON, F.R.C.S., Sir E. WILSON Lecturer on Pathology, R.C.S. ; Assistant Surgeon to, and Lecturer on Anatomy at, Middlesex Hospital. With 149 Engravings. 8vo, 14s.

VIRCHOW.—Post-Mortem Examinations : a Description and Explanation of the Method of Performing them, with especial reference to Medico-Legal Practice. By Professor RUDOLPH VIRCHOW, Berlin Charité Hospital. Translated by Dr. T. B. SMITH. Second Edition, with 4 Plates. Fcap. 8vo, 3s. 6d.

PSYCHOLOGY.

BUCKNILL AND TUKE.—A Manual of Psychological Medicine : containing the Lunacy Laws, Nosology, Ætiology, Statistics, Description, Diagnosis, Pathology, and Treatment of Insanity, with an Appendix of Cases. By JOHN C. BUCKNILL, M.D., F.R.S., and D. HACK TUKE, M.D., F.R.C.P. Fourth Edition with 12 Plates (30 Figures). 8vo, 25s.

PSYCHOLOGY—*continued.*

CLOUSTON.—**Clinical Lectures on Mental Diseases.** By THOMAS S. CLOUSTON, M.D., and F.R.C.P. Edin.; Lecturer on Mental Diseases in the University of Edinburgh. With 8 Plates (6 Coloured). Crown 8vo, 12s. 6d.

PHYSICS.

DRAPER.—**A Text Book of Medical Physics,** for the use of Students and Practitioners of Medicine By JOHN C. DRAPER, M.D., LL.D., Professor of Chemistry and Physics in the University of New York. With 377 Engravings. 8vo, 18s.

PHYSIOLOGY.

CARPENTER.—**Principles of Human Physiology.** By WILLIAM B. CARPENTER, C.B., M.D., F.R.S. Ninth Edition. Edited by Henry Power, M.B., F.R.C.S. With 3 Steel Plates and 377 Wood Engravings. 8vo, 31s. 6d.

DALTON.—**A Treatise on Human Physiology :** designed for the use of Students and Practitioners of Medicine. By JOHN C. DALTON, M.D., Professor of Physiology and Hygiene in the College of Physicians and Surgeons, New York. Seventh Edition. With 252 Engravings. Royal 8vo, 20s.

FREY.—**The Histology and Histo-Chemistry of Man.** A Treatise on the Elements of Composition and Structure of the Human Body. By HEINRICH FREY, Professor of Medicine in Zurich. Translated by ARTHUR E. BARKER, Assistant-Surgeon to the University College Hospital. With 608 Engravings. 8vo, 21s..

PYE-SMITH.—**Syllabus of a Course of Lectures on Physiology.** By PHILIP H. PYE-SMITH, B.A., M.D., F.R.C.P., Physician to Guy's Hospital. With Diagrams, and an Appendix of Notes and Tables. Crown 8vo, 5s.

SANDERSON.—**Handbook for the Physiological Laboratory :** containing an Exposition of the fundamental facts of the Science, with explicit Directions for their demonstration. By J. BURDON SANDERSON, M.D., F.R.S.; E. KLEIN, M.D., F.R.S.; MICHAEL FOSTER, M.D., F.R.S., and T. LAUDER BRUNTON, M.D., F.R.S. 2 Vols., with 123 Plates. 8vo, 24s.

YEO.—**A Manual of Physiology for the Use of Junior Students of Medicine.** By GERALD F. YEO, M.D., F.R.C.S., Professor of Physiology in King's College, London. With 301 Engravings. Crown 8vo, 14s.

SURGERY.

BELLAMY.—*The Student's Guide to Surgical Anatomy*; an Introduction to Operative Surgery. By EDWARD BELLAMY, F.R.C.S., and Member of the Board of Examiners; Surgeon to, and Lecturer on Anatomy at, Charing Cross Hospital. Third Edition. With 80 Engravings. Fcap. 8vo, 7s. 6d.

BRYANT.—*A Manual for the Practice of Surgery.* By THOMAS BRYANT, F.R.C.S., Surgeon to, and Lecturer on Surgery at, Guy's Hospital. Fourth Edition. With 750 Illustrations (many being coloured), and including 6 Chromo-Lithographic Plates. 2 Vols. Crown 8vo, 32s.

CLARK AND WAGSTAFFE. — *Outlines of Surgery and Surgical Pathology.* By F. LE GROS CLARK, F.R.C.S., F.R.S., Consulting Surgeon to St. Thomas's Hospital. Second Edition. Revised and expanded by the Author, assisted by W. W. WAGSTAFFE, F.R.C.S., Assistant Surgeon to St. Thomas's Hospital. 8vo, 10s. 6d.

DRUITT.—*The Surgeon's Vade-Mecum*; a Manual of Modern Surgery. By ROBERT DRUITT, F.R.C.S. Twelfth Edition. With many Engravings. Crown 8vo. [*Nearly ready.*]

FERGUSON.—*A System of Practical Surgery.* By Sir WILLIAM FERGUSON, Bart., F.R.C.S., F.R.S., late Surgeon and Professor of Clinical Surgery to King's College Hospital. With 463 Engravings. Fifth Edition. 8vo, 21s.

HEATH.—*A Manual of Minor Surgery and Bandaging*, for the use of House-Surgeons, Dressers, and Junior Practitioners. By CHRISTOPHER HEATH, F.R.C.S., Holme Professor of Clinical Surgery in University College and Surgeon to the Hospital. Eighth Edition. With 142 Engravings. Fcap. 8vo, 6s.

By the same Author.

A Course of Operative Surgery: with Twenty Plates (containing many figures) drawn from Nature by M. LÉVEILLÉ, and Coloured. Second Edition. Large 8vo, 30s.

ALSO,

The Student's Guide to Surgical Diagnosis. Second Edition. Fcap. 8vo, 6s. 6d.

11, *NEW BURLINGTON STREET.*

SURGERY—*continued.*

SOUTHAM.—**Regional Surgery :** including Surgical Diagnosis. A Manual for the use of Students. By FREDERICK A. SOUTHAM, M.A., M.B. Oxon, F.R.C.S., Assistant-Surgeon to the Royal Infirmary, and Assistant-Lecturer on Surgery in the Owen's College School of Medicine, Manchester.

Part I. The Head and Neck. Crown 8vo, 6s. 6d.

„ II. The Upper Extremity and Thorax. Crown 8vo, 7s. 6d.

„ III. The Abdomen and Lower Extremity. Crown 8vo, 7s.

TERMINOLOGY.

DUNGLISON.—**Medical Lexicon :** a Dictionary of Medical Science, containing a concise Explanation of its various Subjects and Terms, with Accentuation, Etymology, Synonyms, &c. By ROBERT DUNGLISON, M.D. New Edition, thoroughly revised by RICHARD J. DUNGLISON, M.D. Royal 8vo, 28s.

MAYNE.—**A Medical Vocabulary :** being an Explanation of all Terms and Phrases used in the various Departments of Medical Science and Practice, giving their Derivation, Meaning, Application, and Pronunciation. By ROBERT G. MAYNE, M.D., LL.D., and JOHN MAYNE, M.D., L.R.C.S.E. Fifth Edition. Crown 8vo, 10s. 6d.

WOMEN, DISEASES OF.

BARNES.—**A Clinical History of the Medical and Surgical Diseases of Women.** By ROBERT BARNES, M.D., F.R.C.P., Obstetric Physician to, and Lecturer on Diseases of Women, &c., at, St. George's Hospital. Second Edition. With 181 Engravings. 8vo, 28s.

COURTY.—**Practical Treatise on Diseases of the Uterus, Ovaries, and Fallopian Tubes.** By Professor COURTY, Montpellier. Translated from the Third Edition by his Pupil, AGNES M'LAREN, M.D., M.K.Q.C.P. With Preface by Dr. MATTHEWS DUNCAN. With 424 Engravings. 8vo, 24s.

DUNCAN.—**Clinical Lectures on the Diseases of Women.** By J. MATTHEWS DUNCAN, M.D., F.R.C.P., F.R.S.E., Obstetric Physician to St. Bartholomew's Hospital. Third Edition. 8vo.

[*In the press.*]

EMMET.—**The Principles and Practice of Gynæcology.** By THOMAS ADDIS EMMET, M.D., Surgeon to the Woman's Hospital of the State of New York. Third Edition. With 150 Engravings. Royal 8vo, 24s.

WOMEN, DISEASES OF—*continued.*

GALABIN.—**The Student's Guide to the Diseases of Women.** By ALFRED L. GALABIN, M.D., F.R.C.P., Obstetric Physician to, and Lecturer on Obstetric Medicine at, Guy's Hospital. Third Edition. With 78 Engravings. Fcap. 8vo, 7s. 6d.

REYNOLDS.—**Notes on Diseases of Women.** Specially designed to assist the Student in preparing for Examination. By J. J. REYNOLDS, L.R.C.P., M.R.C.S. Third Edition. Fcap. 8vo, 2s. 6d.

SAVAGE.—**The Surgery of the Female Pelvic Organs.** By HENRY SAVAGE, M.D., Lond., F.R.C.S., one of the Consulting Medical Officers of the Samaritan Hospital for Women. Fifth Edition, with 17 Lithographic Plates (15 Coloured), and 52 Woodcuts. Royal 4to, 35s.

WEST AND DUNCAN.—**Lectures on the Diseases of Women.** By CHARLES WEST, M.D., F.R.C.P. Fourth Edition. Revised and in part re-written by the Author, with numerous additions by J. MATTHEWS DUNCAN, M.D., F.R.C.P., F.R.S.E., Obstetric Physician to St. Bartholomew's Hospital. 8vo, 16s.

ZOOLOGY.

CHAUVEAU AND FLEMING.—**The Comparative Anatomy of the Domesticated Animals.** By A. CHAUVEAU, Professor at the Lyons Veterinary School; and GEORGE FLEMING, Veterinary Surgeon, Royal Engineers. With 450 Engravings. 8vo, 31s. 6d.

HUXLEY.—**Manual of the Anatomy of Invertebrated Animals.** By THOMAS H. HUXLEY, LL.D., F.R.S. With 156 Engravings. Post 8vo, 16s.

By the same Author.

Manual of the Anatomy of Vertebrated Animals. With 110 Engravings. Post 8vo, 12s.

WILSON.—**The Student's Guide to Zoology:** a Manual of the Principles of Zoological Science. By ANDREW WILSON, Lecturer on Natural History, Edinburgh. With Engravings. Fcap. 8vo, 6s. 6d.

11, NEW BURLINGTON STREET.

